

## BASIC COMPARISON THEOREMS FOR WEAK AND WEAKER MATRIX SPLITTINGS\*

ZBIGNIEW I. WOŹNICKI†

**Abstract.** The main goal of this paper is to present comparison theorems proven under natural conditions such as  $N_2 \geq N_1$  and  $M_1^{-1} \geq M_2^{-1}$  for weak and weaker splittings of  $A = M_1 - N_1 = M_2 - N_2$  in the cases when  $A^{-1} \geq 0$  and  $A^{-1} \leq 0$ .

**Key words.** Systems of linear equation, convergence conditions, comparison theorems, weak splittings, weaker splittings.

**AMS subject classifications.** 65C20, 65F10, 65F15

**1. Introduction.** A large class of iterative methods for solving system of linear equations of the form

$$Ax = b,$$

where  $A \in \mathbb{R}^{n \times n}$  is a nonsingular matrix and  $x, b \in \mathbb{R}^n$ , can be formulated by means of the splitting

$$(1.1) \quad A = M - N \quad \text{with} \quad M \text{ nonsingular},$$

and the approximate solution  $x^{(t+1)}$  is generated as follows

$$Mx^{(t+1)} = Nx^{(t)} + b, \quad t \geq 0,$$

or equivalently,

$$x^{(t+1)} = M^{-1}Nx^{(t)} + M^{-1}b, \quad t \geq 0,$$

where the starting vector  $x^{(0)}$  is given.

The above iterative method is convergent to the unique solution  $x = A^{-1}b$  for each  $x^{(0)}$  if and only if  $\rho(M^{-1}N) < 1$ , which means that the splitting of  $A = M - N$  is convergent. The convergence analysis of the above method is based on the spectral radius of the iteration matrix  $\rho(M^{-1}N)$ . As is well known, the smaller is  $\rho(M^{-1}N)$ , the faster is the convergence; see, e.g., [1].

The definitions of splittings, with progressively weaker conditions and consistent from the viewpoint of names, are collected in the following definition.

**DEFINITION 1.1.** Let  $M, N \in \mathbb{R}^{n \times n}$ . Then the decomposition  $A = M - N$  is called

(a) a *regular splitting* of  $A$  if  $M^{-1} \geq 0$  and  $N \geq 0$ ,

---

\*Received by the editors on 28 April 2000. Accepted for publication on 1 February 2001. Handling Editor: Miroslav Fiedler. This paper was presented at the Eleventh Haifa Matrix Theory Conference, Technion, Haifa, Israel, June 21-25, 1999.

†Institute of Atomic Energy 05-400 Otwock-Świerk, Poland (woznicki@hp2.cyf.gov.pl).

- (b) a *nonnegative splitting* of  $A$  if  $M^{-1} \geq 0$ ,  $M^{-1}N \geq 0$  and  $NM^{-1} \geq 0$ ,
- (c) a *weak nonnegative splitting* of  $A$  if  $M^{-1} \geq 0$  and either  $M^{-1}N \geq 0$  (the *first type*) or  $NM^{-1} \geq 0$  (the *second type*),
- (d) a *weak splitting* of  $A$  if  $M$  is nonsingular,  $M^{-1}N \geq 0$  and  $NM^{-1} \geq 0$ ,
- (e) a *weaker splitting* of  $A$  if  $M$  is nonsingular and either  $M^{-1}N \geq 0$  (the *first type*) or  $NM^{-1} \geq 0$  (the *second type*),
- (f) a *convergent splitting* of  $A$  if  $\varrho(M^{-1}N) = \varrho(NM^{-1}) < 1$ .

The splittings defined in the successive items extend progressively a class of splittings of  $A = M - N$  for which the matrices  $N$  and  $M^{-1}$  may lose the property of nonnegativity. Distinguishing both types of weak nonnegative and weaker splittings leads to further extensions allowing us to analyze cases when  $M^{-1}N$  may have negative entries if only  $NM^{-1}$  is a nonnegative matrix.

Different splittings were extensively analyzed by many authors, see, e.g., [2] and the references therein.

Conditions ensuring that a splitting of a nonsingular matrix  $A = M - N$  is convergent are unknown in a general case. As was pointed out in [2], the splittings defined in first three items of Definition 1.1 are convergent if and only if  $A^{-1} \geq 0$ , which means that both conditions  $A^{-1} \geq 0$  and  $\varrho(M^{-1}N) = \varrho(NM^{-1}) < 1$  are equivalent. We write this formally as the following lemma.

**LEMMA 1.2.** *Each weak nonnegative (as well as nonnegative and regular) splitting of  $A = M - N$  is convergent if and only if  $A^{-1} \geq 0$ . In other words, if  $A$  is not a monotone matrix, it is impossible to construct a convergent weak nonnegative splitting.*

In the case of weak and weaker splittings, the assumption  $A^{-1} \geq 0$  is not a sufficient condition in order to ensure the convergence of a given splitting of  $A$ ; it is also possible to construct a convergent weak or weaker splitting when  $A^{-1} \not\geq 0$ . Moreover, as can be shown by examples the conditions  $A^{-1}N \geq 0$  or  $NA^{-1} \geq 0$  may not ensure that a given splitting of  $A$  will be a weak or weaker splitting.

The properties of weaker splittings are summarized in the following theorem.

**THEOREM 1.3.** *Let  $A = M - N$  be a weaker splitting of  $A$ . If  $A^{-1} \geq 0$ , then*

1. *If  $M^{-1}N \geq 0$ , then  $A^{-1}N \geq M^{-1}N$  and if  $NM^{-1} \geq 0$ , then  $NA^{-1} \geq NM^{-1}$ .*
2.  $\varrho(M^{-1}N) = \frac{\varrho(A^{-1}N)}{1 + \varrho(A^{-1}N)} = \frac{\varrho(NA^{-1})}{1 + \varrho(NA^{-1})}$ .

Thus, we can conclude that for a convergent weaker splitting of a monotone matrix  $A$  there are three conditions  $M^{-1}N \geq 0$  (or  $NM^{-1} \geq 0$ ),  $A^{-1}N \geq 0$  (or  $NA^{-1} \geq 0$ ) and  $\varrho(M^{-1}N) = \varrho(NM^{-1}) < 1$ , and any two conditions imply the third.

The main goal of this paper is to present comparison theorems proven under natural conditions such as  $N_2 \geq N_1$  and  $M_1^{-1} \geq M_2^{-1}$  for weak and weaker splittings of  $A = M_1 - N_1 = M_2 - N_2$  in the cases when  $A^{-1} \geq 0$  and  $A^{-1} \leq 0$ .

**2. Comparison theorems.** When both convergent weaker splittings of a monotone matrix

$$(2.1) \quad A = M_1 - N_1 = M_2 - N_2$$

are of the same type, the inequality

$$(2.2) \quad N_2 \geq N_1$$

implies either

$$A^{-1}N_2 \geq A^{-1}N_1 \geq 0 \quad \text{or} \quad N_2A^{-1} \geq N_1A^{-1} \geq 0.$$

Hence, by the Perron-Frobenius theory of nonnegative matrices (see, e.g., [1]), we have  $\varrho(A^{-1}N_1) \leq \varrho(A^{-1}N_2)$  or  $\varrho(N_1A^{-1}) \leq \varrho(N_2A^{-1})$  and by Theorem 1.3 we can conclude the following result.

**THEOREM 2.1.** [2] *Let  $A = M_1 - N_1 = M_2 - N_2$  be two convergent weaker splittings of  $A$  of the same type, that is, either  $M_1^{-1}N_1 \geq 0$  and  $M_2^{-1}N_2 \geq 0$  or  $N_1M_1^{-1} \geq 0$  and  $N_2M_2^{-1} \geq 0$ , where  $A^{-1} \geq 0$ . If  $N_2 \geq N_1$ , then*

$$\varrho(M_1^{-1}N_1) \leq \varrho(M_2^{-1}N_2).$$

This theorem, proven originally by Varga [1] for regular splittings, carries over to the case when both weaker splittings are of the same type. As is pointed out in [3] when both splittings in (2.1) are of different types, the condition (2.2) may not hold.

In the case when  $A^{-1} \leq 0$ , then the inequality (2.2) implies either

$$0 \leq A^{-1}N_2 \leq A^{-1}N_1 \quad \text{or} \quad 0 \leq N_2A^{-1} \leq N_1A^{-1}.$$

Hence, one can deduce the following theorem.

**THEOREM 2.2.** *Let  $A = M_1 - N_1 = M_2 - N_2$  be two convergent weaker splittings of  $A$  of the same type, that is, either  $M_1^{-1}N_1 \geq 0$  and  $M_2^{-1}N_2 \geq 0$  or  $N_1M_1^{-1} \geq 0$  and  $N_2M_2^{-1} \geq 0$ , where  $A^{-1} \leq 0$ . If  $N_2 \geq N_1$ , then*

$$\varrho(M_1^{-1}N_1) \geq \varrho(M_2^{-1}N_2).$$

Similarly as in the case of  $A^{-1} \geq 0$ , it can be shown that when both splittings in (2.1) are of different types for  $A^{-1} \leq 0$ , condition (2.2) may not arise.

In the case of the weaker condition

$$(2.3) \quad M_1^{-1} \geq M_2^{-1}$$

the contrary behavior is observed. As is demonstrated on examples in [2], when both weak nonnegative splittings of a monotone matrix  $A$  are the same type, with  $M_1^{-1} \geq M_2^{-1}$  (or even  $M_1^{-1} > M_2^{-1}$ ) it may occur that  $\varrho(M_1^{-1}N_1) > \varrho(M_2^{-1}N_2)$ .

Let us assume that both convergent weaker splittings in (2.1) are of different types such that  $M_1^{-1}N_1 \geq 0$  and  $N_2M_2^{-1} \geq 0$ , and let  $v_1 \geq 0$  and  $y_2 \geq 0$  be the eigenvectors such that

$$(2.4) \quad v_1^T M_1^{-1} N_1 = \lambda_1 v_1^T$$

and

$$(2.5) \quad N_2 M_2^{-1} y_2 = \lambda_2 y_2,$$

where  $\lambda_1 = \varrho(M_1^{-1}N_1)$  and  $\lambda_2 = \varrho(M_2^{-1}N_2) = \varrho(N_2M_2^{-1})$ . Multiplying (2.4) on the right by  $A^{-1}y_2$  and (2.5) on the left by  $v_1^T A^{-1}$ , one obtains

$$v_1^T M_1^{-1} N_1 A^{-1} y_2 = \lambda_1 v_1^T A^{-1} y_2$$

and

$$v_1^T A^{-1} N_2 M_2^{-1} y_2 = \lambda_2 v_1^T A^{-1} y_2,$$

and after subtraction we obtain

$$v_1^T (A^{-1} N_2 M_2^{-1} - M_1^{-1} N_1 A^{-1}) y_2 = (\lambda_2 - \lambda_1) v_1^T A^{-1} y_2.$$

From (1.1) we have

$$M^{-1} = (A + N)^{-1} = A^{-1} (I + N A^{-1})^{-1} = (I + A^{-1} N)^{-1} A^{-1},$$

or

$$A^{-1} = M^{-1} + M^{-1} N A^{-1} = M^{-1} + A^{-1} N M^{-1}$$

which implies that

$$A^{-1} N_2 M_2^{-1} - M_1^{-1} N_1 A^{-1} = M_1^{-1} - M_2^{-1}.$$

Hence, one obtains

$$(2.6) \quad v_1^T (M_1^{-1} - M_2^{-1}) y_2 = (\lambda_2 - \lambda_1) v_1^T A^{-1} y_2.$$

Let us consider the following cases.

*Case I.* When  $A^{-1} > 0$ , then  $v_1^T A^{-1} y_2 > 0$ .

1. If  $M_1^{-1} > M_2^{-1}$ , then  $M_1^{-1} - M_2^{-1} > 0$  and  $v_1^T (M_1^{-1} - M_2^{-1}) y_2 > 0$ , hence  $\lambda_2 - \lambda_1 > 0$  and  $\lambda_2 > \lambda_1$ .

2. If  $M_1^{-1} \geq M_2^{-1}$ , then  $M_1^{-1} - M_2^{-1} \geq 0$  and

a) if  $v_1^T (M_1^{-1} - M_2^{-1}) y_2 > 0$ , hence  $\lambda_2 - \lambda_1 > 0$  and  $\lambda_2 > \lambda_1$ .

b) if  $v_1^T (M_1^{-1} - M_2^{-1}) y_2 = 0$ , hence  $\lambda_2 - \lambda_1 = 0$  and  $\lambda_2 = \lambda_1$ .

*Case II.* When  $A^{-1} \geq 0$ , then  $v_1^T A^{-1} y_2 \geq 0$ .

1. If  $v_1^T (M_1^{-1} - M_2^{-1}) y_2 > 0$ , then  $v_1^T A^{-1} y_2 > 0$ , hence  $\lambda_2 - \lambda_1 > 0$  and  $\lambda_2 > \lambda_1$ .

2. If  $v_1^T (M_1^{-1} - M_2^{-1}) y_2 = 0$ , then

a) for  $v_1^T A^{-1} y_2 > 0$ ,  $\lambda_2 - \lambda_1 = 0$  and  $\lambda_2 = \lambda_1$ .

b) for  $v_1^T A^{-1} y_2 = 0$ , the relation (2.6) is satisfied for arbitrary values of  $\lambda_1$  and  $\lambda_2$ .

The following examples of regular splittings illustrate the case II.2.b).

$$A = \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix} = M_1 - N_1 = M_2 - N_2, \text{ where}$$

$$M_1 = \begin{bmatrix} 6 & 0 \\ 0 & 5 \end{bmatrix}, \quad N_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad M_1^{-1} N_1 = \begin{bmatrix} \frac{1}{6} & 0 \\ 0 & 0 \end{bmatrix} \text{ and } v_1^T = [1 \quad 0],$$

$$M_2 = \begin{bmatrix} 6 & 0 \\ 0 & 7 \end{bmatrix}, \quad N_2 = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}, \quad M_2^{-1}N_2 = \begin{bmatrix} \frac{1}{6} & 0 \\ 0 & \frac{2}{7} \end{bmatrix} \quad \text{and} \quad y_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$\text{Evidently, } v_1^T(M_1^{-1} - M_2^{-1})y_2 = \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & \frac{2}{35} \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = 0$$

$$\text{and } v_1^T A^{-1}y_2 = \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{5} & 0 \\ 0 & \frac{1}{5} \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = 0.$$

However, a simple modification allows us to avoid this apparent difficulty appearing in the case II.2.b). Assuming a matrix  $B > 0$ , then instead the equations (2.4) and (2.5) the following equations may be taken in consideration

$$(2.7) \quad \tilde{v}_1^T(\varepsilon A^{-1}B + M_1^{-1}N_1) = \tilde{\lambda}_1 \tilde{v}_1^T$$

and

$$(2.8) \quad (\varepsilon BA^{-1} + N_2 M_2^{-1})\tilde{y}_2 = \tilde{\lambda}_2 \tilde{y}_2.$$

Since for  $\varepsilon > 0$  both matrices  $\varepsilon A^{-1}B + M_1^{-1}N_1$  and  $\varepsilon BA^{-1} + N_2 M_2^{-1}$  are irreducible, their eigenvalues  $\tilde{\lambda}_1$  and  $\tilde{\lambda}_2$  corresponding to spectral radii are strictly increasing functions of  $\varepsilon \geq 0$  [1], and  $\tilde{\lambda}_1 = \lambda_1$ ,  $\tilde{\lambda}_2 = \lambda_2$ ,  $\tilde{v}_1^T = v_1^T$  and  $\tilde{y}_2 = y_2$  with  $\varepsilon = 0$ . Multiplying (2.7) on the right by  $A^{-1}\tilde{y}_2$  and (2.8) on the left by  $\tilde{v}_1^T A^{-1}$  and proceeding similarly as with the derivation of (2.6), one obtains finally

$$(2.9) \quad \tilde{v}_1^T(M_1^{-1} - M_2^{-1})\tilde{y}_2 = (\tilde{\lambda}_2 - \tilde{\lambda}_1)\tilde{v}_1^T A^{-1}\tilde{y}_2.$$

Since for  $\varepsilon > 0$  both eigenvectors  $\tilde{v}_1$  and  $\tilde{y}_2$  are positive, it can be concluded that  $\tilde{v}_1^T(M_1^{-1} - M_2^{-1})\tilde{y}_2 > 0$  and  $\tilde{v}_1^T A^{-1}\tilde{y}_2 > 0$ , which implies that  $\tilde{\lambda}_2 - \tilde{\lambda}_1 > 0$  hence  $\tilde{\lambda}_2 > \tilde{\lambda}_1$ . Taking the limit for  $\varepsilon \rightarrow 0$ , it follows that  $\tilde{\lambda}_1 \rightarrow \lambda_1$  and  $\tilde{\lambda}_2 \rightarrow \lambda_2$  which allows us to conclude that  $\lambda_2 \geq \lambda_1$ .

In the case when both convergent weaker splittings are of different type but such that  $N_1 M_1^{-1} \geq 0$  and  $M_2^{-1}N_2 \geq 0$ , then instead of the equations (2.4) and (2.5) we can consider the equations

$$N_1 M_1^{-1}y_1 = \lambda_1 y_1 \quad \text{and} \quad v_2^T M_2^{-1}N_2 = \lambda_2 v_2^T$$

providing us the following equation

$$v_2^T(M_1^{-1} - M_2^{-1})y_1 = (\lambda_2 - \lambda_1)v_2^T A^{-1}y_1,$$

from which in a similar way we can conclude that  $\lambda_2 \geq \lambda_1$ .

Thus, from the above considerations we obtain the following result.

**THEOREM 2.3.** [2] *Let  $A = M_1 - N_1 = M_2 - N_2$  be two convergent weaker splittings of different types, that is, either  $M_1^{-1}N_1 \geq 0$  and  $N_2 M_2^{-1} \geq 0$  or  $N_1 M_1^{-1} \geq 0$  and  $M_2^{-1}N_2 \geq 0$ , where  $A^{-1} \geq 0$ . If  $M_1^{-1} \geq M_2^{-1}$ , then*

$$\varrho(M_1^{-1}N_1) \leq \varrho(M_2^{-1}N_2).$$

In particular, if  $A^{-1} > 0$  and  $M_1^{-1} > M_2^{-1}$ , then

$$\varrho(M_1^{-1}N_1) < \varrho(M_2^{-1}N_2).$$

Assuming now that both convergent weaker splittings of different types in (2.1) are derived from a non-monotone matrix  $A$ . Referring back to (2.6) the following cases can be analyzed.

*Case III.* When  $A^{-1} < 0$ , then  $v_1^T A^{-1} y_2 < 0$ .

1. If  $M_1^{-1} > M_2^{-1}$ , then  $M_1^{-1} - M_2^{-1} > 0$  and  $v_1^T (M_1^{-1} - M_2^{-1}) y_2 > 0$ , hence  $\lambda_2 - \lambda_1 < 0$  and  $\lambda_2 < \lambda_1$ .

2. If  $M_1^{-1} \geq M_2^{-1}$ , then  $M_1^{-1} - M_2^{-1} \geq 0$  and

a) if  $v_1^T (M_1^{-1} - M_2^{-1}) y_2 > 0$ , hence  $\lambda_2 - \lambda_1 < 0$  and  $\lambda_2 < \lambda_1$ .

b) if  $v_1^T (M_1^{-1} - M_2^{-1}) y_2 = 0$ , hence  $\lambda_2 - \lambda_1 = 0$  and  $\lambda_2 = \lambda_1$ .

*Case IV.* When  $A^{-1} \leq 0$ , then  $v_1^T A^{-1} y_2 \leq 0$ .

1. If  $v_1^T (M_1^{-1} - M_2^{-1}) y_2 > 0$ , then  $v_1^T A^{-1} y_2 < 0$ , hence  $\lambda_2 - \lambda_1 < 0$  and  $\lambda_2 < \lambda_1$ .

2. If  $v_1^T (M_1^{-1} - M_2^{-1}) y_2 = 0$ , then

a) for  $v_1^T A^{-1} y_2 < 0$ ,  $\lambda_2 - \lambda_1 = 0$  and  $\lambda_2 = \lambda_1$ .

b) for  $v_1^T A^{-1} y_2 = 0$ , the relation (2.6) is satisfied for arbitrary values of  $\lambda_1$  and  $\lambda_2$ .

The following examples of weaker splittings illustrate the case IV.2.b).

$$A = \begin{bmatrix} -5 & 0 \\ 0 & -5 \end{bmatrix} = M_1 - N_1 = M_2 - N_2 \text{ where}$$

$$M_1 = \begin{bmatrix} -6 & 0 \\ 0 & -7 \end{bmatrix}, N_1 = \begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix}, M_1^{-1} N_1 = \begin{bmatrix} \frac{1}{6} & 0 \\ 0 & \frac{2}{7} \end{bmatrix} \text{ and } v_1^T = \begin{bmatrix} 0 & 1 \end{bmatrix},$$

$$M_2 = \begin{bmatrix} -6 & 0 \\ 0 & -5 \end{bmatrix}, N_2 = \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix}, M_2^{-1} N_2 = \begin{bmatrix} \frac{1}{6} & 0 \\ 0 & 0 \end{bmatrix} \text{ and } y_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

$$\text{Evidently, } v_1^T (M_1^{-1} - M_2^{-1}) y_2 = \begin{bmatrix} 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & \frac{2}{35} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 0$$

$$\text{and } v_1^T A^{-1} y_2 = \begin{bmatrix} 0 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{5} & 0 \\ 0 & -\frac{1}{5} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 0.$$

Assuming now a matrix  $B < 0$ , and repeating the same procedure as in the case of the case II.2.b), one can obtain again (2.9) from which, taking the limit for  $\varepsilon \rightarrow 0$ , we can conclude that  $\lambda_2 \leq \lambda_1$  for the case IV.2.b). Hence, the following theorem holds.

**THEOREM 2.4.** *Let  $A = M_1 - N_1 = M_2 - N_2$  be two convergent weaker splittings of different types, that is, either  $M_1^{-1} N_1 \geq 0$  and  $N_2 M_2^{-1} \geq 0$  or  $N_1 M_1^{-1} \geq 0$  and  $M_2^{-1} N_2 \geq 0$ , where  $A^{-1} \leq 0$ . If  $M_1^{-1} \geq M_2^{-1}$ , then*

$$\varrho(M_1^{-1} N_1) \geq \varrho(M_2^{-1} N_2).$$

In particular, if  $A^{-1} < 0$  and  $M_1^{-1} > M_2^{-1}$ , then

$$\varrho(M_1^{-1} N_1) > \varrho(M_2^{-1} N_2).$$

Thus, we see that for the conditions (2.2) and (2.3) passing from the assumption  $A^{-1} \geq 0$  to the assumption  $A^{-1} \leq 0$  implies the change of the inequality sign in the inequalities for spectral radii.

Finally, it is evident that the following corollary holds.

**COROLLARY 2.5.** *Let  $A = M_1 - N_1 = M_2 - N_2$  be two convergent weak splittings or one of them is weak and the second is weaker, then Theorems 2.1, 2.2, 2.3, and 2.4 hold.*

#### REFERENCES

- [1] R.S.Varga. *Matrix iterative analysis*. Prentice Hall, Englewood Cliffs, N.J., 1962. Second edition, revised and expanded, Springer, Berlin, Heidelberg, New York, 2000.
- [2] Z.I.Woźnicki. Nonnegative splitting theory. *Japan J. Industr. Appl. Math.*, 11:289–342, 1994.
- [3] H.A.Jedrzejec and Z.I.Woźnicki. On properties of some matrix splittings. *Electron. J. Linear Algebra*, 8:47–52, 2001.