

## CONSTRUCTIONS OF TRACE ZERO SYMMETRIC STOCHASTIC MATRICES FOR THE INVERSE EIGENVALUE PROBLEM\*

ROBERT REAMS†

**Abstract.** In the special case of where the spectrum  $\sigma = \{\lambda_1, \lambda_2, \lambda_3, 0, 0, \dots, 0\}$  has at most three nonzero eigenvalues  $\lambda_1, \lambda_2, \lambda_3$  with  $\lambda_1 \geq 0 \geq \lambda_2 \geq \lambda_3$ , and  $\lambda_1 + \lambda_2 + \lambda_3 = 0$ , the inverse eigenvalue problem for symmetric stochastic  $n \times n$  matrices is solved. Constructions are provided for the appropriate matrices where they are readily available. It is shown that when  $n$  is odd it is not possible to realize the spectrum  $\sigma$  with an  $n \times n$  symmetric stochastic matrix when  $\lambda_3 \neq 0$  and  $\frac{3}{2n-3} > \frac{\lambda_2}{\lambda_3} \geq 0$ , and it is shown that this bound is best possible.

**Key words.** Inverse eigenvalue problem, Symmetric stochastic matrix, Symmetric nonnegative matrix, Distance matrix.

**AMS subject classifications.** 15A18, 15A48, 15A51, 15A57

**1. Introduction.** Let  $e_1, \dots, e_n$  denote the standard basis in  $\mathbb{R}^n$ , so  $e_i$  denotes the vector with a 1 in the  $i$ th position and zeroes elsewhere. We will denote by  $e$  the vector of all ones, i.e.  $e = [1, 1, \dots, 1]^T \in \mathbb{R}^n$ . A matrix  $A \in \mathbb{R}^{n \times n}$  is said to be stochastic when all of its entries are nonnegative and all its row sums are equal to 1, i.e.  $A$  has right eigenvector  $e$  corresponding to the eigenvalue 1. We will be concerned with symmetric stochastic matrices, so that these matrices are in fact doubly stochastic. Also, the eigenvalues will necessarily be real. If  $A \in \mathbb{R}^{n \times n}$  is nonnegative, has eigenvalue  $\lambda_1 > 0$  corresponding to the right eigenvector  $e$  then  $\frac{1}{\lambda_1}A$  is stochastic, and for convenience we will state our results in the form, for example, of a matrix  $A$  having eigenvector  $e$  corresponding to  $1 + \epsilon$ , where the spectrum  $\sigma = \{1 + \epsilon, -1, -\epsilon, 0, 0, \dots, 0\}$ , with  $0 \leq \epsilon \leq 1$ . We will say that  $\sigma = \{\lambda_1, \dots, \lambda_n\} \subset \mathbb{R}$  is realized as the spectrum of a matrix  $A$  in the event that the  $n \times n$  matrix  $A$  has eigenvalues  $\lambda_1, \dots, \lambda_n$ .

The nonnegative inverse eigenvalue problem is to find necessary and sufficient conditions that the elements of the set  $\sigma = \{\lambda_1, \dots, \lambda_n\} \subset \mathbb{C}$  are the eigenvalues of a matrix with nonnegative entries. This problem is currently unsolved except in special cases [1], [7], [8], [9], [10]. The restriction of this problem to symmetric nonnegative matrices for which the eigenvalues  $\lambda_1, \dots, \lambda_n$  satisfy  $\lambda_1 \geq 0 \geq \lambda_2 \geq \dots \geq \lambda_n$  is solved in [3], where it is shown that the only necessary and sufficient condition is that  $\sum_{i=1}^n \lambda_i \geq 0$ . Distance matrices are necessarily symmetric, nonnegative, have trace zero, and must have  $\lambda_1 \geq 0 \geq \lambda_2 \geq \dots \geq \lambda_n$ , although these are not sufficient conditions to be a distance matrix. We conjectured in [6], after solving numerous special cases including  $n = 2, 3, 4, 5, 6$ , that the only necessary and sufficient conditions for the existence of a distance matrix with a given  $\sigma$  is that  $\lambda_1 \geq 0 \geq \lambda_2 \geq \dots \geq \lambda_n$  and  $\sum_{i=1}^n \lambda_i = 0$ . Distance matrices with eigenvector  $e$  were previously studied in [5],

\* Received by the editors on 13 October 2001. Final manuscript accepted for publication on 4 September 2002. Handling Editor: Tom Laffey.

† Department of Mathematics, National University of Ireland Galway, Galway, Ireland (robert.reams@nuigalway.ie).

although not in the context of these matrices eigenvalues. Bounds on the eigenvalues of symmetric stochastic matrices are given in [2] and [4], although they do not provide a restriction on the eigenvalues for the matrices under consideration here.

In Section 2 we provide an explicit construction of an  $n \times n$  symmetric stochastic matrix which realizes the spectrum  $\{2, -1, -1, 0, 0, \dots, 0\}$ , followed by showing that it is not possible to realize the spectrum  $\{1, -1, 0, 0, \dots, 0\}$  with a symmetric stochastic matrix when  $n$  is odd, although it is possible to realize this spectrum when  $n$  is even. In Section 3 we provide explicit constructions of symmetric stochastic matrices to realize  $\{1 + \epsilon, -1, -\epsilon, 0, 0, \dots, 0\}$ , for  $1 \geq \epsilon \geq 0$ , when  $n$  is even. We then show that it is not possible to realize this spectrum with a symmetric stochastic matrix when  $\frac{3}{2n-3} > \epsilon \geq 0$ , and  $n$  is odd. Although we can realize this spectrum with a symmetric stochastic matrix when  $1 \geq \epsilon \geq \frac{3}{2n-3}$ , and  $n$  is odd. In the latter case we do not provide an explicit construction, instead making use of the Intermediate Value Theorem in several variables.

**2. Freedom and restrictions on spectra.** Lemma 2.1 will be used to establish that the matrix under consideration is nonnegative.

**LEMMA 2.1.** *Let  $A = (a_{ij}) \in \mathbb{R}^{n \times n}$  be a symmetric matrix with eigenvalues  $\lambda_1, \dots, \lambda_n$  which satisfy  $\lambda_1 \geq 0 \geq \lambda_2 \geq \dots \geq \lambda_n$ . Suppose that  $A$  has eigenvector  $e$  corresponding to  $\lambda_1$ , and that  $A$  has all diagonal entries equal to zero. Then  $A$  is a nonnegative matrix.*

*Proof.* Write  $A = \lambda_1 \frac{ee^T}{n} + \lambda_2 u_2 u_2^T + \dots + \lambda_n u_n u_n^T$ , where  $u_2, \dots, u_n$  are unit eigenvectors corresponding to  $\lambda_2, \dots, \lambda_n$ , respectively. Then  $-2a_{ij} = (e_i - e_j)^T A (e_i - e_j) = \lambda_2 ((e_i - e_j)^T u_2)^2 + \dots + \lambda_n ((e_i - e_j)^T u_n)^2 \leq 0$ , for all  $i, j$ ,  $1 \leq i, j \leq n$ .  $\square$

Our next two theorems give some foretaste of what is and isn't possible with the inverse eigenvalue problem for symmetric nonnegative matrices. Note first that a  $3 \times 3$  symmetric nonnegative matrix with eigenvector  $e$  and of trace zero must be a nonnegative scalar multiple of  $A = ee^T - I$ , which has spectrum  $\sigma = \{2, -1, -1\}$ .

**THEOREM 2.2.** *Let  $\sigma = \{2, -1, -1, 0, 0, \dots, 0\}$ . Then  $\sigma$  can be realized as the spectrum of a symmetric nonnegative matrix  $A \in \mathbb{R}^{n \times n}$  with eigenvector  $e$  corresponding to 2, for  $n \geq 3$ .*

*Proof.* Let  $u = [u_1, \dots, u_n]^T, v = [v_1, \dots, v_n]^T \in \mathbb{R}^n$  be given by  $u_j = \sqrt{\frac{2}{n}} \cos \theta_j$ ,  $v_j = \sqrt{\frac{2}{n}} \sin \theta_j$ , where  $\theta_j = \frac{2\pi}{n} j$ , for  $1 \leq j \leq n$ . Then  $A = 2 \frac{ee^T}{n} - uu^T - vv^T$  is symmetric and has zero diagonal entries by construction. Since the roots of  $x^n - 1$  are  $e^{\frac{2\pi i}{n} j}$ ,  $1 \leq j \leq n$ , then (coefficient of  $x^{n-1}$ )  $= 0 = \sum_{j=1}^n -e^{\frac{2\pi i}{n} j}$  and (coefficient of  $x^{n-2}$ )  $= 0 = \sum_{j,k=1, j \neq k}^n e^{\frac{2\pi i}{n} j} e^{\frac{2\pi i}{n} k} = (\sum_{j=1}^n e^{\frac{2\pi i}{n} j})^2 - \sum_{j=1}^n e^{\frac{2\pi i}{n} 2j}$ . It follows that  $\sum_{j=1}^n \cos \theta_j = \sum_{j=1}^n \sin \theta_j = \sum_{j=1}^n \sin 2\theta_j = 0$ , which tells us that  $u^T e = v^T e = u^T v = 0$ , respectively. It also follows that  $\sum_{j=1}^n \cos 2\theta_j = 0$ , so that  $\sum_{j=1}^n \cos^2 \theta_j - \sum_{j=1}^n \sin^2 \theta_j = 0$ . We also know that  $\sum_{j=1}^n \cos^2 \theta_j + \sum_{j=1}^n \sin^2 \theta_j = n$ , so we can conclude that  $\sum_{j=1}^n \cos^2 \theta_j = \sum_{j=1}^n \sin^2 \theta_j = \frac{n}{2}$ . This tells us that  $u$  and  $v$  are unit vectors. Thus  $A$  has spectrum  $\sigma$  and is nonnegative from Lemma 2.1.  $\square$

**THEOREM 2.3.** *Let  $\lambda_1 > 0$  and  $\sigma = \{\lambda_1, -\lambda_1, 0, \dots, 0\}$ . Then  $\sigma$  cannot be realized as the spectrum of an  $n \times n$  symmetric nonnegative matrix with eigenvector*

$e$  corresponding to  $\lambda_1$ , when  $n$  is odd.

*Proof.* Suppose  $A$  is a matrix of the given form that realizes  $\sigma$ . Write  $A = \lambda_1 \frac{ee^T}{n} - \lambda_1 uu^T$ , where  $u = [u_1, \dots, u_n]^T \in \mathbb{R}^n$ ,  $\|u\| = 1$  and  $u^T e = 0$ .  $A$  has trace zero and is nonnegative by hypothesis so all diagonal entries are zero, and thus  $0 = \frac{\lambda_1}{n} - \lambda_1 u_i^2$ , for  $1 \leq i \leq n$ . But then  $u_i = \frac{\pm 1}{\sqrt{n}}$  and  $0 = \sum_{i=1}^n u_i = \sum_{i=1}^n \frac{\pm 1}{\sqrt{n}}$ , which is not possible when  $n$  is odd.  $\square$

REMARK 2.4. The same  $\sigma$  of Theorem 2.3 can be realized by an  $n \times n$  symmetric nonnegative matrix with eigenvector  $e$  corresponding to  $\lambda_1$  when  $n$  is even, by taking in the proof the unit vector  $u = \frac{1}{\sqrt{n}}[1, 1, \dots, 1, -1, -1, \dots, -1]^T \in \mathbb{R}^n$ .

**3. At most three nonzero eigenvalues.** The idea in Remark 2.4 can be extended to the case where  $n$  is a multiple of 4 for the case of at most three nonzero eigenvalues.

THEOREM 3.1. Let  $\sigma = \{1 + \epsilon, -1, -\epsilon, 0, 0, \dots, 0\}$ , where  $1 \geq \epsilon \geq 0$ . Then  $\sigma$  can be realized by a symmetric nonnegative matrix  $A \in \mathbb{R}^{n \times n}$  with eigenvector  $e$  corresponding to  $1 + \epsilon$ , when  $n = 4m$  for some  $m$ .

*Proof.* Let  $u, v \in \mathbb{R}^n$  be given by

$$u = \frac{1}{\sqrt{n}}[1, 1, \dots, 1, 1, 1, \dots, 1, -1, -1, \dots, -1, -1, -1, \dots, -1]^T,$$

$$v = \frac{1}{\sqrt{n}}[1, 1, \dots, 1, -1, -1, \dots, -1, 1, 1, \dots, 1, -1, -1, \dots, -1]^T.$$

Then  $\frac{e}{\sqrt{n}}$ ,  $u$  and  $v$  are orthonormal. Let  $A = \frac{(1+\epsilon)}{n}ee^T - uu^T - \epsilon vv^T$ , and note that all the diagonal entries of  $A$  are zero.  $\square$

However, the way we deal with the remaining cases for  $n$  even is somewhat more complicated.

THEOREM 3.2. Let  $\sigma = \{1 + \epsilon, -1, -\epsilon, 0, 0, \dots, 0\}$ , where  $1 \geq \epsilon \geq 0$ . Then  $\sigma$  can be realized by a symmetric nonnegative matrix  $A \in \mathbb{R}^{n \times n}$  with eigenvector  $e$  corresponding to  $1 + \epsilon$ , when  $n = 4m + 2$  for some  $m$ .

*Proof.* Let  $u = [u_1, \dots, u_n]^T$ ,  $v = [v_1, \dots, v_n]^T \in \mathbb{R}^n$ , and  $A = (1 + \epsilon)\frac{ee^T}{n} - uu^T - \epsilon vv^T$ . We will require that  $0 = \frac{1+\epsilon}{n} - u_i^2 - \epsilon v_i^2$ , for  $1 \leq i \leq n$ , so that  $A$  has all zero diagonal entries. Let  $u_i = \sqrt{\frac{1+\epsilon}{n}} \cos \theta_i$  and  $v_i = \sqrt{\frac{1+\epsilon}{n\epsilon}} \sin \theta_i$ , where the  $\theta_i$ 's are chosen below. Our  $\theta_i$ 's must satisfy  $\sum_{i=1}^n \cos \theta_i = \sum_{i=1}^n \sin \theta_i = \sum_{i=1}^n \cos \theta_i \sin \theta_i = 0$ . So that  $u$  and  $v$  are unit vectors we must have  $\sum_{i=1}^n \cos^2 \theta_i = \frac{n}{1+\epsilon}$  and  $\sum_{i=1}^n \sin^2 \theta_i = \frac{n\epsilon}{1+\epsilon}$ , which will be achieved if  $\sum_{i=1}^n \cos^2 \theta_i - \sum_{i=1}^n \sin^2 \theta_i = \frac{n(1-\epsilon)}{1+\epsilon}$ , and  $\sum_{i=1}^n \cos^2 \theta_i + \sum_{i=1}^n \sin^2 \theta_i = n$ . So for any given  $\epsilon \in [0, 1]$  we also require for our choice of  $\theta_i$ 's that  $\sum_{i=1}^n \cos 2\theta_i = \frac{n(1-\epsilon)}{1+\epsilon}$ .

Let  $0 \leq \delta < \frac{2\pi}{n}$ , and choose the angles  $\theta_i$ ,  $1 \leq i \leq n$ , around the origin in the plane.

Let  $\theta_i$  for  $1 \leq i \leq m$  be, respectively, the 1st quadrant angles

$$\frac{2\pi}{n} - \delta, 2(\frac{2\pi}{n} - \delta), \dots, (m-1)(\frac{2\pi}{n} - \delta), m(\frac{2\pi}{n} - \delta).$$

Let  $\theta_i$  for  $m+1 \leq i \leq 2m$  be, respectively, the 2nd quadrant angles

$$(m+1)\frac{2\pi}{n} + m\delta, (m+2)\frac{2\pi}{n} + (m-1)\delta, \dots, (2m-1)\frac{2\pi}{n} + 2\delta, 2m\frac{2\pi}{n} + \delta.$$

Let  $\theta_{2m+1} = \pi$ .

Let  $\theta_i$  for  $2m+2 \leq i \leq 3m+1$  be, respectively, the 3rd quadrant angles

$$-2m\frac{2\pi}{n} - \delta, -(2m-1)\frac{2\pi}{n} - 2\delta, \dots, -(m+2)\frac{2\pi}{n} - (m-1)\delta, -(m+1)\frac{2\pi}{n} - m\delta.$$

Let  $\theta_i$  for  $3m+2 \leq i \leq 4m+1$  be, respectively, the 4th quadrant angles  $-m(\frac{2\pi}{n}-\delta)$ ,  $-(m-1)(\frac{2\pi}{n}-\delta)$ ,  $\dots$ ,  $-2(\frac{2\pi}{n}-\delta)$ ,  $-(\frac{2\pi}{n}-\delta)$ .

Let  $\theta_{4m+2} = 0$ .

For each  $\theta_i$  there is a  $\theta_i + \pi$  in the above list, by pairing off 1st quadrant angles with appropriate 3rd quadrant angles, and pairing 2nd quadrant angles with appropriate 4th quadrant angles. Therefore since  $\cos(\theta_i + \pi) = -\cos \theta_i$  we conclude that  $\sum_{i=1}^n \cos \theta_i = 0$ .

For each  $\theta_i \neq 0$  and  $\theta_i \neq \pi$  in the above list we can pair off any  $\theta_i$  with a corresponding  $-\theta_i$  and conclude that  $\sum_{i=1}^n \sin \theta_i = 0$ .

Similarly,  $\sum_{i=1}^n \sin 2\theta_i = 0$ .

Pairing off each  $\theta_i$  with a  $\theta_i + \pi$  in the same way as before, and since  $\cos 2\theta_i = \cos(2\theta_i + 2\pi)$ , we must have

$$\sum_{i=1}^n \cos 2\theta_i = 2 + \sum_{\text{1st, 2nd, 3rd, 4th quadrant}} \cos 2\theta_i = 2 + 2 \sum_{\text{1st, 4th quadrant}} \cos 2\theta_i.$$

Because we can pair off each  $\theta_i$  in the 1st quadrant with each  $-\theta_i$  in the 4th quadrant  $\sum_{i=1}^n \cos 2\theta_i = 2 + 4 \sum_{\text{1st quadrant}} \cos 2\theta_i$ . Next using the trigonometric formula

$$\sum_{j=0}^n \cos j\alpha = \frac{\sin(\frac{n+1}{2}\alpha) \cos(\frac{n}{2}\alpha)}{\sin \frac{\alpha}{2}}$$

(found in [11], for example), where  $\alpha = 2(\frac{2\pi}{n}-\delta)$  and some simplification we obtain that  $\sum_{i=1}^n \cos 2\theta_i = 2 \frac{\sin((2m+1)\delta)}{\sin(\frac{\pi}{2m+1}-\delta)}$ . Let  $f(\delta) = 2 \frac{\sin((2m+1)\delta)}{\sin(\frac{\pi}{2m+1}-\delta)}$ , then  $f(0) = 0$  and  $\lim_{\delta \rightarrow \frac{2\pi}{n}} f(\delta) = n$ , and notice that  $f$  is continuous on the interval  $[0, \frac{2\pi}{n})$ . Also, since  $0 \leq \epsilon \leq 1$  we have that  $0 \leq \frac{n(1-\epsilon)}{1+\epsilon} \leq n$ , but then by the Intermediate Value Theorem for each  $\epsilon \in (0, 1]$  there is a  $\delta \in [0, \frac{2\pi}{n})$  such that  $f(\delta) = \frac{n(1-\epsilon)}{1+\epsilon}$ . The case  $\epsilon = 0$  and  $\delta = \frac{2\pi}{n}$  is covered by the remark after Theorem 2.3.  $\square$

In order to deal with the case where  $n$  is odd we will improve on Theorem 2.3.

**THEOREM 3.3.** Let  $\sigma = \{1+\epsilon, -1, -\epsilon, 0, 0, \dots, 0\}$ , where  $\frac{3}{2n-3} > \epsilon \geq 0$ , and  $n \geq 3$ . Then  $\sigma$  cannot be realized by a symmetric nonnegative matrix  $A \in \mathbb{R}^{n \times n}$  with eigenvector  $e$  corresponding to  $1+\epsilon$ , when  $n = 2m+1$  for some  $m$ .

*Proof.*  $\sigma$  is realizable when  $\epsilon = 1$  from Theorem 2.2. We wish to determine the minimum value of  $\epsilon$  for which it is possible to construct a matrix of the desired form. Using the same notation as in the proof of Theorem 3.2 we wish to determine the minimum value of  $\epsilon$  as a function of  $\theta_1, \dots, \theta_n$  subject to the three constraints  $\sum_{i=1}^n \cos \theta_i = \sum_{i=1}^n \sin \theta_i = 0$  and  $\sum_{i=1}^n \cos \theta_i \sin \theta_i = 0$  (we know from Theorem 2.3 that  $\epsilon > 0$ ). Observe that finding the minimum  $\epsilon$  is equivalent to finding the maximum value of the function  $\frac{n}{1+\epsilon} = F(\theta_1, \dots, \theta_n) = \sum_{i=1}^n \cos^2 \theta_i$  subject to the three constraints. For the moment let us determine the maximum value of  $F$  subject to the one constraint  $\sum_{i=1}^n \cos \theta_i = 0$ . Let  $\lambda$  denote the Lagrange multiplier so that  $-2 \cos \theta_i \sin \theta_i - \lambda \sin \theta_i = 0$ , i.e.  $\sin \theta_i (2 \cos \theta_i + \lambda) = 0$ , for each  $i$ ,  $1 \leq i \leq n$ . Then

for each  $i$  we have  $\sin \theta_i = 0$  or else  $\cos \theta_i = -\lambda/2$ . We cannot have all  $\theta_i$ 's equal to 0 or  $\pi$ , since then we would not have  $\sum_{i=1}^n \cos \theta_i = 0$ , because  $n$  is odd. Suppose now that  $k$  of the  $\cos \theta_i$ 's are equal, but not equal to  $\pm 1$ , then  $k \cos \theta_i = \pm 1 \pm 1 \dots$ . Then in order to maximize  $F$  we must have as many  $\pm 1$ 's as possible, thus we must have  $k = 2$  with  $\cos \theta_p = \cos \theta_q$  (say) and the remaining  $\theta_i$ 's all either 0 or  $\pi$ . Then  $2 \cos \theta_i = 1$  or  $-1$ , and there are two possibilities:

Case 1:  $\cos \theta_p = \cos \theta_q = \frac{1}{2}$  with  $m - 1$  of the remaining  $\cos \theta_i$ 's equal to 1 and the other  $m$  of the  $\cos \theta_i$ 's equal to  $-1$ .

Case 2:  $\cos \theta_p = \cos \theta_q = -\frac{1}{2}$  with  $m - 1$  of the remaining  $\cos \theta_i$ 's equal to  $-1$  and the other  $m$  of the  $\cos \theta_i$ 's equal to 1.

In either case the maximum value of  $F$  is  $\frac{1}{2} + 2m - 1 = n - \frac{3}{2}$ , in which case  $\epsilon = \frac{3}{2n-3}$ .

Notice now that this maximum value for  $F$  can be just as easily achieved when  $\theta_p = -\theta_q$  and that then  $\sum_{i=1}^n \sin \theta_i = 0$  and  $\sum_{i=1}^n \cos \theta_i \sin \theta_i = 0$ . So  $\epsilon$  has in effect been minimized subject to all three constraints.  $\square$

**COROLLARY 3.4.** Let  $\sigma = \{1 + \epsilon, -1, -\epsilon, 0, \dots, 0\}$ , where  $1 \geq \epsilon \geq \frac{3}{2n-3}$ , and  $n \geq 3$ . Then  $\sigma$  can be realized as the spectrum of a symmetric nonnegative matrix  $A \in \mathbb{R}^{n \times n}$  with eigenvector  $e$  corresponding to  $1 + \epsilon$ , when  $n$  is odd.

*Proof.* Let  $F(\theta_1, \dots, \theta_n) = \sum_{j=1}^n \cos^2 \theta_j$ . Then

$$F\left(\frac{2\pi}{n}, \frac{2\pi}{n}, \dots, \frac{2\pi}{n}(n-1), \frac{2\pi}{n}n\right) = \frac{n}{2},$$

and

$$F\left(\frac{2\pi}{3}, -\frac{2\pi}{3}, 0, 0, \dots, 0, \pi, \pi, \dots, \pi\right) = n - \frac{3}{2}.$$

Moreover,  $F$  is continuous as a function of  $\cos \theta_1, \dots, \cos \theta_n$ , particularly on the following intervals for the  $\cos \theta_i$ 's:

For  $n = 4k + 3$  let the  $\cos \theta_i$ 's satisfy  $\cos \frac{2\pi(k+1)}{n} \leq \cos \theta_{k+1} \leq \cos \frac{2\pi}{3}$  and  $\cos \frac{2\pi(3k+3)}{n} \leq \cos \theta_{3k+3} \leq \cos \frac{2\pi}{3}$ . Also, let  $\cos \frac{2\pi}{n}j \leq \cos \theta_j \leq \cos 0$  for  $1 \leq j \leq k$  and  $3k + 4 \leq j \leq 4k + 3$ , and let  $\cos \pi \leq \cos \theta_j \leq \cos \frac{2\pi}{n}j$  for  $k + 2 \leq j \leq 3k + 2$  except that  $\cos \theta_{2k+2} = \cos \pi$ .

For  $n = 4k + 1$  let the  $\cos \theta_i$ 's satisfy  $\cos \frac{2\pi(k+1)}{n} \leq \cos \theta_{k+1} \leq \cos \frac{2\pi}{3}$  and  $\cos \frac{2\pi(3k+2)}{n} \leq \cos \theta_{3k+2} \leq \cos \frac{2\pi}{3}$ . Also, let  $\cos \frac{2\pi}{n}j \leq \cos \theta_j \leq \cos 0$  for  $1 \leq j \leq k$  and  $3k + 3 \leq j \leq 4k + 1$ , and let  $\cos \pi \leq \cos \theta_j \leq \cos \frac{2\pi}{n}j$  for  $k + 2 \leq j \leq 3k + 1$  except that  $\cos \theta_1 = \cos 0$ .

For each  $\epsilon$  such that  $1 \geq \epsilon \geq \frac{3}{2n-3}$  we have  $\frac{n}{2} \leq \frac{n}{1+\epsilon} \leq n - \frac{3}{2}$ , then from the Intermediate Value Theorem for real valued functions of several variables it follows that for each  $\epsilon \in [\frac{3}{2n-3}, 1]$  there is a  $(\theta_1, \dots, \theta_n)$  such that  $F(\theta_1, \dots, \theta_n) = \frac{n}{1+\epsilon}$ .  $\square$

The author does not see a natural extension of the above methods to deal with the case of at most four nonzero eigenvalues.

**Acknowledgement.** I am grateful to Ron Smith and Charles R. Johnson for some interesting conversations.

# REFERENCES

- [1] M. Boyle and D. Handelman. The spectra of nonnegative matrices via symbolic dynamics. *Annals of Mathematics*, 133:249–316, 1991.
- [2] M. Fiedler. Bounds for eigenvalues of doubly stochastic matrices. *Linear Algebra and Its Applications*, 5:299–310, 1972.
- [3] M. Fiedler. Eigenvalues of nonnegative symmetric matrices. *Linear Algebra and Its Applications*, 9:119–142, 1974.
- [4] M. Fiedler. Additive compound matrices and an inequality for eigenvalues of symmetric stochastic matrices. *Czechoslovak Mathematical Journal*, 24(99):392–402, 1974.
- [5] T.L. Hayden and P. Tarazaga. Distance matrices and regular figures. *Linear Algebra and its Applications*, 195:9-16, 1993.
- [6] T.L. Hayden, R. Reams, and J. Wells. Methods for constructing distance matrices and the inverse eigenvalue problem. *Linear Algebra and Its Applications*, 295:97–112, 1999.
- [7] T.J. Laffey and E. Meehan. A characterization of trace zero nonnegative  $5 \times 5$  matrices. *Linear Algebra and Its Applications*, 302/303:295–302, 1999.
- [8] R. Loewy and D. London. A note on an inverse problem for nonnegative matrices. *Linear and Multilinear Algebra*, 6:83–90, 1978.
- [9] H. Minc. *Nonnegative Matrices*. John Wiley and Sons, New York, 1988.
- [10] R. Reams. An inequality for nonnegative matrices and the inverse eigenvalue problem. *Linear and Multilinear Algebra*, 41:367-375, 1996.
- [11] D.O. Shklarsky, N.N. Chentzov, and I.M. Yaglom. *The USSR Olympiad Problem Book*. Dover Publications, Inc., New York, 1993.