

ON A CONJECTURE REGARDING CHARACTERISTIC POLYNOMIAL OF A MATRIX PAIR*

C. M. DA FONSECA†

Abstract. For n -by- n Hermitian matrices $A(> 0)$ and B , define

$$\eta(A, B) = \sum_S \det A(S) \det B(S'),$$

where the summation is over all subsets of $\{1, \dots, n\}$, S' is the complement of S , and by convention $\det A(\emptyset) = 1$. Bapat proved for $n = 3$ that the zeros of $\eta(\lambda A, -B)$ and the zeros of $\eta(\lambda A(23), -B(23))$ interlace. This result is generalized to a broader class of matrices.

Key words. Symmetric matrices, Cycles, Characteristic polynomial, Interlacing.

AMS subject classifications. 15A15, 15A42.

1. Introduction. Let $A = (a_{ij})$ and $B = (b_{ij})$ be matrices of order n . For index sets $S \subset \{1, \dots, n\}$, we denote by $A(S)$ the $|S| \times |S|$ principal submatrix lying in the rows and columns indexed by S . We may also denote $A(S')$ by A_S , with S' indexed the complement of S .

Define

$$(1.1) \quad \eta(A, B) := \sum_S \det A(S) \det B(S')$$

where the summation is over all subsets of $\{1, \dots, n\}$ and, by convention, $\det A(\emptyset) = \det B(\emptyset) = 1$. Notice that

$$(1.2) \quad \eta(\lambda I_n, -B) = \det(\lambda I_n - B),$$

i.e., $\eta(\lambda I_n, -B)$ is the characteristic polynomial of B . It is well-known that, if B is Hermitian, then the roots of (1.2), the eigenvalues of B , are all real. Motivated by this result, Johnson [3] considered the polynomial (of degree n)

$$(1.3) \quad \eta(\lambda A, -B) = \sum_{k=0}^n \sum_{|S|=k} (-1)^{n-k} \det A(S) \det B(S') \lambda^k,$$

and stated the conjecture:

CONJECTURE 1.1 (Johnson [3]). *If A and B are Hermitian and A is positive semidefinite, then the polynomial $\eta(\lambda A, -B)$ has only real roots.*

For a square matrix A , we write $A > 0$ to denote that A is positive definite. If all roots of the polynomial (1.3), say $\lambda_\ell^A(B)$, for $\ell = 1, \dots, n$, are real, we assume that

*Received by the editors 27 April 2005. Accepted for publication 25 May 2005. Handling Editor: Ravindra B. Bapat.

†Departamento de Matemática, Universidade de Coimbra, 3001-454 Coimbra, PORTUGAL (cmf@mat.uc.pt). This work was supported by CMUC - Centro de Matemática da Universidade Coimbra.

they have been arranged in increasing order $\lambda_1^A(B) \leq \dots \leq \lambda_n^A(B)$. Bapat in [1] and Johnson in [4] conjectured:

CONJECTURE 1.2. *If $A > 0$ and B are Hermitian, then $\lambda_\ell^{A_1}(B_1)$, for $\ell = 1, \dots, n-1$, interlace $\lambda_\ell^A(B)$, for $\ell = 1, \dots, n$, i.e.,*

$$\lambda_\ell^A(B) \leq \lambda_\ell^{A_1}(B_1) \leq \lambda_{\ell+1}^A(B), \quad \ell = 1, \dots, n-1.$$

Conjecture 1.1 has been verified for the case $n = 3$ by Rublein in [5] in a very complicated way. On the other hand, Bapat in [1] gave concise solutions for the cases $n \leq 3$. Bapat also verified that Conjectures 1.1 and 1.2 are true when both A and B are tridiagonal. Recently, the author generalized these results for matrices whose graph is a tree [2].

In this note we generalize the result of Bapat when $n = 3$ to matrices whose graph is a cycle.

For sake of simplicity we consider only symmetric matrices throughout. All the results can be easily generalized to Hermitian matrices.

2. Results on tridiagonal matrices. We define the weights of a symmetric matrix A as $w_{ij}(A) = -a_{ij}^2$ if $i \neq j$, and $w_{ii}(A) = a_{ii}$. Sometimes we abbreviate to w_{ij} , with no mention of A .

LEMMA 2.1 (Bapat [1]). *Let A and B be symmetric tridiagonal matrices and let $S = \{1, 2\}$. Then*

$$(2.1) \quad \eta(A, B) = \sum_{\ell \in S} (w_{1\ell}(A) + w_{1\ell}(B)) \eta(A_{1\ell}, B_{1\ell}).$$

For tridiagonal matrices we also state the following result, which can be proved by induction.

LEMMA 2.2. *Let A and B be symmetric tridiagonal matrices of order n . Then*

$$(2.2) \quad \eta(A, B) \eta(A_{1n}, B_{1n}) = \eta(A_1, B_1) \eta(A_n, B_n) - (a_{12}^2 + b_{12}^2) \cdots (a_{n-1,n}^2 + b_{n-1,n}^2).$$

Notice that (2.2) holds up to permutation similarity.

3. An interlacing theorem. Bapat proved the veracity of Conjectures 1.1 and 1.2 in the case $n \leq 3$.

THEOREM 3.1 (Bapat [1]). *Let A and B be Hermitian matrices of order 3 with $A > 0$ and B has all nonzero subdiagonal entries. Then $\eta(\lambda A, -B)$ has three real roots, say $\lambda_1 < \lambda_2 < \lambda_3$. Furthermore, if $\mu_1 < \mu_2$ are the roots of $\eta(\lambda A_1, -B_1)$, then $\lambda_1 < \mu_1 < \lambda_2 < \mu_2 < \lambda_3$.*

Consider symmetric matrices A and B such that $a_{ij} = b_{ij} = 0$ for $|i - j| > 1$ and $(i, j) \neq (1, n)$. We say, for obvious reasons, that A and B are matrices whose graph is a cycle. The next result generalizes Lemma 2.1, since if $w_{1,n} = 0$, we get (2.1) with $i = 1$.

LEMMA 3.2. Let A and B be symmetric matrices whose graph is a cycle and set $S = \{i-1, i, i+1\}$. Then

$$(3.1) \quad \eta(A, B) = \sum_{\ell \in S} (w_{i\ell}(A) + w_{i\ell}(B)) \eta(A_{i\ell}, B_{i\ell}) + 2(-1)^{n-1} \left(\prod_{\ell=1}^n a_{\ell, \ell+1} + \prod_{\ell=1}^n b_{\ell, \ell+1} \right),$$

with the convention $(n, n+1) = (1, n)$.

Proof. Let $C = \{i-1, i+1\}$. Considering the partition of all subsets of $\{1, \dots, n\}$, define

$$\mathcal{A}_P = \{S \mid i \in S, P \subset S, P' \cap S = \emptyset\}$$

and

$$\mathcal{C}_P = \{S \mid i \notin S, P \subset S, P' \cap S = \emptyset\},$$

for each subset P of C , where P' is the complement of P with respect to C . Evaluating $\det A(S)$ for each $S \in \mathcal{A}_P$, and $\det B(S')$ for each $S \in \mathcal{C}_P$, substituting in (1.3) the expressions obtained and finally rearranging the terms we get (3.1). \square

Without loss of generality, set $i = 1$. Notice that A_i and B_i are permutation similar to tridiagonal matrices. Suppose that B_1 is nonsingular and the subdiagonal entries of B are nonzero. From (3.1) we have

$$(3.2) \quad \begin{aligned} \eta(\lambda A, -B) &= (\lambda a_{11} - b_{11}) \eta(\lambda A_1, -B_1) \\ &\quad + (a_{12}, a_{1n}) P(\lambda) (a_{12}, a_{1n})^t + (b_{12}, b_{1n}) Q(\lambda) (b_{12}, b_{1n})^t, \end{aligned}$$

where

$$(3.3) \quad P(\lambda) = \begin{pmatrix} -\eta(\lambda A_{12}, -B_{12}) & (-)^{n-1} \lambda^{n-2} a_{23} \cdots a_{n-1, n} \\ (-)^{n-1} \lambda^{n-2} a_{23} \cdots a_{n-1, n} & -\eta(\lambda A_{1n}, -B_{1n}) \end{pmatrix}$$

and

$$(3.4) \quad Q(\lambda) = \begin{pmatrix} -\eta(\lambda A_{12}, -B_{12}) & -b_{23} \cdots b_{n-1, n} \\ -b_{23} \cdots b_{n-1, n} & -\eta(\lambda A_{1n}, -B_{1n}) \end{pmatrix}.$$

Suppose that the conjectures are true for such matrices of order less than $n-1$ in the conditions above, and proceed by induction on n . By hypothesis, $\eta(\lambda A_1, -B_1)$ has $n-1$ real roots, say $\mu_1 < \mu_2 < \cdots < \mu_{n-1}$, which strictly interlace the $n-2$ real roots of $\eta(\lambda A_{12}, -B_{12})$ and the $n-2$ real roots of $\eta(\lambda A_{1n}, -B_{1n})$. Since $\eta(\lambda A_{12}, -B_{12}), \eta(\lambda A_{1n}, -B_{1n}) \rightarrow \infty$ as $\lambda \rightarrow \infty$, the sign of $\eta(\mu_k A_{12}, -B_{12})$ and of $\eta(\mu_k A_{1n}, -B_{1n})$ must be $(-)^{n-k-1}$, for $k = 1, \dots, n-1$. Setting $\lambda = \mu_k$ in (3.3), we have

$$\det P(\mu_k) = \eta(\mu_k A_{12} - B_{12}) \eta(\mu_k A_{1n} - B_{1n}) - \mu_k^{2n-4} a_{23}^2 \cdots a_{n-1, n}^2.$$

According to (2.2), since $\eta(\mu_k A_1, -B_1) = 0$, we have

$$\text{sign } \det P(\mu_k) = +.$$

Analogously we can prove that $\text{sign det } Q(\mu_k) = +$. Therefore, $P(\mu_k)$ and $Q(\mu_k)$ are positive definite if $n - k - 1$ is odd, and negative definite if $n - k - 1$ is even. Hence

$$\begin{aligned} \text{sign } \eta(\mu_k A, -B) &= (-)^{n-k} + (-)^{n-k} \\ &= (-)^{n-k}, \quad k = 1, \dots, n-1. \end{aligned}$$

Since $\eta(\lambda A, -B) \rightarrow (\pm)^n \infty$ as $\lambda \rightarrow \pm\infty$, it follows that $\eta(\lambda A, -B)$ has a root in each of the intervals

$$(-\infty, \mu_1), (\mu_2, \mu_3), \dots, (\mu_{n-2}, \mu_{n-1}), (\mu_{n-1}, \infty),$$

and therefore $\eta(\lambda A, -B)$ has n distinct real roots, which strictly interlace $\mu_1, \mu_2, \dots, \mu_{n-1}$. Relaxing now by a continuity argument the nondegeneracy of the nonsingularity B_1 , we have:

THEOREM 3.3. *Let A and B be Hermitian matrices whose graph is a given cycle, with $A > 0$ and $b_{ij} \neq 0$ for $|i - j| = 1$. Then $\eta(\lambda A, -B)$ has n distinct real roots, say*

$$\lambda_1 < \lambda_2 < \dots < \lambda_n.$$

Furthermore, if

$$\mu_1 < \mu_2 < \dots < \mu_{n-1}$$

are the roots of $\eta(\lambda A_i, -B_i)$, $i = 1, \dots, n$, then

$$\lambda_1 < \mu_1 < \lambda_2 < \mu_2 < \dots < \mu_{n-1} < \lambda_n.$$

4. Example. Let us consider the Hermitian matrices

$$A = \begin{pmatrix} 3 & i & 0 & 1-i \\ -i & 2 & 1 & 0 \\ 0 & 1 & 4 & -2 \\ 1+i & 0 & -2 & 5 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} -1 & 2 & 0 & 1 \\ 2 & 0 & i & 0 \\ 0 & -i & 0 & -2 \\ 1 & 0 & -2 & -1 \end{pmatrix}.$$

The matrix A is positive definite and

$$\eta(\lambda A, -B) = 16 - 32\lambda - 97\lambda^2 + 44\lambda^3 + 47\lambda^4,$$

with roots

$$\begin{aligned} \lambda_1 &= -1.8109 \\ \lambda_2 &= -0.5646 \\ \lambda_3 &= 0.2895 \\ \lambda_4 &= 1.1498 \end{aligned}.$$

On the other hand

$$\eta(\lambda A_2, -B_2) = -4 - 12\lambda + 28\lambda^2 + 40\lambda^3,$$

with roots

$$\begin{aligned}\mu_1 &= -0.9090 \\ \mu_2 &= -0.2432 \\ \mu_3 &= 0.4522\end{aligned}.$$

Hence

$$\lambda_1 \leq \mu_1 \leq \lambda_2 \leq \mu_2 \leq \mu_3 \leq \lambda_4.$$

Finally, note that $\eta(\lambda A, -B)$ has as many positive and negative roots as the inertia of B $(2, 2, 0)$; see [4, (2)].

REFERENCES

- [1] R.B. Bapat. An interlacing theorem for tridiagonal matrices. *Linear Algebra Appl.*, 150:331–340, 1991.
- [2] C.M. da Fonseca. An interlacing theorem for matrices whose graph is a given tree. *J. Math. Sci.* (N.Y.), to appear.
- [3] C.R. Johnson. The permanent-on-top conjecture: A status report. *Current Trends in Matrix Theory*, pp. 167–174, 1987.
- [4] C.R. Johnson. A characteristic polynomial for matrix pairs. *Linear and Multilinear Algebra*, 25:289–290, 1989.
- [5] G. Rublein. On a conjecture of C. Johnson. *Linear and Multilinear Algebra*, 25:257–267, 1989.