



TOTALLY BIPARTITE TRIDIAGONAL PAIRS*

KAZUMASA NOMURA[†] AND PAUL TERWILLIGER[‡]

Abstract. There is a concept in linear algebra called a tridiagonal pair. The concept was motivated by the theory of Q -polynomial distance-regular graphs. We give a tutorial introduction to tridiagonal pairs, working with a special case as a concrete example. The special case is called totally bipartite, or totally bipartite (TB). Starting from first principles, we give an elementary but comprehensive account of TB tridiagonal pairs. The following topics are discussed: (i) the notion of a TB tridiagonal system; (ii) the eigenvalue array; (iii) the standard basis and matrix representations; (iv) the intersection numbers; (v) the Askey–Wilson relations; (vi) a recurrence involving the eigenvalue array; (vii) the classification of TB tridiagonal systems; (viii) self-dual TB tridiagonal pairs and systems; (ix) the \mathbb{Z}_3 -symmetric Askey–Wilson relations; (x) some automorphisms and antiautomorphisms associated with a TB tridiagonal pair; and (xi) an action of the modular group $\mathrm{PSL}_2(\mathbb{Z})$ associated with a TB tridiagonal pair.

Key words. Tridiagonal pair, Leonard pair, Tridiagonal matrix.

AMS subject classifications. 17B37, 15A21.

1. Introduction. This paper is about a linear algebraic object called a tridiagonal pair [71, Definition 1.1]. Before describing our purpose in detail, we give the definition of a tridiagonal pair. Let \mathbb{F} denote a field and let V denote a vector space over \mathbb{F} with finite positive dimension. A *tridiagonal pair* on V is an ordered pair of \mathbb{F} -linear maps $A : V \rightarrow V$ and $A^* : V \rightarrow V$ that satisfy the following conditions (i)–(iv).

- (i) Each of A, A^* is diagonalizable.
- (ii) There exists an ordering $\{V_i\}_{i=0}^d$ of the eigenspaces of A such that

$$(1.1) \quad A^*V_i \subseteq V_{i-1} + V_i + V_{i+1} \quad (0 \leq i \leq d),$$

where $V_{-1} = 0$ and $V_{d+1} = 0$.

- (iii) There exists an ordering $\{V_i^*\}_{i=0}^\delta$ of the eigenspaces of A^* such that

$$(1.2) \quad AV_i^* \subseteq V_{i-1}^* + V_i^* + V_{i+1}^* \quad (0 \leq i \leq \delta),$$

where $V_{-1}^* = 0$ and $V_{\delta+1}^* = 0$.

- (iv) There does not exist a subspace $W \subseteq V$ such that $AW \subseteq W, A^*W \subseteq W, W \neq 0, W \neq V$.

We summarize the history behind the tridiagonal pair concept. There is a type of finite undirected graph said to be distance-regular (see [31]). In [39], Delsarte investigated a class of distance-regular graphs said to be Q -polynomial. Given a Q -polynomial distance-regular graph, Delsarte obtained two sequences of orthogonal polynomials that are related by what is now called Askey–Wilson duality. In [93], Leonard classified the pairs of orthogonal polynomial sequences that obey this duality. He found that all the solutions come from the terminating branch of the Askey scheme, which consists of the q -Racah polynomials and their

*Received by the editors on November 1, 2017. Accepted for publication on May 13, 2021. Handling Editor: Bryan Shader. Corresponding Author: Kazumasa Nomura.

[†]Tokyo Medical and Dental University, Kohnodai Ichikawa, 272-0827 Japan (knomura@pop11.odn.ne.jp).

[‡]Department of Mathematics, University of Wisconsin, 480 Lincoln Drive, Madison, Wisconsin, 53706 USA (terwilli@math.wisc.edu).

limits (see [13]). In [15], Bannai and Ito gave a thorough study of Q -polynomial distance-regular graphs, including a detailed version of Leonard's theorem (see [15, Theorem 5.1]). In [132–134], the second author introduced an algebra $T = T(x)$, called the *subconstituent algebra* (or *Terwilliger algebra*), to analyze the subconstituents of a Q -polynomial distance-regular graph Γ with respect to a fixed vertex x . The algebra T is generated by the adjacency matrix A of Γ and a certain diagonal matrix A^* , called the dual adjacency matrix of Γ with respect to x . In [132, Lemmas 3.9, 3.12], it was observed that each of A, A^* acts on the eigenspaces of the other one in a block-tridiagonal fashion. This observation motivated the definition of a tridiagonal pair (see [71, Definition 1.1]).

This paper is meant for graduate students and researchers who seek an introduction to tridiagonal pairs. The theory of tridiagonal pairs is extensive, and some proofs are rather intricate. So for the beginner, it is best to start with a special case. In this paper, we consider a special case said to be totally bipartite (TB). A pair of \mathbb{F} -linear maps $A : V \rightarrow V$ and $A^* : V \rightarrow V$ is called a TB tridiagonal pair whenever it satisfies the above conditions (i)–(iv), with (1.1) replaced by

$$(1.3) \quad A^*V_i \subseteq V_{i-1} + V_{i+1} \quad (0 \leq i \leq d),$$

and (1.2) replaced by

$$(1.4) \quad AV_i^* \subseteq V_{i-1}^* + V_{i+1}^* \quad (0 \leq i \leq \delta).$$

We are not the first authors to consider the TB tridiagonal pairs. A number of previous articles are effectively about this topic, although they may not use the term. We now summarize these articles. In [32], Brown classified up to isomorphism a certain class of TB tridiagonal pairs, said to have Bannai/Ito type. Later in [64], Gao, Hou, and Wang classified up to isomorphism all the TB tridiagonal pairs. This classification separates the TB tridiagonal pairs into three infinite families, called Krawtchouk, Bannai/Ito, and q -Racah. It has been shown that a TB tridiagonal pair can be extended to a Leonard triple in the sense of Curtin [35]. This result is due to Balmaceda and Maralit for Krawtchouk type (see [14]), Brown for Bannai/Ito type (see [32]), and Gao, Hou, Wang for q -Racah type (see [64]). In [138], the second author showed that a finite-dimensional irreducible module for the Lie algebra \mathfrak{sl}_2 gives a TB tridiagonal pair of Krawtchouk type. In [7], Alnajjar and Curtin obtained a similar result using the equitable basis for \mathfrak{sl}_2 . The anticommutator spin algebra was introduced by Arik and Kayserilioglu [12]. In [32], Brown showed that certain irreducible modules for this algebra give TB tridiagonal pairs of Bannai/Ito type. In [59, 68], Havlíček, Pošta and Huang showed that certain irreducible $U_q(\mathfrak{so}_3)$ -modules give TB tridiagonal pairs of q -Racah type. We remark that in all the above articles except [7, 138] the field \mathbb{F} is assumed to be algebraically closed.

We just summarized the previous articles that are effectively about TB tridiagonal pairs. Among these, the articles [7, 14, 32, 35, 63, 64, 68] explicitly refer to the concept of a tridiagonal pair. What these cited papers have in common is that they invoke results about general tridiagonal pairs and then specialize to the TB case. In our view, this makes the theory more complicated than necessary. In our approach, we examine TB tridiagonal pairs from first principles and do not invoke any results from the literature about general tridiagonal pairs. For most of our results, we do not assume that \mathbb{F} is algebraically closed.

In this paper, we discuss the following topics: (i) the notion of a TB tridiagonal system; (ii) the eigenvalue array; (iii) the standard basis and matrix representations; (iv) the intersection numbers; (v) the Askey–Wilson relations; (vi) a recurrence involving the eigenvalue array; (vii) the classification of TB tridiagonal systems; (viii) self-dual TB tridiagonal pairs and systems; (ix) the \mathbb{Z}_3 -symmetric Askey–Wilson relations;

(x) some automorphisms and antiautomorphisms associated with a TB tridiagonal pair; and (xi) an action of the modular group $\text{PSL}_2(\mathbb{Z})$ associated with a TB tridiagonal pair.

We now summarize our results in detail. Let V denote a vector space over \mathbb{F} with finite positive dimension. Let $\text{End}(V)$ denote the \mathbb{F} -algebra consisting of the \mathbb{F} -linear maps from V to V . Let A, A^* denote a TB tridiagonal pair on V . Fix an ordering $\{V_i\}_{i=0}^d$ (resp. $\{V_i^*\}_{i=0}^\delta$) of the eigenspaces of A (resp. A^*) that satisfies (1.3) (resp. (1.4)). For $0 \leq i \leq d$ let $E_i \in \text{End}(V)$ denote the projection onto V_i . The elements $\{E_i\}_{i=0}^d$ are called the primitive idempotents of A . The primitive idempotents $\{E_i^*\}_{i=0}^\delta$ of A^* are similarly defined. We call the sequence $\Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^\delta)$ a TB tridiagonal system. For $0 \leq i \leq d$ let θ_i denote the eigenvalue of A for the eigenspace $E_i V$, and for $0 \leq i \leq \delta$ let θ_i^* denote the eigenvalue of A^* for the eigenspace $E_i^* V$. We call the sequence $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^\delta)$ the eigenvalue array of Φ . We show that $d = \delta$ (see Corollary 4.11). We show that the eigenspaces $E_i V, E_i^* V$ have dimension 1 for $0 \leq i \leq d$ (see Corollary 4.11). Fix a nonzero $v \in E_0 V$ and define $v_i = E_i^* v$ ($0 \leq i \leq d$). We show that $\{v_i\}_{i=0}^d$ form a basis for V (see Lemma 4.10). With respect to this basis, the matrix representing A^* is diagonal and the matrix representing A is tridiagonal with diagonal entries all zero. Let $\{c_i\}_{i=1}^d$ (resp. $\{b_i\}_{i=0}^{d-1}$) denote the subdiagonal entries (resp. superdiagonal entries) of this tridiagonal matrix. The scalars $\{c_i\}_{i=1}^d$ and $\{b_i\}_{i=0}^{d-1}$ are called the intersection numbers of Φ . We represent these intersection numbers in terms of the eigenvalue array (see Lemma 5.4). Using this we show that a TB tridiagonal system is uniquely determined up to isomorphism by its eigenvalue array (see Corollary 5.5). We show that A, A^* satisfy a pair of equations called the Askey–Wilson relations (see lines (8.42), (8.43)). We show that there exists $\beta \in \mathbb{F}$ such that

$$\theta_{i-1} - \beta\theta_i + \theta_{i+1} = 0, \quad \theta_{i-1}^* - \beta\theta_i^* + \theta_{i+1}^* = 0,$$

for $1 \leq i \leq d-1$ (see Proposition 9.10). We classify the TB tridiagonal systems up to isomorphism in terms of the eigenvalue array (see Theorem 11.1). We represent the eigenvalue array in closed form (see Examples 13.3–13.6). The TB tridiagonal pair A, A^* is said to be self-dual whenever it is isomorphic to A^*, A . We show that there exists $0 \neq \zeta \in \mathbb{F}$ such that $\zeta A, A^*$ is self-dual (see Lemma 14.8). In Theorem 14.9, we show that if A, A^* is self-dual, then the following TB tridiagonal pairs are mutually isomorphic:

$$(1.5) \quad \begin{array}{cccc} A, A^*, & A, -A^*, & -A, A^*, & -A, -A^*, \\ A^*, A, & A^*, -A, & -A^*, A, & -A^*, -A. \end{array}$$

The explicit isomorphisms are given in Section 14. We put the Askey–Wilson relations in a form said to be \mathbb{Z}_3 -symmetric (see Propositions 15.2–15.4). For our remaining results, we assume that \mathbb{F} is algebraically closed, and that A, B is self-dual, where $B = A^*$. We show that there exists $C \in \text{End}(V)$ such that A, B, C is a modular Leonard triple in the sense of Curtin (see [35]). We display some automorphisms and antiautomorphisms of $\text{End}(V)$ that act on A, B, C in an attractive manner. Recall that the modular group $\text{PSL}_2(\mathbb{Z})$ has a presentation by generators r, s and relations $r^3 = 1, s^2 = 1$. We show that $\text{PSL}_2(\mathbb{Z})$ acts on $\text{End}(V)$ as a group of automorphisms such that r sends $A \mapsto B \mapsto C \mapsto A$ and s sends $A \leftrightarrow B$ (see Corollary 16.11 and Proposition 17.9). In the last part of the paper, we describe general tridiagonal pairs, using the TB case as a guide.

The paper is organized as follows. In Section 2, we recall some materials from linear algebra. In Section 3, we define a TB tridiagonal system and its eigenvalue array. In Section 4, we discuss the diameter and eigenspace dimensions of a TB tridiagonal system. In Section 5, we discuss the intersection numbers of a TB tridiagonal system. In Sections 6 and 7 we obtain some linear algebra facts that will be needed later. In Section 8, we discuss the Askey–Wilson relations. In Sections 9 and 10, we investigate a recurrence satisfied by the eigenvalue array of a TB tridiagonal system. In Section 11, we classify the TB tridiagonal systems up



to isomorphism. In Sections 12 and 13, we examine in detail the eigenvalue array of a TB tridiagonal system. In Section 14, we show that in the self-dual case the TB tridiagonal pairs (1.5) are mutually isomorphic. In Section 15, we put the Askey–Wilson relations in \mathbb{Z}_3 -symmetric form. In Sections 16 and 17, we obtain some automorphisms and antiautomorphisms of $\text{End}(V)$ that act on a given TB tridiagonal pair in an attractive manner. In Section 18, we discuss an action of $\text{PSL}_2(\mathbb{Z})$. In Section 19, we describe the general tridiagonal pairs.

2. Preliminaries. In this section, we recall some materials from linear algebra. Throughout the paper, we use the following notation. Let \mathbb{F} denote a field and let $\overline{\mathbb{F}}$ denote the algebraic closure of \mathbb{F} . For an integer $n \geq 0$, let $\text{Mat}_{n+1}(\mathbb{F})$ denote the \mathbb{F} -algebra consisting of the $n + 1$ by $n + 1$ matrices that have all entries in \mathbb{F} . We index the rows and columns by $0, 1, \dots, n$. Let \mathbb{F}^{n+1} denote the vector space over \mathbb{F} consisting of the column vectors of length $n + 1$ that have all entries in \mathbb{F} . We index the rows by $0, 1, \dots, n$. Let V denote a vector space over \mathbb{F} with finite positive dimension. Let $\text{End}(V)$ denote the \mathbb{F} -algebra consisting of the \mathbb{F} -linear maps from V to V . For the \mathbb{F} -algebras $\text{Mat}_{n+1}(\mathbb{F})$ and $\text{End}(V)$, the identity element is denoted by I . For $X \in \text{Mat}_{n+1}(\mathbb{F})$ let X^t denote the transpose of X .

Let A denote an element of $\text{End}(V)$. For $\theta \in \mathbb{F}$ define $V(\theta) = \{v \in V \mid Av = \theta v\}$. Observe that $V(\theta)$ is a subspace of V . The scalar θ is called an *eigenvalue* of A whenever $V(\theta) \neq 0$. In this case, $V(\theta)$ is called the *eigenspace* of A corresponding to θ . We say that A is *diagonalizable* whenever V is spanned by the eigenspaces of A . Assume that A is diagonalizable. Let $\{V_i\}_{i=0}^d$ denote an ordering of the eigenspaces of A . So $V = \sum_{i=0}^d V_i$ (direct sum). For $0 \leq i \leq d$ let θ_i denote the eigenvalue of A corresponding to V_i . For $0 \leq i \leq d$ define $E_i \in \text{End}(V)$ such that $(E_i - I)V_i = 0$ and $E_i V_j = 0$ if $j \neq i$ ($0 \leq j \leq d$). Thus, E_i is the projection onto V_i . Observe that (i) $V_i = E_i V$ ($0 \leq i \leq d$); (ii) $E_i E_j = \delta_{i,j} E_i$ ($0 \leq i, j \leq d$); (iii) $I = \sum_{i=0}^d E_i$; (iv) $A = \sum_{i=0}^d \theta_i E_i$. Also

$$(2.6) \quad E_i = \prod_{\substack{0 \leq j \leq d \\ j \neq i}} \frac{A - \theta_j I}{\theta_i - \theta_j} \quad (0 \leq i \leq d).$$

We call E_i the *primitive idempotent of A for θ_i* ($0 \leq i \leq d$). Let M denote the subalgebra of $\text{End}(V)$ generated by A . Observe that $\{A^i\}_{i=0}^d$ is a basis for the \mathbb{F} -vector space M , and $\prod_{i=0}^d (A - \theta_i I) = 0$. Also observe that $\{E_i\}_{i=0}^d$ is a basis for the \mathbb{F} -vector space M .

Let $\{v_i\}_{i=0}^n$ denote a basis for V . For $A \in \text{End}(V)$ and $X \in \text{Mat}_{n+1}(\mathbb{F})$, we say that X *represents A with respect to $\{v_i\}_{i=0}^n$* whenever $Av_j = \sum_{i=0}^n X_{i,j} v_i$ for $0 \leq j \leq n$. For $A \in \text{End}(V)$ let A^b denote the matrix in $\text{Mat}_{n+1}(\mathbb{F})$ that represents A with respect to $\{v_i\}_{i=0}^n$. Then the map $\text{End}(V) \rightarrow \text{Mat}_{n+1}(\mathbb{F})$, $A \mapsto A^b$ is an isomorphism of \mathbb{F} -algebras.

For an \mathbb{F} -algebra \mathcal{A} , by an *automorphism of \mathcal{A}* we mean an isomorphism of \mathbb{F} -algebras from \mathcal{A} to \mathcal{A} . For an invertible $T \in \mathcal{A}$, the map $X \mapsto T^{-1}XT$ is an automorphism of \mathcal{A} , said to be *inner*. By the Skolem–Noether theorem [126, Corollary 7.125], every automorphism of $\text{End}(V)$ is inner.

At several places in the paper, we will discuss polynomials. Let x denote an indeterminate. Let $\mathbb{F}[x]$ denote the \mathbb{F} -algebra consisting of the polynomials in x that have all coefficients in \mathbb{F} .

3. Totally bipartite tridiagonal pairs. In this section, we define a totally bipartite tridiagonal pair and obtain some basic facts about this object.

DEFINITION 3.1 (See [32, Definition 1.2]). Let V denote a vector space over \mathbb{F} with finite positive dimension. A totally bipartite tridiagonal pair (or TB tridiagonal pair) on V is an ordered pair A, A^* of elements in $\text{End}(V)$ that satisfy the following (i)–(iv).

- (i) Each of A, A^* is diagonalizable.
- (ii) There exists an ordering $\{V_i\}_{i=0}^d$ of the eigenspaces of A such that

$$(3.7) \quad A^*V_i \subseteq V_{i-1} + V_{i+1} \quad (0 \leq i \leq d),$$

where $V_{-1} = 0$ and $V_{d+1} = 0$.

- (iii) There exists an ordering $\{V_i^*\}_{i=0}^\delta$ of the eigenspaces of A^* such that

$$(3.8) \quad AV_i^* \subseteq V_{i-1}^* + V_{i+1}^* \quad (0 \leq i \leq \delta),$$

where $V_{-1}^* = 0$ and $V_{\delta+1}^* = 0$.

- (iv) There does not exist a subspace $W \subseteq V$ such that $AW \subseteq W, A^*W \subseteq W, W \neq 0, W \neq V$.

We say that A, A^* is over \mathbb{F} . We call V the underlying vector space.

NOTE 3.2. According to a common notational convention, A^* denotes the conjugate-transpose of A . We are not using this convention. In a TB tridiagonal pair, the elements A and A^* are arbitrary subject to (i)–(iv) above.

NOTE 3.3. If A, A^* is a TB tridiagonal pair on V , then so is A^*, A .

NOTE 3.4. Let A, A^* denote a TB tridiagonal pair on V . Pick nonzero $\zeta, \zeta^* \in \mathbb{F}$. Then $\zeta A, \zeta^* A^*$ is a TB tridiagonal pair on V .

We mention a special case of a TB tridiagonal pair.

EXAMPLE 3.5. Referring to Definition 3.1, assume that $\dim V = 1$ and $A = 0, A^* = 0$. Then, A, A^* is a TB tridiagonal pair on V , said to be *trivial*.

LEMMA 3.6. With reference to Definition 3.1, assume that A, A^* is a TB tridiagonal pair on V . Then the following (i)–(vi) are equivalent: (i) A, A^* is trivial; (ii) $d = 0$; (iii) $\delta = 0$; (iv) $A = 0$; (v) $A^* = 0$; (vi) $\dim V = 1$.

Proof. Routine. □

For the rest of this section, let V denote a vector space over \mathbb{F} with finite positive dimension.

Let A, A^* denote a TB tridiagonal pair on V . Let V' denote a vector space over \mathbb{F} with finite positive dimension, and let A', A'^* denote a TB tridiagonal pair on V' . By an *isomorphism of TB tridiagonal pairs* from A, A^* to A', A'^* , we mean an \mathbb{F} -linear bijection $\psi : V \rightarrow V'$ such that $\psi A = A' \psi$ and $\psi A^* = A'^* \psi$. We say that A, A^* and A', A'^* are *isomorphic* whenever there exists an isomorphism of TB tridiagonal pairs from A, A^* to A', A'^* .

Let A, A^* denote a TB tridiagonal pair on V . An ordering $\{V_i\}_{i=0}^d$ of the eigenspaces of A is said to be *standard* whenever it satisfies (3.7). Let $\{V_i\}_{i=0}^d$ denote a standard ordering of the eigenspaces of A . Then the ordering $\{V_{d-i}\}_{i=0}^d$ is also standard and no further ordering is standard. An ordering of the eigenvalues or primitive idempotents for A is said to be *standard* whenever the corresponding ordering of the eigenspaces is standard. Similar comments apply to A^* .

DEFINITION 3.7. *By a TB tridiagonal system on V , we mean a sequence*

$$(3.9) \quad \Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^\delta),$$

of elements in $\text{End}(V)$ that satisfy the following (i)–(iii):

- (i) A, A^* is a TB tridiagonal pair on V ;
- (ii) $\{E_i\}_{i=0}^d$ is a standard ordering of the primitive idempotents of A ;
- (iii) $\{E_i^*\}_{i=0}^\delta$ is a standard ordering of the primitive idempotents of A^* .

We say that Φ is over \mathbb{F} . We call V the underlying vector space.

DEFINITION 3.8. *Referring to Definition 3.7, for notational convenience define $E_{-1} = 0$, $E_{d+1} = 0$, $E_{-1}^* = 0$, $E_{\delta+1}^* = 0$.*

DEFINITION 3.9. *Referring to Definition 3.7, we say that the TB tridiagonal pair A, A^* and the TB tridiagonal system Φ are associated.*

DEFINITION 3.10. *A TB tridiagonal system is said to be trivial whenever the associated TB tridiagonal pair is trivial.*

DEFINITION 3.11. *Consider the TB tridiagonal system Φ from (3.9). For $0 \leq i \leq d$ let θ_i denote the eigenvalue of A corresponding to E_i , and for $0 \leq i \leq \delta$ let θ_i^* denote the eigenvalue of A^* corresponding to E_i^* . We call $\{\theta_i\}_{i=0}^d$ (resp. $\{\theta_i^*\}_{i=0}^\delta$) the eigenvalue sequence (resp. dual eigenvalue sequence) of Φ . We call $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^\delta)$ the eigenvalue array of Φ .*

NOTE 3.12. *Referring to Definition 3.11, the scalars $\{\theta_i\}_{i=0}^d$ are mutually distinct and contained in \mathbb{F} . Moreover, the scalars $\{\theta_i^*\}_{i=0}^\delta$ are mutually distinct and contained in \mathbb{F} .*

NOTE 3.13. *Referring to Definition 3.11, assume that Φ is trivial. Then $\theta_0 = 0$ and $\theta_0^* = 0$.*

LEMMA 3.14. *Pick nonzero $\zeta, \zeta^* \in \mathbb{F}$. Let $(A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^\delta)$ denote a TB tridiagonal system on V , with eigenvalue array $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^\delta)$. Then*

$$(\zeta A; \{E_i\}_{i=0}^d; \zeta^* A^*; \{E_i^*\}_{i=0}^\delta),$$

is a TB tridiagonal system on V , with eigenvalue array $(\{\zeta\theta_i\}_{i=0}^d; \{\zeta^\theta_i^*\}_{i=0}^\delta)$.*

Proof. By Definitions 3.7, 3.11 and linear algebra. □

Consider the TB tridiagonal system Φ from (3.9). Let V' denote a vector space over \mathbb{F} with finite positive dimension, and let

$$\Phi' = (A'; \{E_i'\}_{i=0}^d; A'^*; \{E_i'^*\}_{i=0}^\delta),$$

denote a TB tridiagonal system on V' . By an *isomorphism of TB tridiagonal systems* from Φ to Φ' , we mean an \mathbb{F} -linear bijection $\psi : V \rightarrow V'$ such that $\psi A = A' \psi$, $\psi A^* = A'^* \psi$, $\psi E_i = E_i' \psi$ ($0 \leq i \leq d$), $\psi E_i^* = E_i'^* \psi$ ($0 \leq i \leq \delta$). We say that Φ and Φ' are *isomorphic* whenever there exists an isomorphism of TB tridiagonal systems from Φ to Φ' . Note that isomorphic TB tridiagonal systems have the same eigenvalue array.

Given a TB tridiagonal system $\Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^\delta)$ on V , each of the following is a TB tridiagonal system on V :

$$\begin{aligned} \Phi^* &= (A^*; \{E_i^*\}_{i=0}^\delta; A; \{E_i\}_{i=0}^d), \\ \Phi^\downarrow &= (A; \{E_i\}_{i=0}^d; A^*; \{E_{\delta-i}^*\}_{i=0}^\delta), \\ \Phi^\uparrow &= (A; \{E_{d-i}\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^\delta). \end{aligned}$$

Viewing $*$, \downarrow , \Downarrow as permutations on the set of all TB tridiagonal systems,

$$(3.10) \quad *^2 = \downarrow^2 = \Downarrow^2 = 1,$$

$$(3.11) \quad \Downarrow * = * \downarrow, \quad \downarrow * = * \Downarrow, \quad \Downarrow \downarrow = \downarrow \Downarrow.$$

The group generated by the symbols $*$, \downarrow , \Downarrow subject to the relations (3.10), (3.11) is the dihedral group D_4 . Recall that D_4 is the group of symmetries of a square and has 8 elements. The elements $*$, \downarrow , \Downarrow induce an action of D_4 on the set of all TB tridiagonal systems over \mathbb{F} . Two TB tridiagonal systems will be called *relatives* whenever they are in the same orbit of this D_4 action.

DEFINITION 3.15. Let Φ denote a TB tridiagonal system, and let $g \in D_4$. For any object f attached to Φ , let f^g denote the corresponding object attached to $\Phi^{g^{-1}}$.

LEMMA 3.16. Let Φ and Φ' denote TB tridiagonal systems over \mathbb{F} . Assume that Φ and Φ' are isomorphic, and let ψ denote an isomorphism of TB tridiagonal systems from Φ to Φ' . Then for $g \in D_4$ the map ψ is an isomorphism of TB tridiagonal systems from Φ^g to Φ'^g .

Proof. By the construction. □

LEMMA 3.17. Let A, A^* denote a TB tridiagonal pair over \mathbb{F} , and let Φ denote an associated TB tridiagonal system. Then the TB tridiagonal systems associated with A, A^* are $\Phi, \Phi^\downarrow, \Phi^\Downarrow, \Phi^{\downarrow\Downarrow}$.

Proof. By the comments above Definition 3.7. □

DEFINITION 3.18. Let A, A^* denote a TB tridiagonal pair over \mathbb{F} . By an eigenvalue sequence (resp. dual eigenvalue sequence) (resp. eigenvalue array) of A, A^* we mean the eigenvalue sequence (resp. dual eigenvalue sequence) (resp. eigenvalue array) of a TB tridiagonal system associated with A, A^* .

LEMMA 3.19. Let Φ denote a TB tridiagonal system over \mathbb{F} with eigenvalue array $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d)$. Then for $g \in \{*, \downarrow, \Downarrow\}$ the eigenvalue array of Φ^g is as follows:

g	Eigenvalue array of Φ^g
$*$	$(\{\theta_i^*\}_{i=0}^d; \{\theta_i\}_{i=0}^d)$
\downarrow	$(\{\theta_i\}_{i=0}^d; \{\theta_{\delta-i}^*\}_{i=0}^d)$
\Downarrow	$(\{\theta_{d-i}\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d)$

Proof. By the construction. □

LEMMA 3.20. For the TB tridiagonal system in (3.9), we have

$$\begin{aligned} AE_i^*V &\subseteq E_{i-1}^*V + E_{i+1}^*V & (0 \leq i \leq \delta), \\ A^*E_iV &\subseteq E_{i-1}V + E_{i+1}V & (0 \leq i \leq d). \end{aligned}$$

Proof. By Definition 3.1(ii), (iii). □

LEMMA 3.21. For the TB tridiagonal system in (3.9), we have

$$(3.12) \quad E_i^*AE_j^* = \begin{cases} 0 & \text{if } |i-j| \neq 1, \\ \neq 0 & \text{if } |i-j| = 1 \end{cases} \quad (0 \leq i, j \leq \delta),$$

$$(3.13) \quad E_iA^*E_j = \begin{cases} 0 & \text{if } |i-j| \neq 1, \\ \neq 0 & \text{if } |i-j| = 1 \end{cases} \quad (0 \leq i, j \leq d).$$

Proof. We first show that

$$(3.14) \quad E_i^* A E_j^* = 0 \quad \text{if } |i - j| \neq 1 \quad (0 \leq i, j \leq \delta).$$

By Lemma 3.20,

$$(3.15) \quad A E_j^* V \subseteq E_{j-1}^* V + E_{j+1}^* V.$$

In the above line, apply E_i^* to each side. The right-hand side becomes 0 since $i \neq j - 1$ and $i \neq j + 1$. Thus, $E_i^* A E_j^* V = 0$ and so (3.14) holds. Next we show that

$$(3.16) \quad E_i^* A E_j^* \neq 0 \quad \text{if } |i - j| = 1 \quad (1 \leq i, j \leq \delta).$$

Consider the case $i = j - 1$. Assume by way of contradiction that $E_i^* A E_j^* = 0$. For $v \in A E_j^* V$ we have $E_{j-1}^* v = 0$. By this and (3.15) we find $v \in E_{j+1}^* V$. By these comments

$$(3.17) \quad A E_j^* V \subseteq E_{j+1}^* V.$$

Define $W = E_j^* V + \cdots + E_\delta^* V$. We have $AW \subseteq W$ by Lemma 3.20 and (3.17). By construction $A^*W \subseteq W$, $W \neq 0$, $W \neq V$. This contradicts Definition 3.1(iv), and therefore (3.16) holds for $i = j - 1$. A similar argument shows that (3.16) holds for $j = i - 1$. We have shown (3.12). To get (3.13), apply (3.12) to Φ^* . \square

4. The raising and lowering maps. Throughout this section, let V denote a vector space over \mathbb{F} with finite positive dimension, and let

$$\Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^\delta),$$

denote a TB tridiagonal system on V . Our next general goal is to show that $d = \delta$, and that each of $E_i V$ and $E_i^* V$ has dimension one for $0 \leq i \leq d$. To this end, it is convenient to introduce two maps R, L called the raising and lowering maps. Let $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^\delta)$ denote the eigenvalue array of Φ .

DEFINITION 4.1. Define $R, L \in \text{End}(V)$ by

$$R = \sum_{i=1}^{\delta} E_i^* A E_{i-1}^*, \quad L = \sum_{i=1}^{\delta} E_{i-1}^* A E_i^*.$$

We call R (resp. L) the raising map (resp. lowering map) for Φ .

LEMMA 4.2. We have $A = R + L$.

Proof. Evaluate $A = I A I$ using $I = \sum_{i=0}^{\delta} E_i^*$ to get $A = \sum_{i=0}^{\delta} \sum_{j=0}^{\delta} E_i^* A E_j^*$. In this equation evaluate the right-hand side using Lemma 3.21 and Definition 4.1. \square

LEMMA 4.3. The following (i)–(iv) hold:

- (i) $E_i^* R = E_i^* A E_{i-1}^* = R E_{i-1}^* \quad (1 \leq i \leq \delta)$;
- (ii) $E_0^* R = 0, \quad R E_\delta^* = 0$;
- (iii) $E_{i-1}^* L = E_{i-1}^* A E_i^* = L E_i^* \quad (1 \leq i \leq \delta)$;
- (iv) $E_\delta^* L = 0, \quad L E_0^* = 0$.

Proof. Recall that $E_r^* E_s^* = \delta_{r,s} E_r^*$ for $0 \leq r, s \leq \delta$. Using this and Definition 4.1, we routinely obtain the results. \square

LEMMA 4.4. *The following hold:*

- (i) $RE_i^*V \subseteq E_{i+1}^*V \quad (0 \leq i \leq \delta)$;
- (ii) $LE_i^*V \subseteq E_{i-1}^*V \quad (0 \leq i \leq \delta)$.

Proof. Use Lemma 4.3. □

DEFINITION 4.5. *Fix $0 \neq v \in E_0V$. For $0 \leq i \leq \delta$ define $v_i = E_i^*v$. For notational convenience, define $v_{-1} = 0$ and $v_{\delta+1} = 0$.*

LEMMA 4.6. *With reference to Definition 4.5, the following hold:*

- (i) $v_i \in E_i^*V \quad (0 \leq i \leq \delta)$;
- (ii) $v = \sum_{i=0}^{\delta} v_i$.

Proof. (i) By Definition 4.5.

- (ii) Use $I = \sum_{i=0}^{\delta} E_i^*$. □

We now describe the action of R and L on $\{v_i\}_{i=0}^{\delta}$. Note by Lemma 4.4 that $Rv_{\delta} = 0$ and $Lv_0 = 0$.

LEMMA 4.7. *With reference to Definition 4.5,*

$$\theta_0 v_i = Rv_{i-1} + Lv_{i+1} \quad (0 \leq i \leq \delta).$$

Proof. We have $v \in E_0V$ so $Av = \theta_0 v$. Using Lemmas 4.2 and 4.3,

$$\begin{aligned} 0 &= E_i^*(A - \theta_0 I)v \\ &= E_i^*(R + L - \theta_0 I)v \\ &= (RE_{i-1}^* + LE_{i+1}^* - \theta_0 E_i^*)v \\ &= Rv_{i-1} + Lv_{i+1} - \theta_0 v_i. \end{aligned}$$

The result follows. □

LEMMA 4.8. *Assume that Φ is nontrivial. Then with reference to Definition 4.5, the following (i)–(iii) hold:*

- (i) $\theta_1 \theta_0^* v_0 = \theta_1^* Lv_1$;
- (ii) $\theta_1 \theta_i^* v_i = \theta_{i-1}^* Rv_{i-1} + \theta_{i+1}^* Lv_{i+1} \quad (1 \leq i \leq \delta - 1)$;
- (iii) $\theta_1 \theta_{\delta}^* v_{\delta} = \theta_{\delta-1}^* Rv_{\delta-1}$.

Proof. (ii) We have $v \in E_0V$ and $A^*E_0V \subseteq E_1V$, so $A^*v \in E_1V$. Therefore $AA^*v = \theta_1 A^*v$. Using Lemmas 4.2, 4.3 and $E_r^*A^* = \theta_r^* E_r^*$ ($0 \leq r \leq d$),

$$\begin{aligned} 0 &= E_i^*(A - \theta_1 I)A^*v \\ &= E_i^*(R + L - \theta_1 I)A^*v \\ &= (RE_{i-1}^* + LE_{i+1}^* - \theta_1 E_i^*)A^*v \\ &= \theta_{i-1}^* Rv_{i-1} + \theta_{i+1}^* Lv_{i+1} - \theta_1 \theta_i^* v_i. \end{aligned}$$

The result follows.

- (i), (iii) Similar to the proof of (ii) above. □

LEMMA 4.9. *Assume that Φ is nontrivial. Then with reference to Definition 4.5,*

$$\begin{aligned} Rv_{i-1} &= \frac{\theta_1\theta_i^* - \theta_0\theta_{i+1}^*}{\theta_{i-1}^* - \theta_{i+1}^*} v_i & (1 \leq i \leq \delta - 1), & & Rv_{\delta-1} &= \theta_0v_\delta, \\ Lv_{i+1} &= \frac{\theta_1\theta_i^* - \theta_0\theta_{i-1}^*}{\theta_{i+1}^* - \theta_{i-1}^*} v_i & (1 \leq i \leq \delta - 1), & & Lv_1 &= \theta_0v_0. \end{aligned}$$

Also

$$(\theta_0\theta_1^* - \theta_1\theta_0^*)v_0 = 0, \quad (\theta_0\theta_{\delta-1}^* - \theta_1\theta_\delta^*)v_\delta = 0.$$

Proof. Solve the linear equations in Lemmas 4.7 and 4.8. □

LEMMA 4.10. *With reference to Definition 4.5, the following hold:*

- (i) v_i is a basis for E_i^*V ($0 \leq i \leq \delta$);
- (ii) $\{v_i\}_{i=0}^\delta$ is a basis for V .

Proof. Assume that Φ is nontrivial; otherwise, the result holds by Lemma 3.6. The sum $V = \sum_{i=0}^\delta E_i^*V$ is direct. For $0 \leq i \leq \delta$, the subspace E_i^*V is nonzero and contains v_i . So it suffices to show that the vectors $\{v_i\}_{i=0}^\delta$ span V . Let W denote the subspace of V spanned by $\{v_i\}_{i=0}^\delta$. Note that $W \neq 0$ since $0 \neq v \in W$. We have $A^*W \subseteq W$ by construction. Using Lemma 4.9 we find that $RW \subseteq W$ and $LW \subseteq W$. So $AW \subseteq W$ in view of Lemma 4.2. Now $W = V$ by Definition 3.1(iv). Therefore, $\{v_i\}_{i=0}^\delta$ span V . The result follows. □

COROLLARY 4.11. *We have $d = \delta$ and*

$$\dim E_iV = 1, \quad \dim E_i^*V = 1 \quad (0 \leq i \leq d).$$

Moreover $\dim V = d + 1$.

Proof. By Lemma 4.10, $\dim E_i^*V = 1$ ($0 \leq i \leq \delta$) and $\dim V = \delta + 1$. Applying Lemma 4.10 to Φ^* , we obtain $\dim E_iV = 1$ ($0 \leq i \leq d$) and $\dim V = d + 1$. The result follows. □

NOTE 4.12. A tridiagonal pair A, A^* is often called a Leonard pair if all the eigenspaces of A and A^* have dimension one.

DEFINITION 4.13. *Recall from Corollary 4.11 that $d = \delta$. We call this common value the diameter of Φ .*

DEFINITION 4.14. *A basis $\{v_i\}_{i=0}^d$ for V is said to be Φ -standard whenever there exists a nonzero $v \in E_0V$ such that $v_i = E_i^*v$ for $0 \leq i \leq d$.*

A Φ -standard basis is characterized as follows.

LEMMA 4.15. *Given vectors $\{v_i\}_{i=0}^d$ in V , not all zero. Then the following are equivalent:*

- (i) $v_i \in E_i^*V$ for $0 \leq i \leq d$, and $\sum_{i=0}^d v_i \in E_0V$;
- (ii) $\{v_i\}_{i=0}^d$ is a Φ -standard basis for V .

Proof. (i) \Rightarrow (ii) Define $v = \sum_{j=0}^d v_j$. By construction $v \in E_0V$. For $0 \leq i \leq d$, in the equation $v = \sum_{j=0}^d v_j$ apply E_i^* to each side to obtain $E_i^*v = v_i$. Note that $v \neq 0$ since the vectors $\{v_i\}_{i=0}^d$ are not all zero. The vectors $\{v_i\}_{i=0}^d$ form a basis for V by Lemma 4.10. This basis is Φ -standard by Definition 4.14.

(ii) \Rightarrow (i) By Definition 4.14 there exists $0 \neq v \in E_0V$ such that $v_i = E_i^*v$ for $0 \leq i \leq d$. By construction $v_i \in E_i^*V$ for $0 \leq i \leq d$. Also $\sum_{i=0}^d v_i = v \in E_0V$. □

We mention a lemma for later use.

LEMMA 4.16. *Assume that Φ is nontrivial. Then the following (i)–(iii) hold.*

- (i) *Each of $\theta_0, \theta_d, \theta_0^*, \theta_d^*$ is nonzero.*
- (ii) *$\theta_1/\theta_0 = \theta_{d-1}/\theta_d = \theta_1^*/\theta_0^* = \theta_{d-1}^*/\theta_d^*$.*
- (iii) *$\theta_0\theta_{d-1} = \theta_1\theta_d$ and $\theta_0^*\theta_{d-1}^* = \theta_1^*\theta_d^*$.*

Proof. By the last assertion of Lemma 4.9 and since $d = \delta, v_0 \neq 0, v_d \neq 0$,

$$(4.18) \quad \theta_0\theta_1^* = \theta_1\theta_0^*,$$

$$(4.19) \quad \theta_0\theta_{d-1}^* = \theta_1\theta_d^*.$$

Applying (4.19) to Φ^* ,

$$(4.20) \quad \theta_{d-1}\theta_0^* = \theta_d\theta_1^*.$$

Suppose that $\theta_0 = 0$. Then $\theta_1 \neq 0$ since $\{\theta_i\}_{i=0}^d$ are mutually distinct. Now $\theta_0^* = 0$ by (4.18) and $\theta_d^* = 0$ by (4.19). This is a contradiction since $\{\theta_i^*\}_{i=0}^d$ are mutually distinct and $d \geq 1$. Therefore, $\theta_0 \neq 0$. Applying this to $\Phi^*, \Phi^\dagger, \Phi^\natural$, we obtain $\theta_d \neq 0, \theta_0^* \neq 0, \theta_d^* \neq 0$. We have obtained assertion (i). Assertion (ii) follows in view of (4.18)–(4.20). Assertion (iii) follows from assertion (ii). \square

5. The intersection numbers. In this section, we introduce the intersection numbers of a TB tridiagonal system and express these intersection numbers in terms of the eigenvalue array. We use this result to show that a TB tridiagonal system is uniquely determined up to isomorphism by its eigenvalue array. Let V denote a vector space over \mathbb{F} with finite positive dimension, and let

$$\Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d),$$

denote a TB tridiagonal system on V . Let $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d)$ denote the eigenvalue array of Φ . Let $\{v_i\}_{i=0}^d$ denote a Φ -standard basis for V . For $X \in \text{End}(V)$ let X^\natural denote the matrix in $\text{Mat}_{d+1}(\mathbb{F})$ that represents X with respect to the basis $\{v_i\}_{i=0}^d$. By construction,

$$(5.21) \quad (E_i^*)^\natural = \text{diag}(0, \dots, 0, \overset{i}{1}, 0, \dots, 0) \quad (0 \leq i \leq d),$$

$$(5.22) \quad (A^*)^\natural = \text{diag}(\theta_0^*, \theta_1^*, \dots, \theta_d^*).$$

Also by construction there exist scalars $\{c_i\}_{i=1}^d, \{b_i\}_{i=0}^{d-1}$ in \mathbb{F} such that

$$(5.23) \quad A^\natural = \begin{pmatrix} 0 & b_0 & & & \mathbf{0} \\ c_1 & 0 & b_1 & & \\ & c_2 & \cdot & \cdot & \\ & & \cdot & \cdot & \cdot \\ \mathbf{0} & & & \cdot & \cdot & b_{d-1} \\ & & & c_d & & 0 \end{pmatrix}.$$

By Definition 4.1,

$$(5.24) \quad R^\natural = \begin{pmatrix} 0 & & & & \mathbf{0} \\ c_1 & 0 & & & \\ & c_2 & \cdot & & \\ & & \cdot & \cdot & \\ & & & \cdot & \cdot \\ \mathbf{0} & & & c_d & 0 \end{pmatrix}, \quad L^\natural = \begin{pmatrix} 0 & b_0 & & & \mathbf{0} \\ & 0 & b_1 & & \\ & & \cdot & \cdot & \\ & & & \cdot & \cdot \\ & & & & \cdot & b_{d-1} \\ \mathbf{0} & & & & & 0 \end{pmatrix}.$$

The scalars $\{c_i\}_{i=1}^d$, $\{b_i\}_{i=0}^{d-1}$ are called the *intersection numbers of Φ* . The intersection numbers $\{c_i^*\}_{i=1}^d$, $\{b_i^*\}_{i=0}^{d-1}$ of Φ^* are called the *dual intersection numbers of Φ* .

LEMMA 5.1. *Assume that Φ is nontrivial. Then*

$$\begin{aligned} Av_0 &= c_1v_1, \\ Av_i &= b_{i-1}v_{i-1} + c_{i+1}v_{i+1} \quad (1 \leq i \leq d-1), \\ Av_d &= b_{d-1}v_{d-1}. \end{aligned}$$

Proof. By (5.23). □

LEMMA 5.2. *We have*

$$\begin{aligned} Rv_i &= c_{i+1}v_{i+1} \quad (0 \leq i \leq d-1), & Rv_d &= 0, \\ Lv_i &= b_{i-1}v_{i-1} \quad (1 \leq i \leq d), & Lv_0 &= 0. \end{aligned}$$

Proof. By (5.24). □

LEMMA 5.3. *The scalars $\{b_i\}_{i=0}^{d-1}$, $\{c_i\}_{i=1}^d$ are all nonzero.*

Proof. We first show that $c_i \neq 0$ for $1 \leq i \leq d$. Let i be given. We have $E_i^*AE_{i-1}^* \neq 0$ by Lemma 3.21. By (5.21) and (5.23), the matrix $(E_i^*AE_{i-1}^*)^\natural$ has $(i, i-1)$ -entry c_i and all other entries 0. By these comments $c_i \neq 0$. One similarly shows that $b_i \neq 0$ for $0 \leq i \leq d-1$. □

LEMMA 5.4. *Assume that Φ is nontrivial. Then the following hold.*

(i) *The intersection numbers of Φ satisfy*

$$(5.25) \quad c_i = \frac{\theta_1\theta_i^* - \theta_0\theta_{i+1}^*}{\theta_{i-1}^* - \theta_{i+1}^*} \quad (1 \leq i \leq d-1), \quad c_d = \theta_0,$$

$$(5.26) \quad b_i = \frac{\theta_1\theta_i^* - \theta_0\theta_{i-1}^*}{\theta_{i+1}^* - \theta_{i-1}^*} \quad (1 \leq i \leq d-1), \quad b_0 = \theta_0.$$

(ii) *The dual intersection numbers of Φ satisfy*

$$(5.27) \quad c_i^* = \frac{\theta_1^*\theta_i - \theta_0^*\theta_{i+1}}{\theta_{i-1} - \theta_{i+1}} \quad (1 \leq i \leq d-1), \quad c_d^* = \theta_0^*,$$

$$(5.28) \quad b_i^* = \frac{\theta_1^*\theta_i - \theta_0^*\theta_{i-1}}{\theta_{i+1} - \theta_{i-1}} \quad (1 \leq i \leq d-1), \quad b_0^* = \theta_0^*.$$

Proof. (i) By Lemmas 4.9 and 5.2.

(ii) Apply (i) above to Φ^* . □

COROLLARY 5.5. *A TB tridiagonal system is uniquely determined up to isomorphism by its eigenvalue array.*

Proof. By (5.22), (5.23) and Lemma 5.4(i), the entries of A^\natural and $(A^*)^\natural$ are determined by the eigenvalue array. □

LEMMA 5.6. *For nonzero ζ , $\zeta^* \in \mathbb{F}$, consider the TB tridiagonal system*

$$\tilde{\Phi} = (\zeta A; \{E_i\}_{i=0}^d; \zeta^* A^*; \{E_i^*\}_{i=0}^d).$$

Then the (dual) intersection numbers of $\tilde{\Phi}$ are

$$\begin{aligned} \check{c}_i &= \zeta c_i, & \check{c}_i^* &= \zeta^* c_i^* & (1 \leq i \leq d), \\ \check{b}_i &= \zeta b_i, & \check{b}_i^* &= \zeta^* b_i^* & (0 \leq i \leq d-1). \end{aligned}$$

Proof. Use Lemmas 3.14 and 5.4. □

6. Tridiagonal matrices. In this section, we collect some results about tridiagonal matrices that will be used later in the paper. Consider a tridiagonal matrix in $\text{Mat}_{d+1}(\mathbb{F})$:

$$(6.29) \quad A = \begin{pmatrix} a_0 & b_0 & & & \mathbf{0} \\ c_1 & a_1 & b_1 & & \\ & c_2 & \cdot & \cdot & \\ & & \cdot & \cdot & \\ & & & \cdot & \cdot & b_{d-1} \\ \mathbf{0} & & & & c_d & a_d \end{pmatrix}.$$

We say that A is *irreducible* whenever $c_i b_{i-1} \neq 0$ for $1 \leq i \leq d$. For the rest of this section, assume that A is irreducible.

We have some remarks about the minimal polynomial of A . For $0 \leq r \leq d$ consider the matrix A^r . For $0 \leq i, j \leq d$ the (i, j) -entry of A^r satisfies

$$(6.30) \quad (A^r)_{i,j} = \begin{cases} 0 & \text{if } |i-j| > r, \\ \neq 0 & \text{if } |i-j| = r. \end{cases}$$

Consequently, the matrices $\{A^r\}_{r=0}^d$ are linearly independent. Therefore, the minimal polynomial of A coincides with the characteristic polynomial of A . So each eigenspace of A has dimension one. We remark that A might not be diagonalizable.

For $0 \leq i \leq d$ define $E_i^* \in \text{Mat}_{d+1}(\mathbb{F})$ by

$$(6.31) \quad E_i^* = \text{diag}(0, \dots, 0, \overset{i}{1}, 0, \dots, 0).$$

Observe that $E_i^* E_j^* = \delta_{i,j} E_i^*$ ($0 \leq i, j \leq d$) and $I = \sum_{i=0}^d E_i^*$.

LEMMA 6.1. For $0 \leq i, j, r \leq d$,

$$(6.32) \quad E_i^* A^r E_j^* = \begin{cases} 0 & \text{if } |i-j| > r, \\ \neq 0 & \text{if } |i-j| = r. \end{cases}$$

Proof. This is a reformulation of (6.30). □

LEMMA 6.2. The elements

$$(6.33) \quad \{A^r E_0^* A^s \mid 0 \leq r, s \leq d\}$$

form a basis for the \mathbb{F} -vector space $\text{Mat}_{d+1}(\mathbb{F})$.

Proof. For $0 \leq r, s \leq d$ consider the matrix $A^r E_0^* A^s$. For $0 \leq i, j \leq d$, we compute its (i, j) -entry using (6.31) and evaluate the result using (6.30):

$$(A^r E_0^* A^s)_{i,j} = (A^r)_{i,0} (A^s)_{0,j} = \begin{cases} 0 & \text{if } i > r \text{ or } j > s, \\ \neq 0 & \text{if } i = r \text{ and } j = s. \end{cases}$$

From the pattern of zero/nonzero entries, we see that the elements (6.33) are linearly independent. The set (6.33) contains $(d+1)^2$ elements, and this is the dimension of $\text{Mat}_{d+1}(\mathbb{F})$. The result follows. □

COROLLARY 6.3. *The elements A, E_0^* generate the \mathbb{F} -algebra $\text{Mat}_{d+1}(\mathbb{F})$.*

Proof. By Lemma 6.2. □

DEFINITION 6.4. *Let $\{\theta_i^*\}_{i=0}^d$ denote scalars in \mathbb{F} . Define $A^* \in \text{Mat}_{d+1}(\mathbb{F})$ by*

$$(6.34) \quad A^* = \text{diag}(\theta_0^*, \theta_1^*, \dots, \theta_d^*).$$

Observe that

$$(6.35) \quad A^* = \sum_{i=0}^d \theta_i^* E_i^*.$$

LEMMA 6.5. *Assume that $\theta_0^* \neq \theta_i^*$ for $1 \leq i \leq d$. Then the following (i)–(iii) hold.*

(i) *The element E_0^* is a polynomial in A^* :*

$$E_0^* = \prod_{i=1}^d \frac{A^* - \theta_i^* I}{\theta_0^* - \theta_i^*}.$$

(ii) *The elements A, A^* generate the \mathbb{F} -algebra $\text{Mat}_{d+1}(\mathbb{F})$.*

(iii) *There does not exist a subspace $W \subseteq \mathbb{F}^{d+1}$ such that $W \neq 0, W \neq \mathbb{F}^{d+1}, AW \subseteq W, A^*W \subseteq W$.*

Proof. (i) By matrix multiplication.

(ii) By (i) and Corollary 6.3.

(iii) By (ii) above and since \mathbb{F}^{d+1} is an irreducible $\text{Mat}_{d+1}(\mathbb{F})$ -module. □

DEFINITION 6.6. *Define scalars $\{k_i\}_{i=0}^d$ by*

$$(6.36) \quad k_i = \frac{b_0 b_1 \cdots b_{i-1}}{c_1 c_2 \cdots c_i} \quad (0 \leq i \leq d).$$

Observe that $k_0 = 1$, and $k_i \neq 0$ for $0 \leq i \leq d$. Moreover,

$$(6.37) \quad k_{i-1} b_{i-1} = k_i c_i \quad (1 \leq i \leq d).$$

DEFINITION 6.7. *Define $K \in \text{Mat}_{d+1}(\mathbb{F})$ by*

$$(6.38) \quad K = \text{diag}(k_0, k_1, \dots, k_d).$$

LEMMA 6.8. *We have $A^\dagger K = KA$.*

Proof. Use (6.29) and (6.37). □

For an \mathbb{F} -algebra \mathcal{A} , by an *antiautomorphism* of \mathcal{A} , we mean an \mathbb{F} -linear bijection $\xi : \mathcal{A} \rightarrow \mathcal{A}$ such that $(xy)^\xi = y^\xi x^\xi$ for $x, y \in \mathcal{A}$.

DEFINITION 6.9. *Define the map*

$$\dagger : \text{Mat}_{d+1}(\mathbb{F}) \rightarrow \text{Mat}_{d+1}(\mathbb{F}), \quad X \mapsto K^{-1} X^\dagger K.$$

LEMMA 6.10. *The map \dagger is an antiautomorphism of $\text{Mat}_{d+1}(\mathbb{F})$. Moreover, $(X^\dagger)^\dagger = X$ for all $X \in \text{Mat}_{d+1}(\mathbb{F})$.*

Proof. The map \dagger is \mathbb{F} -linear. For $X \in \text{Mat}_{d+1}(\mathbb{F})$ we have $(X^\dagger)^\dagger = X$, since

$$(X^\dagger)^\dagger = (K^{-1}X^tK)^\dagger = K^{-1}(K^{-1}X^tK)^tK = K^{-1}KXK^{-1}K = X.$$

Consequently, the map \dagger is bijective. For $X, Y \in \text{Mat}_{d+1}(\mathbb{F})$, we have

$$(XY)^\dagger = K^{-1}(XY)^tK = K^{-1}Y^tX^tK = K^{-1}Y^tKK^{-1}X^tK = Y^\dagger X^\dagger.$$

Thus \dagger is an antiautomorphism. □

The map \dagger is characterized as follows.

LEMMA 6.11. *The map \dagger is the unique antiautomorphism of $\text{Mat}_{d+1}(\mathbb{F})$ that fixes each of $A, E_0^*, E_1^*, \dots, E_d^*$.*

Proof. Using Lemma 6.8 and Definition 6.9,

$$A^\dagger = K^{-1}A^tK = K^{-1}KA = A,$$

so \dagger fixes A . For $0 \leq i \leq d$ the map \dagger fixes E_i^* by Definition 6.9 and since E_i^*, K are diagonal. Concerning the uniqueness, let ξ denote an antiautomorphism of $\text{Mat}_{d+1}(\mathbb{F})$ that fixes each of $A, E_0^*, E_1^*, \dots, E_d^*$. We show that $\xi = \dagger$. The composition $\xi\dagger$ is an automorphism of $\text{Mat}_{d+1}(\mathbb{F})$ that fixes $A, E_0^*, E_1^*, \dots, E_d^*$. Consequently $\xi\dagger = 1$ in view of Corollary 6.3. So $\xi = \dagger$. □

7. Recurrent sequences. Let $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d)$ denote the eigenvalue array of a TB tridiagonal system over \mathbb{F} . Later in the paper we will show that there exists $\beta \in \mathbb{F}$ such that

$$\theta_{i-1} - \beta\theta_i + \theta_{i+1} = 0, \quad \theta_{i-1}^* - \beta\theta_i^* + \theta_{i+1}^* = 0 \quad (1 \leq i \leq d-1).$$

In this section, we have some comments about the above recurrence. For the rest of this section, fix an integer $d \geq 0$ and let $\{\sigma_i\}_{i=0}^d$ denote a sequence of scalars taken from \mathbb{F} .

DEFINITION 7.1. *For $\beta \in \mathbb{F}$, the sequence $\{\sigma_i\}_{i=0}^d$ is said to be β -recurrent whenever*

$$(7.39) \quad \sigma_{i-1} - \beta\sigma_i + \sigma_{i+1} = 0 \quad (1 \leq i \leq d-1).$$

We say that $\{\sigma_i\}_{i=0}^d$ is recurrent whenever it is β -recurrent for some $\beta \in \mathbb{F}$.

LEMMA 7.2. *For $\beta \in \mathbb{F}$ the following hold.*

(i) *Assume that $\{\sigma_i\}_{i=0}^d$ is β -recurrent. Then there exists $\varrho \in \mathbb{F}$ such that*

$$(7.40) \quad \varrho = \sigma_{i-1}^2 - \beta\sigma_{i-1}\sigma_i + \sigma_i^2 \quad (1 \leq i \leq d).$$

(ii) *Assume that there exists $\varrho \in \mathbb{F}$ that satisfies (7.40). Also assume that $\sigma_{i-1} \neq \sigma_{i+1}$ for $1 \leq i \leq d-1$. Then $\{\sigma_i\}_{i=0}^d$ is β -recurrent.*

Proof. For $1 \leq i \leq d$ define

$$S_i = \sigma_{i-1}^2 - \beta\sigma_{i-1}\sigma_i + \sigma_i^2.$$

Observe that for $1 \leq i \leq d-1$,

$$(7.41) \quad S_i - S_{i+1} = (\sigma_{i-1} - \sigma_{i+1})(\sigma_{i-1} - \beta\sigma_i + \sigma_{i+1}).$$

(i) In (7.41), the right-hand side is zero by (7.39). So $S_i - S_{i+1} = 0$. Thus, S_i is independent of i for $1 \leq i \leq d$.

(ii) In (7.41), the left-hand side is zero. In the right-hand side, the first factor is nonzero so the second factor is zero. Consequently $\{\sigma_i\}_{i=0}^d$ is β -recurrent. \square

8. The Askey–Wilson relations. Let A, A^* denote a TB tridiagonal pair over \mathbb{F} . In this section, we show that there exist scalars $\beta, \varrho, \varrho^*$ in \mathbb{F} such that both

$$(8.42) \quad A^2A^* - \beta AA^*A + A^*A^2 = \varrho A^*,$$

$$(8.43) \quad A^{*2}A - \beta A^*AA^* + AA^{*2} = \varrho^* A.$$

The equations (8.42), (8.43) are special cases of the Askey–Wilson relations [154, 166].

For the rest of this section, fix an integer $d \geq 0$, let V denote a vector space over \mathbb{F} with dimension $d + 1$, and let $\Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d)$ denote a TB tridiagonal system on V with eigenvalue array $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d)$.

First, we explain how (8.42), (8.43) are related to the recurrence discussed in Lemma 7.2.

LEMMA 8.1. *For $\beta, \varrho \in \mathbb{F}$ the following are equivalent:*

- (i) $\theta_{i-1}^2 - \beta\theta_{i-1}\theta_i + \theta_i^2 = \varrho \quad (1 \leq i \leq d)$;
- (ii) $A^2A^* - \beta AA^*A + A^*A^2 = \varrho A^*$.

Proof. Define $D = A^2A^* - \beta AA^*A + A^*A^2 - \varrho A^*$. Using $I = \sum_{i=0}^d E_i$, we obtain

$$D = IDI = \sum_{0 \leq i, j \leq d} E_i D E_j.$$

Thus $D = 0$ if and only if $E_i D E_j = 0$ ($0 \leq i, j \leq d$). Pick integers i, j such that $0 \leq i, j \leq d$. Using $E_i A = \theta_i E_i$ and $A E_j = \theta_j E_j$, we find that

$$E_i D E_j = E_i A^* E_j (\theta_i^2 - \beta \theta_i \theta_j + \theta_j^2 - \varrho).$$

By (3.13), $E_i A^* E_j \neq 0$ if and only if $|i - j| = 1$. The result follows from these comments. \square

Our next goal is to prove (8.42), (8.43); this is accomplished in Proposition 8.23.

LEMMA 8.2. *The following hold.*

- (i) *The elements A, A^* generate $\text{End}(V)$.*
- (ii) *For $0 \leq i, j, r \leq d$,*

$$E_i^* A^r E_j^* = \begin{cases} 0 & \text{if } |i - j| > r, \\ \neq 0 & \text{if } |i - j| = r. \end{cases}$$

Proof. Let $\{v_i\}_{i=0}^d$ denote a Φ -standard basis for V . For $X \in \text{End}(V)$ let X^\natural denote the matrix in $\text{Mat}_{d+1}(\mathbb{F})$ that represents X with respect to $\{v_i\}_{i=0}^d$. The matrices $(E_i^*)^\natural, (A^*)^\natural, A^\natural$ are given in (5.21)–(5.23). Now (i) follows from Lemma 6.5(ii) and (ii) follows from Lemma 6.1. \square

LEMMA 8.3. For $0 \leq i, j, k, r, s \leq d$ consider the expression

$$(8.44) \quad E_i^* A^r E_k^* A^s E_j^*.$$

- (i) Assume that $|i - j| > r + s$. Then (8.44) is zero.
- (ii) Assume that $i - j = r + s$. Then (8.44) is nonzero if and only if $k = j + s$.
- (iii) Assume that $j - i = r + s$. Then (8.44) is nonzero if and only if $k = i + r$.

Proof. Routine verification using Lemma 8.2(ii). □

LEMMA 8.4. For $0 \leq i, j, r, s \leq d$,

$$(8.45) \quad E_i^* A^r A^* A^s E_j^* = \begin{cases} 0 & \text{if } |i - j| > r + s, \\ \theta_{j+s}^* E_i^* A^{r+s} E_j^* & \text{if } i - j = r + s, \\ \theta_{i+r}^* E_i^* A^{r+s} E_j^* & \text{if } j - i = r + s. \end{cases}$$

Proof. Using $A^* = \sum_{k=0}^d \theta_k^* E_k^*$,

$$E_i^* A^r A^* A^s E_j^* = \sum_{k=0}^d \theta_k^* E_i^* A^r E_k^* A^s E_j^*.$$

Using $I = \sum_{k=0}^d E_k^*$,

$$E_i^* A^{r+s} E_j^* = E_i^* A^r I A^s E_j^* = \sum_{k=0}^d E_i^* A^r E_k^* A^s E_j^*.$$

By these comments and Lemma 8.3, we obtain the result. □

Let M (resp. M^*) denote the subalgebra of $\text{End}(V)$ generated by A (resp. A^*). Recall that each of $\{A^i\}_{i=0}^d$ and $\{E_i\}_{i=0}^d$ (resp. $\{A^{*i}\}_{i=0}^d$ and $\{E_i^*\}_{i=0}^d$) is a basis for the \mathbb{F} -vector space M (resp. M^*).

LEMMA 8.5. There exists a unique antiautomorphism \dagger of $\text{End}(V)$ that fixes each of A and A^* . The map \dagger fixes each element of M and each element of M^* . In particular, \dagger fixes E_i and E_i^* for $0 \leq i \leq d$. Moreover, $(X^\dagger)^\dagger = X$ for all $X \in \text{End}(V)$.

Proof. Follows from Lemmas 6.10 and 6.11. □

DEFINITION 8.6. Define a subspace MA^*M of $\text{End}(V)$ by

$$MA^*M = \text{Span}\{XA^*Y \mid X, Y \in M\}.$$

LEMMA 8.7. The following elements form a basis for the \mathbb{F} -vector space MA^*M :

$$(8.46) \quad \{E_{i-1}A^*E_i, E_iA^*E_{i-1} \mid 1 \leq i \leq d\}.$$

Moreover $\dim(MA^*M) = 2d$.

Proof. We assume $d \geq 1$; otherwise the result follows from Lemma 3.6. The \mathbb{F} -vector space MA^*M is spanned by $\{E_iA^*E_j \mid 0 \leq i, j \leq d\}$. By (3.13) we have $E_iA^*E_j = 0$ if $|i - j| \neq 1$ ($0 \leq i, j \leq d$). By these comments, the elements (8.46) span MA^*M . We show that the elements (8.46) are linearly independent. Suppose that there exist scalars $\{\alpha_i\}_{i=1}^d, \{\beta_i\}_{i=1}^d$ in \mathbb{F} such that

$$(8.47) \quad 0 = \sum_{i=1}^d \alpha_i E_{i-1}A^*E_i + \sum_{i=1}^d \beta_i E_iA^*E_{i-1}.$$

Pick any integer j ($1 \leq j \leq d$). We show that $\alpha_j = 0$. In (8.47), multiply each side on the left by E_{j-1} and right by E_j . This gives $\alpha_j E_{j-1} A^* E_j = 0$. This forces $\alpha_j = 0$ since $E_{j-1} A^* E_j \neq 0$ by (3.13). A similar argument shows that $\beta_j = 0$. We have shown that the elements (8.46) are linearly independent. By the above comments, the elements (8.46) form a basis for the \mathbb{F} -vector space MA^*M . The set (8.46) has cardinality $2d$ so $\dim(MA^*M) = 2d$. \square

LEMMA 8.8. For $0 \leq i \leq d$,

$$(8.48) \quad E_i A^* = E_i A^* E_{i-1} + E_i A^* E_{i+1}.$$

Proof. Using $I = \sum_{j=0}^d E_j$,

$$E_i A^* = E_i A^* I = \sum_{j=0}^d E_i A^* E_j.$$

By (3.13), $E_i A^* E_j = 0$ if $|i - j| \neq 1$ ($0 \leq j \leq d$). The result follows. \square

LEMMA 8.9. For $0 \leq i \leq d$,

$$A^* E_i = E_{i-1} A^* E_i + E_{i+1} A^* E_i.$$

Proof. Apply the antiautomorphism \dagger to each side of (8.48). \square

LEMMA 8.10. For $1 \leq i \leq d$,

$$\begin{aligned} E_{i-1} A^* E_i &= E_{i-1} A^* - A^* E_{i-2} + E_{i-3} A^* - A^* E_{i-4} + \cdots \\ E_i A^* E_{i-1} &= A^* E_{i-1} - E_{i-2} A^* + A^* E_{i-3} - E_{i-4} A^* + \cdots \end{aligned}$$

Moreover

$$(8.49) \quad \sum_{\substack{0 \leq i \leq d \\ i \text{ even}}} E_i A^* = \sum_{\substack{0 \leq i \leq d \\ i \text{ odd}}} A^* E_i, \quad \sum_{\substack{0 \leq i \leq d \\ i \text{ odd}}} E_i A^* = \sum_{\substack{0 \leq i \leq d \\ i \text{ even}}} A^* E_i.$$

Proof. Solve the equations in Lemmas 8.8 and 8.9. \square

LEMMA 8.11. We have $\sum_{i=0}^d (-1)^i (E_i A^* + A^* E_i) = 0$.

Proof. Follows from (8.49). \square

LEMMA 8.12. The element A^* is equal to each of the following sums:

$$\sum_{\substack{0 \leq i \leq d \\ i \text{ even}}} (E_i A^* + A^* E_i), \quad \sum_{\substack{0 \leq i \leq d \\ i \text{ odd}}} (E_i A^* + A^* E_i).$$

Proof. We have $A^* = \sum_{i=0}^d E_i A^*$. Evaluate this sum using (8.49). \square

LEMMA 8.13. The following elements form a basis for the \mathbb{F} -vector space MA^*M :

$$(8.50) \quad \{E_i A^*, A^* E_i \mid 0 \leq i \leq d - 1\}.$$

Proof. Let \mathcal{X} denote the subspace of $\text{End}(V)$ spanned by (8.50). By construction $\mathcal{X} \subseteq MA^*M$. By Lemmas 8.7 and 8.10, $MA^*M \subseteq \mathcal{X}$. So $MA^*M = \mathcal{X}$. By Lemma 8.7, the dimension of MA^*M is $2d$. The set (8.50) has cardinality $2d$. Therefore, the elements (8.50) form a basis for the \mathbb{F} -vector space MA^*M . \square

COROLLARY 8.14. We have $MA^*M = MA^* + A^*M$.

Proof. By construction $MA^* + A^*M \subseteq MA^*M$. By Lemma 8.13, $MA^*M \subseteq MA^* + A^*M$. \square

DEFINITION 8.15. Define a subspace $(MA^*M)^{\text{sym}}$ of $\text{End}(V)$ by

$$(MA^*M)^{\text{sym}} = \{Z \in MA^*M \mid Z^\dagger = Z\}.$$

We call $(MA^*M)^{\text{sym}}$ the symmetric part of MA^*M .

LEMMA 8.16. The subspace $(MA^*M)^{\text{sym}}$ contains XA^*X for all X in M . Moreover, $(MA^*M)^{\text{sym}}$ contains $XA^*Y + YA^*X$ for all X, Y in M .

Proof. Observe that $(XA^*Y)^\dagger = YA^*X$ for all $X, Y \in M$. \square

LEMMA 8.17. The following elements form a basis for the \mathbb{F} -vector space $(MA^*M)^{\text{sym}}$:

$$(8.51) \quad \{E_{i-1}A^*E_i + E_iA^*E_{i-1} \mid 1 \leq i \leq d\}.$$

Moreover, $\dim(MA^*M)^{\text{sym}} = d$.

Proof. Let \mathcal{Z} denote the subspace of $\text{End}(V)$ spanned by (8.51). We show that $\mathcal{Z} = (MA^*M)^{\text{sym}}$. By Lemma 8.16, we have $\mathcal{Z} \subseteq (MA^*M)^{\text{sym}}$. To show the reverse inclusion, pick any $Z \in (MA^*M)^{\text{sym}}$. By construction $Z^\dagger = Z$. By Lemma 8.7, there exist scalars $\{\alpha_i\}_{i=1}^d, \{\beta_i\}_{i=1}^d$ in \mathbb{F} such that

$$Z = \sum_{i=1}^d \alpha_i E_{i-1}A^*E_i + \sum_{i=1}^d \beta_i E_iA^*E_{i-1}.$$

We have

$$Z^\dagger = \sum_{i=1}^d \alpha_i E_iA^*E_{i-1} + \sum_{i=1}^d \beta_i E_{i-1}A^*E_i.$$

By $Z^\dagger = Z$ and the above two equations, we find that $\alpha_i = \beta_i$ for $1 \leq i \leq d$. Thus,

$$Z = \sum_{i=1}^d \alpha_i (E_{i-1}A^*E_i + E_iA^*E_{i-1}),$$

and so $Z \in \mathcal{Z}$. We have shown that $\mathcal{Z} = (MA^*M)^{\text{sym}}$. Therefore, the elements (8.51) span $(MA^*M)^{\text{sym}}$. Using Lemma 8.7 one checks that the elements (8.51) are linearly independent. The result follows. \square

LEMMA 8.18. The following elements form a basis for the \mathbb{F} -vector space $(MA^*M)^{\text{sym}}$:

$$(8.52) \quad \{E_iA^* + A^*E_i \mid 0 \leq i \leq d-1\}.$$

Proof. Each of the elements in (8.52) is fixed by \dagger , and therefore contained in $(MA^*M)^{\text{sym}}$. The elements (8.52) are linearly independent by Lemma 8.13. The set (8.52) has cardinality d . The subspace $(MA^*M)^{\text{sym}}$ has dimension d by Lemma 8.17. The result follows from these comments. \square

LEMMA 8.19. We have

$$(8.53) \quad (MA^*M)^{\text{sym}} = \{XA^* + A^*X \mid X \in M\}.$$

Proof. The inclusion \subseteq follows from Lemma 8.18. The inclusion \supseteq follows from Lemma 8.16. \square

LEMMA 8.20. The \mathbb{F} -vector space $(MA^*M)^{\text{sym}}$ is spanned by

$$(8.54) \quad \{A^*, AA^* + A^*A, A^2A^* + A^*A^2, \dots, A^dA^* + A^*A^d\}.$$

Proof. Let \mathcal{Z} denote the subspace of $\text{End}(V)$ spanned by (8.54). We show that $\mathcal{Z} = (MA^*M)^{\text{sym}}$. We have $\mathcal{Z} \subseteq (MA^*M)^{\text{sym}}$ by Lemma 8.16. By Lemma 8.19,

$$(MA^*M)^{\text{sym}} = \text{Span}\{A^i A^* + A^* A^i \mid 0 \leq i \leq d\} \subseteq \mathcal{Z}.$$

The result follows. □

By Lemma 8.17 the \mathbb{F} -vector space $(MA^*M)^{\text{sym}}$ has dimension d . The set (8.54) contains $d+1$ elements. So the vectors (8.54) are linearly dependent. We now find the dependency. To avoid trivialities we assume that $d \geq 1$.

LEMMA 8.21. *Assume that $d \geq 1$. Then the following (i)–(iii) hold.*

- (i) $\text{Char}(\mathbb{F}) \neq 2$.
- (ii) *There exists a unique integer n ($1 \leq n \leq d$) such that $\theta_n^* = -\theta_0^*$.*
- (iii) *The element $A^n A^* + A^* A^n$ is contained in the span of*

$$(8.55) \quad \{A^*, AA^* + A^*A, A^2A^* + A^*A^2, \dots, A^{n-1}A^* + A^*A^{n-1}\}.$$

Proof. Since the vectors (8.54) are linearly dependent, there exist scalars $\{\alpha_i\}_{i=0}^d$ in \mathbb{F} , not all zero, such that

$$\alpha_0 A^* + \sum_{i=1}^d \alpha_i (A^i A^* + A^* A^i) = 0.$$

Define $n = \max\{i \mid 0 \leq i \leq d, \alpha_i \neq 0\}$. Note that $n \geq 1$, since $A^* \neq 0$ by Lemma 3.6. So

$$(8.56) \quad \alpha_0 A^* + \sum_{i=1}^n \alpha_i (A^i A^* + A^* A^i) = 0, \quad \alpha_n \neq 0.$$

By (8.56) the element $A^n A^* + A^* A^n$ is contained in the span of (8.55). In (8.56), multiply each side on the left by E_0^* and right by E_n^* . Simplify the result using $E_0^* A^* = \theta_0^* E_0^*$, $A^* E_n^* = \theta_n^* E_n^*$, $E_0^* E_n^* = 0$ to get

$$(\theta_0^* + \theta_n^*) \sum_{i=1}^n \alpha_i E_0^* A^i E_n^* = 0.$$

By Lemma 8.2(ii), $E_0^* A^i E_n^* = 0$ for $1 \leq i \leq n-1$. So the above line becomes

$$\alpha_n (\theta_0^* + \theta_n^*) E_0^* A^n E_n^* = 0.$$

We have $\alpha_n \neq 0$, and $E_0^* A^n E_n^* \neq 0$ by Lemma 8.2(ii). Thus, $\theta_0^* + \theta_n^* = 0$, and so $\theta_n^* = -\theta_0^*$. The $\{\theta_i^*\}_{i=0}^d$ are mutually distinct, so no dual eigenvalue other than θ_n^* is equal to $-\theta_0^*$. We have $\text{Char}(\mathbb{F}) \neq 2$; otherwise $\theta_n^* = \theta_0^*$. The result follows. □

NOTE 8.22. Referring to Lemma 8.21(ii), it turns out that $n = d$; this will be established in Lemma 11.3.

PROPOSITION 8.23. *There exist $\beta, \varrho, \varrho^*$ in \mathbb{F} that satisfy both (8.42), (8.43).*

Proof. We first show that

$$(8.57) \quad AA^*A \in \text{Span}\{A^*, AA^* + A^*A, A^2A^* + A^*A^2\}.$$

We assume that $d \geq 3$; otherwise (8.57) holds by Lemma 8.20. Recall the integer n from Lemma 8.21. By Lemmas 8.20 and 8.21(iii), there exist scalars $\{\alpha_i\}_{i=0}^d$ in \mathbb{F} with $\alpha_n = 0$ such that

$$AA^*A = \alpha_0 A^* + \sum_{i=1}^d \alpha_i (A^i A^* + A^* A^i).$$

We show that $\alpha_i = 0$ for $3 \leq i \leq d$. Suppose not. Then there exists an integer t ($3 \leq t \leq d$) such that $\alpha_t \neq 0$. We choose t to be maximal. Then $t \neq n$ and

$$AA^*A = \alpha_0 A^* + \sum_{i=1}^t \alpha_i (A^i A^* + A^* A^i).$$

In the above line, multiply each side on the left by E_0^* and right by E_t^* . Simplify the result to get

$$E_0^* AA^* A E_t^* = (\theta_0^* + \theta_t^*) \sum_{i=1}^t \alpha_i E_0^* A^i E_t^*.$$

In the above line the left-hand side is zero by Lemma 8.4, and $E_0^* A^i E_t^* = 0$ ($1 \leq i \leq t-1$) by Lemma 8.2(ii). Thus,

$$(8.58) \quad 0 = \alpha_t (\theta_0^* + \theta_t^*) E_0^* A^t E_t^*.$$

We examine the factors in (8.58). By construction $\alpha_t \neq 0$. We have $\theta_0^* + \theta_t^* \neq 0$ by Lemma 8.21(ii) and $t \neq n$. Also $E_0^* A^t E_t^* \neq 0$ by Lemma 8.2(ii). Therefore, the right-hand side of (8.58) is nonzero, for a contradiction. We have shown (8.57).

Next we show that there exists $\beta \in \mathbb{F}$ such that

$$(8.59) \quad \theta_{i-1}^* - \beta \theta_i^* + \theta_{i+1}^* = 0 \quad (1 \leq i \leq d-1).$$

We assume that $d \geq 2$; otherwise the assertion is vacuous. By (8.57) there exist scalars $\alpha_0, \alpha_1, \alpha_2$ in \mathbb{F} such that

$$0 = \alpha_0 A^* + \alpha_1 (AA^* + A^*A) + \alpha_2 (A^2 A^* + A^* A^2) - AA^*A.$$

For $1 \leq i \leq d-1$, in the above line multiply each side on the left by E_{i-1}^* and right by E_{i+1}^* . Simplify the result using Lemmas 8.2(ii) and 8.4 to get

$$0 = E_{i-1}^* A^2 E_{i+1}^* (\alpha_2 \theta_{i+1}^* + \alpha_2 \theta_{i-1}^* - \theta_i^*). \quad (1 \leq i \leq d-1).$$

We have $E_{i-1}^* A^2 E_{i+1}^* \neq 0$ by Lemma 8.2(ii). Thus,

$$\alpha_2 (\theta_{i-1}^* + \theta_{i+1}^*) = \theta_i^* \quad (1 \leq i \leq d-1).$$

Assume for the moment that $\alpha_2 \neq 0$. Then (8.59) holds for $\beta = \alpha_2^{-1}$. Next assume that $\alpha_2 = 0$. Then $\theta_i^* = 0$ for $1 \leq i \leq d-1$. This forces $d = 2$ and $\theta_1^* = 0$. In this case $\theta_2^* = -\theta_0^*$ by Lemma 8.21(ii) and (8.59) holds for any $\beta \in \mathbb{F}$. We have shown that there exists $\beta \in \mathbb{F}$ that satisfies (8.59).

By (8.59) and Lemma 7.2(i) there exists $\varrho^* \in \mathbb{F}$ such that

$$\theta_{i-1}^{*2} - \beta \theta_{i-1}^* \theta_i^* + \theta_i^{*2} = \varrho^* \quad (1 \leq i \leq d).$$

By this and Lemma 8.1 (applied to Φ^*) we obtain (8.43). For $1 \leq i \leq d-1$, multiply each side of (8.43) on the left by E_{i-1} and right by E_{i+1} . Simplify the result using Lemmas 8.2(ii) and 8.4 (applied to Φ^*) to get

$$0 = E_{i-1} A^{*2} E_{i+1} (\theta_{i-1} - \beta \theta_i + \theta_{i+1}) \quad (1 \leq i \leq d-1).$$

Applying Lemma 8.2(ii) to Φ^* , we obtain $E_{i-1}A^{*2}E_{i+1} \neq 0$. Thus,

$$\theta_{i-1} - \beta\theta_i + \theta_{i+1} = 0 \quad (1 \leq i \leq d-1).$$

By this and Lemma 7.2(i), there exists $\varrho \in \mathbb{F}$ such that

$$\theta_{i-1}^2 - \beta\theta_{i-1}\theta_i + \theta_i^2 = \varrho \quad (1 \leq i \leq d).$$

By this and Lemma 8.1 we obtain (8.42). □

We mention some results for later use.

LEMMA 8.24. *For $X \in \text{End}(V)$ the following are equivalent:*

- (i) X commutes with each of A, A^* ;
- (ii) there exists $\lambda \in \mathbb{F}$ such that $X = \lambda I$.

Proof. (i) \Rightarrow (ii) By Lemma 8.2(i) the element X is contained in the center of $\text{End}(V)$. The center of $\text{End}(V)$ is spanned by I . The result follows.

(ii) \Rightarrow (i) Clear. □

The next two results follow from Lemma 8.24.

COROLLARY 8.25. *For $\psi \in \text{End}(V)$ the following are equivalent:*

- (i) ψ is an isomorphism of TB tridiagonal pairs from A, A^* to A, A^* ;
- (ii) there exists nonzero $\lambda \in \mathbb{F}$ such that $\psi = \lambda I$.

COROLLARY 8.26. *For $\psi \in \text{End}(V)$ the following are equivalent:*

- (i) ψ is an isomorphism of TB tridiagonal systems from Φ to Φ ;
- (ii) there exists nonzero $\lambda \in \mathbb{F}$ such that $\psi = \lambda I$.

9. The Askey–Wilson sequence and the fundamental parameter. In this section, we introduce the notion of an Askey–Wilson sequence and fundamental parameter for TB tridiagonal pairs and systems.

Throughout this section let A, A^* denote a TB tridiagonal pair over \mathbb{F} . Let Φ denote an associated TB tridiagonal system.

DEFINITION 9.1. *By an Askey–Wilson sequence for A, A^* , we mean a sequence $\beta, \varrho, \varrho^*$ of scalars in \mathbb{F} that satisfy (8.42) and (8.43).*

DEFINITION 9.2. *By an Askey–Wilson sequence for Φ , we mean an Askey–Wilson sequence for A, A^* .*

LEMMA 9.3. *The following are the same:*

- (i) an Askey–Wilson sequence for Φ ;
- (ii) an Askey–Wilson sequence for Φ^\downarrow ;
- (iii) an Askey–Wilson sequence for Φ^\updownarrow .

Proof. The TB tridiagonal systems $\Phi, \Phi^\downarrow, \Phi^\updownarrow$ have the same associated TB tridiagonal pair A, A^* . The result follows. □

Let $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d)$ denote the eigenvalue array of Φ .

PROPOSITION 9.4. *Let $\beta, \varrho, \varrho^*$ denote a sequence of scalars taken from \mathbb{F} . This sequence is an Askey–Wilson sequence for Φ if and only if both*

$$(9.60) \quad \theta_{i-1}^2 - \beta\theta_{i-1}\theta_i + \theta_i^2 = \varrho \quad (1 \leq i \leq d),$$

$$(9.61) \quad \theta_{i-1}^{*2} - \beta\theta_{i-1}^*\theta_i^* + \theta_i^{*2} = \varrho^* \quad (1 \leq i \leq d).$$

Proof. Apply Lemma 8.1 to Φ and Φ^* . □

LEMMA 9.5. *Let $\beta, \varrho, \varrho^*$ denote an Askey–Wilson sequence for A, A^* . Then $\beta, \varrho^*, \varrho$ is an Askey–Wilson sequence for A^*, A .*

Proof. By (8.42) and (8.43). □

LEMMA 9.6. *Let $\beta, \varrho, \varrho^*$ denote an Askey–Wilson sequence for A, A^* . Then for nonzero $\zeta, \zeta^* \in \mathbb{F}$ the TB tridiagonal pair $\zeta A, \zeta^* A^*$ has an Askey–Wilson sequence $\beta, \zeta^2 \varrho, \zeta^{*2} \varrho^*$.*

Proof. One verifies that $\beta, \zeta^2 \varrho, \zeta^{*2} \varrho^*$ satisfy the Askey–Wilson relations for $\zeta A, \zeta^* A^*$. □

DEFINITION 9.7. *A scalar $\beta \in \mathbb{F}$ is called a fundamental parameter for A, A^* whenever there exist $\varrho, \varrho^* \in \mathbb{F}$ such that $\beta, \varrho, \varrho^*$ is an Askey–Wilson sequence for A, A^* .*

DEFINITION 9.8. *By a fundamental parameter for Φ , we mean a fundamental parameter for A, A^* .*

LEMMA 9.9. *Let β denote a fundamental parameter for Φ . Then β is a fundamental parameter for each of $\Phi, \Phi^\downarrow, \Phi^\uparrow, \Phi^*$.*

Proof. By Lemmas 9.3 and 9.5. □

PROPOSITION 9.10. *A scalar $\beta \in \mathbb{F}$ is a fundamental parameter for Φ if and only if both*

$$(9.62) \quad \theta_{i-1} - \beta\theta_i + \theta_{i+1} = 0 \quad (1 \leq i \leq d-1),$$

$$(9.63) \quad \theta_{i-1}^* - \beta\theta_i^* + \theta_{i+1}^* = 0 \quad (1 \leq i \leq d-1).$$

Proof. By Lemma 7.2 and Proposition 9.4. □

LEMMA 9.11. *Let β denote a fundamental parameter for A, A^* . Then β is a fundamental parameter for A^*, A .*

Proof. By Lemma 9.5 or Lemma 9.9. □

LEMMA 9.12. *Let β denote a fundamental parameter for A, A^* . Then β is a fundamental parameter for $\zeta A, \zeta^* A^*$ for all nonzero $\zeta, \zeta^* \in \mathbb{F}$.*

Proof. By Lemma 9.6. □

The uniqueness of the Askey–Wilson sequence and fundamental parameter will be discussed in Lemmas 12.6, 12.7.

10. More on recurrent sequences. Recall the recurrent sequences from Definition 7.1. In this section, we obtain a detailed description of these sequences. In our description, two special cases come up, said to be symmetric and antisymmetric. Throughout this section fix an integer $d \geq 1$.

DEFINITION 10.1. *Let $\{\sigma_i\}_{i=0}^d$ denote a sequence of scalars taken from \mathbb{F} . This sequence is said to be over \mathbb{F} and have diameter d . The sequence $\{\sigma_i\}_{i=0}^d$ is said to be symmetric whenever $\sigma_i = \sigma_{d-i}$ for $0 \leq i \leq d$. The sequence $\{\sigma_i\}_{i=0}^d$ is said to be antisymmetric whenever $\sigma_i + \sigma_{d-i} = 0$ for $0 \leq i \leq d$.*

For the rest of this section fix $\beta \in \mathbb{F}$.

DEFINITION 10.2. Let \mathcal{R} denote the set of β -recurrent sequences over \mathbb{F} that have diameter d . Let \mathcal{R}^{sym} (resp. $\mathcal{R}^{\text{asym}}$) denote the set of symmetric (resp. antisymmetric) elements in \mathcal{R} . Note that \mathcal{R} is a subspace of the \mathbb{F} -vector space \mathbb{F}^{d+1} . Moreover, \mathcal{R}^{sym} and $\mathcal{R}^{\text{asym}}$ are subspaces of \mathcal{R} .

LEMMA 10.3. If $\text{Char}(\mathbb{F}) = 2$ then $\mathcal{R}^{\text{sym}} = \mathcal{R}^{\text{asym}}$. If $\text{Char}(\mathbb{F}) \neq 2$ then

$$\mathcal{R} = \mathcal{R}^{\text{sym}} + \mathcal{R}^{\text{asym}} \quad (\text{direct sum}).$$

Proof. The first assertion is clear. Concerning the second assertion, assume that $\text{Char}(\mathbb{F}) \neq 2$. For a sequence $\{\sigma_i\}_{i=0}^d$ in \mathcal{R} define

$$\sigma_i^+ = \frac{\sigma_i + \sigma_{d-i}}{2}, \quad \sigma_i^- = \frac{\sigma_i - \sigma_{d-i}}{2} \quad (0 \leq i \leq d).$$

We have $\sigma_i = \sigma_i^+ + \sigma_i^-$ for $0 \leq i \leq d$. The sequence $\{\sigma_i^+\}_{i=0}^d$ is β -recurrent and symmetric. The sequence $\{\sigma_i^-\}_{i=0}^d$ is β -recurrent and antisymmetric. By these comments $\mathcal{R} = \mathcal{R}^{\text{sym}} + \mathcal{R}^{\text{asym}}$. We show that this sum is direct. Pick an element $\{\sigma_i\}_{i=0}^d$ of $\mathcal{R}^{\text{sym}} \cap \mathcal{R}^{\text{asym}}$. For $0 \leq i \leq d$ we have both $\sigma_{d-i} = \sigma_i$ and $\sigma_{d-i} = -\sigma_i$, and so $\sigma_i = 0$. Therefore, $\mathcal{R}^{\text{sym}} \cap \mathcal{R}^{\text{asym}} = 0$ and consequently the sum $\mathcal{R}^{\text{sym}} + \mathcal{R}^{\text{asym}}$ is direct. \square

Our next goal is to display a basis for \mathcal{R}^{sym} and $\mathcal{R}^{\text{asym}}$, under the assumption that $\text{Char}(\mathbb{F}) \neq 2$. Our strategy is to first display some nonzero elements in \mathcal{R}^{sym} and $\mathcal{R}^{\text{asym}}$, and a bit later show that these elements form a basis. The cases $\beta = 2$, $\beta = -2$ will be handled separately.

LEMMA 10.4. Assume that $\text{Char}(\mathbb{F}) \neq 2$ and $\beta = \pm 2$. For $0 \leq i \leq d$ define σ_i as follows:

Case	σ_i
$\beta = 2$	1
$\beta = -2, d \text{ even}$	$(-1)^i$
$\beta = -2, d \text{ odd}$	$(d - 2i)(-1)^i$

Then $\{\sigma_i\}_{i=0}^d$ is nonzero and contained in \mathcal{R}^{sym} .

Proof. One routinely checks that $\{\sigma_i\}_{i=0}^d$ is nonzero, β -recurrent, and symmetric. The result follows. \square

LEMMA 10.5. Assume that $\text{Char}(\mathbb{F}) \neq 2$ and $\beta = \pm 2$. For $0 \leq i \leq d$ define σ_i as follows:

Case	σ_i
$\beta = 2$	$d - 2i$
$\beta = -2, d \text{ even}$	$(d - 2i)(-1)^i$
$\beta = -2, d \text{ odd}$	$(-1)^i$

Then $\{\sigma_i\}_{i=0}^d$ is nonzero and contained in $\mathcal{R}^{\text{asym}}$.

Proof. One routinely checks that $\{\sigma_i\}_{i=0}^d$ is nonzero, β -recurrent, and antisymmetric. The result follows. \square

LEMMA 10.6. Assume that $\text{Char}(\mathbb{F}) \neq 2$ and $\beta \neq \pm 2$. Let $q \in \overline{\mathbb{F}}$ be such that $\beta = q^2 + q^{-2}$. For $0 \leq i \leq d$ define σ_i by

$$\sigma_i = \begin{cases} q^{d-2i} + q^{2i-d} & \text{if } d \text{ is even,} \\ \frac{q^{d-2i} + q^{2i-d}}{q + q^{-1}} & \text{if } d \text{ is odd.} \end{cases}$$

Then $\{\sigma_i\}_{i=0}^d$ is nonzero and contained in \mathcal{R}^{sym} .

Proof. Clearly $\{\sigma_i\}_{i=0}^d$ is nonzero. We verify that $\{\sigma_i\}_{i=0}^d$ is contained in \mathcal{R}^{sym} . One routinely checks that $\{\sigma_i\}_{i=0}^d$ is β -recurrent and symmetric. We show that $\sigma_i \in \mathbb{F}$ for $0 \leq i \leq d$. First assume that d is even. Observe that $\sigma_{d/2} = 2$ and $\sigma_{d/2-1} = q^2 + q^{-2} = \beta$. Therefore, each of $\sigma_{d/2}, \sigma_{d/2-1}$ is contained in \mathbb{F} . By this and since $\{\sigma_i\}_{i=0}^d$ is β -recurrent, we obtain $\sigma_i \in \mathbb{F}$ for $0 \leq i \leq d$. Next assume that d is odd. Observe that $\sigma_{(d-1)/2} = 1$ and $\sigma_{(d+1)/2} = 1$. Therefore, each of $\sigma_{(d-1)/2}, \sigma_{(d+1)/2}$ is contained in \mathbb{F} . By this and since $\{\sigma_i\}_{i=0}^d$ is β -recurrent, we obtain $\sigma_i \in \mathbb{F}$ for $0 \leq i \leq d$. Thus, $\{\sigma_i\}_{i=0}^d$ is contained in \mathcal{R}^{sym} . The result follows. \square

LEMMA 10.7. Assume that $\text{Char}(\mathbb{F}) \neq 2$ and $\beta \neq \pm 2$. Let $q \in \overline{\mathbb{F}}$ be such that $\beta = q^2 + q^{-2}$. For $0 \leq i \leq d$ define σ_i by

$$\sigma_i = \begin{cases} \frac{q^{d-2i} - q^{2i-d}}{q^2 - q^{-2}} & \text{if } d \text{ is even,} \\ \frac{q^{d-2i} - q^{2i-d}}{q - q^{-1}} & \text{if } d \text{ is odd.} \end{cases}$$

Then $\{\sigma_i\}_{i=0}^d$ is nonzero and contained in $\mathcal{R}^{\text{asym}}$.

Proof. Clearly $\{\sigma_i\}_{i=0}^d$ is nonzero. We verify that $\{\sigma_i\}_{i=0}^d$ is contained in $\mathcal{R}^{\text{asym}}$. One routinely checks that $\{\sigma_i\}_{i=0}^d$ is β -recurrent and antisymmetric. We show that $\sigma_i \in \mathbb{F}$ for $0 \leq i \leq d$. First assume that d is even. Observe that $\sigma_{d/2} = 0$ and $\sigma_{d/2-1} = 1$. Therefore, each of $\sigma_{d/2}, \sigma_{d/2-1}$ is contained in \mathbb{F} . By this and since $\{\sigma_i\}_{i=0}^d$ is β -recurrent, we obtain $\sigma_i \in \mathbb{F}$ for $0 \leq i \leq d$. Next assume that d is odd. Observe that $\sigma_{(d-1)/2} = 1$ and $\sigma_{(d+1)/2} = -1$. Therefore, each of $\sigma_{(d-1)/2}, \sigma_{(d+1)/2}$ is contained in \mathbb{F} . By this and since $\{\sigma_i\}_{i=0}^d$ is β -recurrent, we obtain $\sigma_i \in \mathbb{F}$ for $0 \leq i \leq d$. Thus, $\{\sigma_i\}_{i=0}^d$ is contained in $\mathcal{R}^{\text{asym}}$. The result follows. \square

NOTE 10.8. The parameter q in Lemmas 10.6, 10.7 is not uniquely determined by $\{\sigma_i\}_{i=0}^d$. To clarify the situation, let y denote an indeterminate, and consider the equation

$$(10.66) \quad \beta = y^2 + y^{-2}.$$

Let $0 \neq q \in \mathbb{F}$ denote a solution of (10.66). Then the solutions of (10.66) are $q, -q, q^{-1}, -q^{-1}$. In Lemmas 10.6 and 10.7, replacing q by $-q, q^{-1}, -q^{-1}$ does not change σ_i . Thus, the given sequence $\{\sigma_i\}_{i=0}^d$ depends only on d and β and not on the choice of q .

In Lemmas 10.6 and 10.7, each scalar σ_i can be represented as a polynomial in β . The polynomial has Chebychev type, as we now explain. Assume that $\text{Char}(\mathbb{F}) \neq 2$. For scalars a, b, c in \mathbb{F} define polynomials $T_0(x), T_1(x), T_2(x), \dots$ by

$$\begin{aligned} T_0(x) &= a, \\ T_1(x) &= bx + c, \\ T_i(x) &= 2xT_{i-1}(x) - T_{i-2}(x) \quad i = 2, 3, \dots \end{aligned}$$

Then $T_i(x)$ is the i th Chebychev polynomial with parameters a, b, c [11, Remark 2.5.3]. Now consider the sequence $\{\sigma_i\}_{i=0}^d$ from Lemma 10.6.

(i) Assume that d is even. Then for $n = d/2$ and $a = 2, b = 2, c = 0$,

$$\sigma_i = T_{n-i}(\beta/2) \quad (0 \leq i \leq n).$$

(ii) Assume that d is odd. Then for $n = (d - 1)/2$ and $a = 1, b = 2, c = -1$,

$$\sigma_i = T_{n-i}(\beta/2) \quad (0 \leq i \leq n).$$

Next consider the sequence $\{\sigma_i\}_{i=0}^d$ from Lemma 10.7.

(i) Assume that d is even. Then for $n = d/2$ and $a = 1, b = 2, c = 0$,

$$\sigma_i = T_{n-1-i}(\beta/2) \quad (0 \leq i \leq n - 1), \quad \sigma_n = 0.$$

(ii) Assume that d is odd. Then for $n = (d - 1)/2$ and $a = 1, b = 2, c = 1$,

$$\sigma_i = T_{n-i}(\beta/2) \quad (0 \leq i \leq n).$$

LEMMA 10.9. Assume that $\text{Char}(\mathbb{F}) \neq 2$. Then

$$\dim \mathcal{R} = 2, \quad \dim \mathcal{R}^{\text{sym}} = 1, \quad \dim \mathcal{R}^{\text{asym}} = 1.$$

Proof. Pick a sequence $\{\sigma_i\}_{i=0}^d$ in \mathcal{R} . By the β -recurrence, the scalars $\{\sigma_i\}_{i=0}^d$ are uniquely determined by σ_0, σ_1 . Thus, $\dim \mathcal{R} \leq 2$. The subspace \mathcal{R}^{sym} is nonzero by Lemmas 10.4 and 10.6. The subspace $\mathcal{R}^{\text{asym}}$ is nonzero by Lemmas 10.5 and 10.7. By these comments and Lemma 10.3, we obtain the results. \square

COROLLARY 10.10. The following hold.

- (i) Assume that $\beta = \pm 2$. Then the sequence $\{\sigma_i\}_{i=0}^d$ from Lemma 10.4 (resp. Lemma 10.5) is a basis for \mathcal{R}^{sym} (resp. $\mathcal{R}^{\text{asym}}$).
- (ii) Assume that $\beta \neq \pm 2$. Then the sequence $\{\sigma_i\}_{i=0}^d$ from Lemma 10.6 (resp. Lemma 10.7) is a basis for \mathcal{R}^{sym} (resp. $\mathcal{R}^{\text{asym}}$).

Proof. By Lemma 10.9. \square

For a sequence $\{\sigma_i\}_{i=0}^d$ in \mathcal{R} , we sometimes consider the case in which $\{\sigma_i\}_{i=0}^d$ are mutually distinct. Of course this does not happen if $\{\sigma_i\}_{i=0}^d$ is symmetric. But it could happen if $\{\sigma_i\}_{i=0}^d$ is antisymmetric. We now give the details.

DEFINITION 10.11. A sequence $\{\sigma_i\}_{i=0}^d$ in \mathcal{R} is said to be MutDist whenever $\sigma_0, \sigma_1, \dots, \sigma_d$ are mutually distinct.

DEFINITION 10.12. A subspace of \mathcal{R} is said to be MutDist whenever the subspace is nonzero and every nonzero element of the subspace is MutDist.

LEMMA 10.13. The following (i), (ii) are equivalent:

- (i) $\mathcal{R}^{\text{asym}}$ is MutDist;
- (ii) $\mathcal{R}^{\text{asym}}$ contains a MutDist element.

Assume that (i), (ii) hold. Then $\text{Char}(\mathbb{F}) \neq 2$.

Proof. First assume that (i) holds. Then (ii) follows from Definition 10.12. Next assume that (ii) holds. Let $\{\sigma_i\}_{i=0}^d$ denote a MutDist element in $\mathcal{R}^{\text{asym}}$. Then $\sigma_d = -\sigma_0$ and $\sigma_d \neq \sigma_0$. Thus, $\text{Char}(\mathbb{F}) \neq 2$. By this and Lemma 10.9, the space $\mathcal{R}^{\text{asym}}$ has dimension 1. Now (i) follows. Next assume that (i), (ii) hold. We mentioned in the proof of (ii) \Rightarrow (i) that $\text{Char}(\mathbb{F}) \neq 2$. \square

LEMMA 10.14. Assume that $d \geq 2$ and $\beta = -2$. Assume that $\mathcal{R}^{\text{asym}}$ is MutDist. Then d is even.

Proof. By Lemma 10.13, $\text{Char}(\mathbb{F}) \neq 2$. Let $\{\sigma_i\}_{i=0}^d$ be from Lemma 10.5. Then $\{\sigma_i\}_{i=0}^d$ is MutDist. If d is odd, then $\sigma_0 = \sigma_2$, for a contradiction. Thus, d is even. \square

LEMMA 10.15. *Let the sequence $\{\sigma_i\}_{i=0}^d$ be from Lemmas 10.5 and 10.7. Then for $0 \leq i, j \leq d$ the scalar $\sigma_i - \sigma_j$ is given as follows:*

Case	$\sigma_i - \sigma_j$	
$\beta = 2$	$2(j - i)$	
$\beta = -2, d \text{ even}$	$j \text{ even}$	$j \text{ odd}$
	$i \text{ even}$	$2(j - i) \quad 2(d - i - j)$
	$i \text{ odd}$	$2(i + j - d) \quad 2(i - j)$
$\beta \neq \pm 2, d \text{ even}$	$\frac{(q^{2j} - q^{2i})(1 + q^{2d-2i-2j})}{q^d(q^2 - q^{-2})}$	
$\beta \neq \pm 2, d \text{ odd}$	$\frac{(q^{2j} - q^{2i})(1 + q^{2d-2i-2j})}{q^d(q - q^{-1})}$	

In the above table the scalar q is from Lemma 10.7.

Proof. Routine. \square

LEMMA 10.16. *For $d = 1$, $\mathcal{R}^{\text{asym}}$ is MutDist if and only if $\text{Char}(\mathbb{F}) \neq 2$. For $d \geq 2$, $\mathcal{R}^{\text{asym}}$ is MutDist if and only if the following conditions hold:*

Case	Conditions
$\beta = 2$	$\text{Char}(\mathbb{F})$ is 0 or greater than d
$\beta = -2$	d is even, $\text{Char}(\mathbb{F})$ is 0 or greater than d
$\beta \neq \pm 2$	$\text{Char}(\mathbb{F}) \neq 2, \quad q^{2i} \neq 1 \quad (1 \leq i \leq d), \quad q^{2i} \neq -1 \quad (1 \leq i \leq d - 1)$

In the above table the scalar q is from Lemma 10.7.

Proof. Use Lemmas 10.13–10.15. \square

We mention a lemma for later use.

LEMMA 10.17. *Let $\{\sigma_i\}_{i=0}^d$ denote a MutDist element in $\mathcal{R}^{\text{asym}}$. Then σ_0, σ_d are nonzero and*

$$\sigma_1\sigma_i \neq \sigma_0\sigma_{i-1}, \quad \sigma_1\sigma_i \neq \sigma_0\sigma_{i+1} \quad (1 \leq i \leq d - 1).$$

Proof. Without loss of generality, we may assume that $\{\sigma_i\}_{i=0}^d$ is the sequence from Lemmas 10.5, 10.7. One routinely verifies the results using Lemma 10.16. \square

Motivated by Lemma 4.16(iii) we consider the relation $\sigma_1\sigma_d = \sigma_0\sigma_{d-1}$. We will need the following result.

LEMMA 10.18. *For sequences $\{\sigma_i\}_{i=0}^d$ and $\{\tau_i\}_{i=0}^d$ in \mathcal{R} the following are equivalent:*

- (i) $\sigma_0\tau_1 \neq \sigma_1\tau_0$;
- (ii) $\{\sigma_i\}_{i=0}^d$ and $\{\tau_i\}_{i=0}^d$ are linearly independent.

Proof. (i) \Rightarrow (ii) Clear.

(ii) \Rightarrow (i) Consider the $d + 1$ by 2 matrix H with row i equal to σ_i, τ_i for $0 \leq i \leq d$. The rank of H is 2. By the β -recurrence, each row of H is a linear combination of rows 0 and 1. Therefore, rows 0 and 1 are linearly independent. Now take the determinant of the submatrix of H consisting of rows 0 and 1. \square

PROPOSITION 10.19. Assume that $\text{Char}(\mathbb{F}) \neq 2$. Then for a sequence $\{\sigma_i\}_{i=0}^d$ in \mathcal{R} the following are equivalent:

- (i) $\sigma_1\sigma_d = \sigma_0\sigma_{d-1}$;
- (ii) the sequence $\{\sigma_i\}_{i=0}^d$ is symmetric or antisymmetric.

Proof. By Lemma 10.3 there exist $\{\tau_i\}_{i=0}^d$ in \mathcal{R}^{sym} and $\{\mu_i\}_{i=0}^d$ in $\mathcal{R}^{\text{asym}}$ such that $\sigma_i = \tau_i + \mu_i$ for $0 \leq i \leq d$. We have

$$\begin{aligned} \sigma_0\sigma_{d-1} - \sigma_1\sigma_d &= (\tau_0 + \mu_0)(\tau_{d-1} + \mu_{d-1}) - (\tau_1 + \mu_1)(\tau_d + \mu_d) \\ &= (\tau_0 + \mu_0)(\tau_1 - \mu_1) - (\tau_1 + \mu_1)(\tau_0 - \mu_0) \\ &= 2(\tau_0\mu_1 - \tau_1\mu_0). \end{aligned}$$

By this and Lemma 10.18, $\sigma_0\sigma_{d-1} = \sigma_1\sigma_d$ if and only if $\{\tau_i\}_{i=0}^d$ and $\{\mu_i\}_{i=0}^d$ are linearly dependent. Since the sum $\mathcal{R}^{\text{sym}} + \mathcal{R}^{\text{asym}}$ is direct, the sequences $\{\tau_i\}_{i=0}^d, \{\mu_i\}_{i=0}^d$ are linearly dependent if and only if at least one of $\{\tau_i\}_{i=0}^d, \{\mu_i\}_{i=0}^d$ is zero. The result follows. \square

COROLLARY 10.20. Assume that $\text{Char}(\mathbb{F}) \neq 2$. Let $\{\sigma_i\}_{i=0}^d$ denote a MutDist element in \mathcal{R} . Then the following are equivalent:

- (i) $\{\sigma_i\}_{i=0}^d$ is antisymmetric;
- (ii) $\sigma_1\sigma_d = \sigma_0\sigma_{d-1}$.

Proof. By Proposition 10.19 and since no element of \mathcal{R}^{sym} is MutDist. \square

11. The classification of TB tridiagonal systems. In this section, we classify up to isomorphism the TB tridiagonal systems. Fix an integer $d \geq 1$ and consider a sequence of scalars taken from \mathbb{F} :

$$(11.68) \quad (\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d).$$

We now state the classification.

THEOREM 11.1. There exists a TB tridiagonal system Φ over \mathbb{F} that has eigenvalue array (11.68) if and only if the following (i)–(iii) hold:

- (i) $\theta_i \neq \theta_j, \theta_i^* \neq \theta_j^*$ if $i \neq j$ ($0 \leq i, j \leq d$);
- (ii) there exists $\beta \in \mathbb{F}$ such that both

$$(11.69) \quad \theta_{i-1} - \beta\theta_i + \theta_{i+1} = 0, \quad \theta_{i-1}^* - \beta\theta_i^* + \theta_{i+1}^* = 0 \quad (1 \leq i \leq d-1);$$

- (iii) $\theta_i + \theta_{d-i} = 0, \theta_i^* + \theta_{d-i}^* = 0$ ($0 \leq i \leq d$).

In this case Φ is unique up to isomorphism of TB tridiagonal systems.

NOTE 11.2. For $d \leq 2$, Theorem 11.1 remains valid if condition (ii) is deleted. Here is the reason. First assume that $d = 1$. Then (11.69) is vacuous and so holds for any $\beta \in \mathbb{F}$. Next assume that $d = 2$. Then condition (ii) follows from conditions (i), (iii). Indeed by condition (iii) we have $\theta_0 + \theta_2 = 0$ and $2\theta_1 = 0$. By this and condition (i) we have $\text{Char}(\mathbb{F}) \neq 2$ and $\theta_1 = 0$. Similarly $\theta_0^* + \theta_2^* = 0$ and $\theta_1^* = 0$. Therefore, (11.69) holds for any $\beta \in \mathbb{F}$.

The proof of Theorem 11.1 takes up most of this section.

LEMMA 11.3. *Assume that there exists a TB tridiagonal system Φ over \mathbb{F} that has eigenvalue array (11.68). Then this eigenvalue array satisfies conditions (i)–(iii) in Theorem 11.1. Moreover, Φ is unique up to isomorphism.*

Proof. Condition (i) holds by construction. Condition (ii) holds by Proposition 9.10. Condition (iii) holds by Lemmas 4.16(iii), 8.21(i), and Corollary 10.20. The uniqueness follows from Corollary 5.5. \square

For the rest of this section, we assume that the sequence (11.68) satisfies Conditions (i)–(iii) in Theorem 11.1. We will construct a TB tridiagonal system Φ over \mathbb{F} that has eigenvalue array (11.68). Fix $\beta \in \mathbb{F}$ that satisfies (11.69). We will refer to Section 10 with this β . Observe that each of the sequences $\{\theta_i\}_{i=0}^d$, $\{\theta_i^*\}_{i=0}^d$ is MutDist and contained in $\mathcal{R}^{\text{asym}}$. Note that $\text{Char}(\mathbb{F}) \neq 2$ by Lemma 10.13.

LEMMA 11.4. *There exists $0 \neq \zeta \in \mathbb{F}$ such that $\theta_i^* = \zeta\theta_i$ for $0 \leq i \leq d$.*

Proof. By Lemma 10.9 the subspace $\mathcal{R}^{\text{asym}}$ has dimension one. Each of $\{\theta_i\}_{i=0}^d$, $\{\theta_i^*\}_{i=0}^d$ is a nonzero element of $\mathcal{R}^{\text{asym}}$. The result follows. \square

LEMMA 11.5. *The following hold.*

- (i) *Each of θ_0 , θ_d , θ_0^* , θ_d^* is nonzero.*
- (ii) *$\theta_1/\theta_0 = \theta_{d-1}/\theta_d = \theta_1^*/\theta_0^* = \theta_{d-1}^*/\theta_d^*$.*

Proof. (i) By Lemmas 10.17 and 11.4.

(ii) We have $\theta_1\theta_d = \theta_0\theta_{d-1}$ by Corollary 10.20. The result follows from this and Lemma 11.4. \square

The following definition is motivated by Lemma 5.4.

DEFINITION 11.6. *Define scalars $\{c_i\}_{i=1}^d$, $\{b_i\}_{i=0}^{d-1}$ by*

$$(11.70) \quad c_i = \frac{\theta_1\theta_i^* - \theta_0\theta_{i+1}^*}{\theta_{i-1}^* - \theta_{i+1}^*} \quad (1 \leq i \leq d-1), \quad c_d = \theta_0,$$

$$(11.71) \quad b_i = \frac{\theta_1\theta_i^* - \theta_0\theta_{i-1}^*}{\theta_{i+1}^* - \theta_{i-1}^*} \quad (1 \leq i \leq d-1), \quad b_0 = \theta_0.$$

LEMMA 11.7. *We have $c_i = b_{d-i}$ for $1 \leq i \leq d$.*

Proof. Compare (11.70), (11.71) using the fact that the sequence $\{\theta_i^*\}_{i=0}^d$ is antisymmetric. \square

LEMMA 11.8. *The scalars $\{c_i\}_{i=1}^d$, $\{b_i\}_{i=0}^{d-1}$ are all nonzero.*

Proof. By Lemma 11.4, c_i is a nonzero scalar multiple of $\theta_1\theta_i - \theta_0\theta_{i+1}$ for $1 \leq i \leq d-1$ and $c_d = \theta_0$. By this and Lemma 10.17, $c_i \neq 0$ for $1 \leq i \leq d$. By this and Lemma 11.7, $b_i \neq 0$ for $0 \leq i \leq d-1$. \square

LEMMA 11.9. *The following hold:*

(i) $c_d = \theta_0$, $b_0 = \theta_0$ and

$$(11.72) \quad c_i + b_i = \theta_0 \quad (1 \leq i \leq d-1);$$

(ii) $c_d\theta_{d-1}^* = \theta_1\theta_d^*$, $b_0\theta_1^* = \theta_1\theta_0^*$ and

$$(11.73) \quad c_i\theta_{i-1}^* + b_i\theta_{i+1}^* = \theta_1\theta_i^* \quad (1 \leq i \leq d-1).$$

Proof. (i) Use Definition 11.6.

(ii) To get (11.73) use Definition 11.6. Concerning the first two equations, use $c_d = \theta_0$, $b_0 = \theta_0$, and Lemma 11.5. □

DEFINITION 11.10. *Define matrices A, A^* in $\text{Mat}_{d+1}(\mathbb{F})$ by*

$$A = \begin{pmatrix} 0 & b_0 & & & \mathbf{0} \\ c_1 & 0 & b_1 & & \\ & c_2 & \cdot & \cdot & \\ & & \cdot & \cdot & \\ \mathbf{0} & & & \cdot & b_{d-1} \\ & & & c_d & 0 \end{pmatrix}, \quad A^* = \text{diag}(\theta_0^*, \theta_1^*, \dots, \theta_d^*).$$

Recall the vector space $V = \mathbb{F}^{d+1}$. We are going to show that A, A^* is a TB tridiagonal pair on V and $\{\theta_i\}_{i=0}^d$ (resp. $\{\theta_i^*\}_{i=0}^d$) is a standard ordering of the eigenvalues of A (resp. A^*).

We first consider the eigenspaces of A^* , and the action of A on these eigenspaces. Define matrices $\{E_i^*\}_{i=0}^d$ in $\text{Mat}_{d+1}(\mathbb{F})$ by

$$E_i^* = \text{diag}(0, \dots, 0, \overset{i}{1}, 0, \dots, 0) \quad (0 \leq i \leq d).$$

For $0 \leq i \leq d$ define $V_i^* = E_i^*V$. Observe that V_i^* is a subspace of V with dimension 1.

LEMMA 11.11. *For $0 \leq i \leq d$, V_i^* is the eigenspace of A^* with eigenvalue θ_i^* . Moreover*

$$AV_i^* \subseteq V_{i-1}^* + V_{i+1}^* \quad (0 \leq i \leq d),$$

where $V_{-1}^* = 0$, $V_{d+1}^* = 0$.

Proof. By the form of the matrices A, A^* in Definition 11.10. □

Next we consider the eigenspaces of A , and the action of A^* on these eigenspaces. Each eigenspace of A has dimension 1, as we saw below (6.30). But conceivably A is not diagonalizable. For $0 \leq i \leq d$ define

$$V_i = \{v \in V \mid Av = \theta_i v\}.$$

The subspace V_i is nonzero if and only if θ_i is an eigenvalue of A , and in this case V_i has dimension 1. Our next goal is to show that each of $\{\theta_i\}_{i=0}^d$ is an eigenvalue of A , and $A^*V_i \subseteq V_{i-1} + V_{i+1}$ for $0 \leq i \leq d$, where $V_{-1} = 0$ and $V_{d+1} = 0$.

By Lemma 7.2 there exists $\varrho^* \in \mathbb{F}$ such that

$$(11.74) \quad \varrho^* = \theta_{i-1}^{*2} - \beta\theta_{i-1}^*\theta_i^* + \theta_i^{*2} \quad (1 \leq i \leq d).$$

LEMMA 11.12. *We have*

$$(11.75) \quad A^{*2}A - \beta A^*AA^* + AA^{*2} = \varrho^*A.$$

Proof. Let Z denote the left-hand side of (11.75) minus the right-hand side of (11.75). We show that $Z = 0$. To do this, we use Definition 11.10 and matrix multiplication to show that each entry of Z is zero. For $0 \leq i, j \leq d$ the (i, j) -entry of Z is

$$(11.76) \quad Z_{i,j} = (\theta_i^{*2} - \beta\theta_i^*\theta_j^* + \theta_j^{*2} - \varrho^*)A_{i,j}.$$

First assume that $|i - j| \neq 1$. Then $A_{i,j} = 0$ so $Z_{i,j} = 0$. Next assume that $|i - j| = 1$. Then in (11.76) the first factor on the right is zero so $Z_{i,j} = 0$. We have shown $Z = 0$, and the result follows. \square

LEMMA 11.13. *For $0 \leq i \leq d$ the scalar θ_i is an eigenvalue of A . Moreover,*

$$(11.77) \quad A^*V_i \subseteq V_{i-1} + V_{i+1} \quad (0 \leq i \leq d),$$

where $V_{-1} = 0$ and $V_{d+1} = 0$.

Proof. We first show that for $0 \leq i \leq d - 1$,

$$(11.78) \quad A^*V_i \subseteq V_0 + V_1 + \cdots + V_{i-1} + V_{i+1},$$

$$(11.79) \quad A^*V_i \not\subseteq V_0 + V_1 + \cdots + V_{i-1}.$$

We prove this using induction on i . Define vectors v_0, v_1 in V by

$$(11.80) \quad v_0 = (1, 1, \dots, 1)^t, \quad v_1 = (\theta_0^*, \theta_1^*, \dots, \theta_d^*)^t.$$

The vectors v_0, v_1 are nonzero. Using Lemma 11.9 one finds that $Av_0 = \theta_0v_0$ and $Av_1 = \theta_1v_1$. So $v_0 \in V_0$ and $v_1 \in V_1$. Using the form of A^* in Definition 11.10, we find $A^*v_0 = v_1$. So (11.78), (11.79) hold for $i = 0$. Assume that $1 \leq i \leq d - 1$. By induction,

$$(11.81) \quad A^*V_{i-1} \subseteq V_0 + V_1 + \cdots + V_{i-2} + V_i,$$

$$(11.82) \quad A^*V_{i-1} \not\subseteq V_0 + V_1 + \cdots + V_{i-2}.$$

By (11.82), $V_{i-1} \neq 0$. Pick $0 \neq v \in V_{i-1}$, and note that v is a basis for V_{i-1} . By (11.81) there exist $w \in V_0 + V_1 + \cdots + V_{i-2}$ and $v' \in V_i$ such that $A^*v = w + v'$. By (11.82) the vector v' is nonzero and hence a basis for V_i . We apply each side of (11.75) to v . Evaluate each term using $A^*v = w + v'$ and simplify the result using $\theta_{i-1} - \beta\theta_i = -\theta_{i+1}$ to get

$$(11.83) \quad (A - \theta_{i+1}I)A^*v' = -\theta_{i-1}A^*w + \beta A^*Aw - AA^*w + \varrho^*\theta_{i-1}v.$$

By construction $Aw \in V_0 + \cdots + V_{i-2}$. By induction $A^*(V_0 + \cdots + V_{i-2}) \subseteq V_0 + \cdots + V_{i-1}$. By these comments, the right-hand side of (11.83) is contained in $V_0 + \cdots + V_{i-1}$. So (11.83) yields

$$(A - \theta_{i+1}I)A^*v' \in V_0 + V_1 + \cdots + V_{i-1}.$$

Thus, there exist vectors $\{v_r\}_{r=0}^{i-1}$ in V such that $v_r \in V_r$ ($0 \leq r \leq i - 1$) and $(A - \theta_{i+1}I)A^*v' = \sum_{r=0}^{i-1} v_r$. Define $v'' \in V$ by

$$v'' = A^*v' + \sum_{r=0}^{i-1} \frac{v_r}{\theta_{i+1} - \theta_r}.$$

One routinely checks $Av'' = \theta_{i+1}v''$. So $v'' \in V_{i+1}$. Therefore, (11.78) holds. We show (11.79). Assume by way of contradiction that $A^*V_i \subseteq V_0 + V_1 + \cdots + V_{i-1}$. Define $W = V_0 + V_1 + \cdots + V_i$. Then $A^*W \subseteq W$. By construction $AW \subseteq W$. We have $W \neq 0$ since $i \geq 0$, and $W \neq V$ since $i \leq d-1$. These comments contradict Lemma 6.5(iii). Thus, (11.79) holds. We have now shown that (11.78) and (11.79) hold for $0 \leq i \leq d-1$.

Next we show that for $1 \leq i \leq d$,

$$(11.84) \quad A^*V_i \subseteq V_d + V_{d-1} + \cdots + V_{i+1} + V_{i-1},$$

$$(11.85) \quad A^*V_i \not\subseteq V_d + V_{d-1} + \cdots + V_{i+1}.$$

We prove this using induction on $i = d, d-1, \dots, 1$. Define vectors v_d, v_{d-1} in V by

$$(11.86) \quad v_d = (1, -1, 1, -1, \dots)^t, \quad v_{d-1} = (\theta_0^*, -\theta_1^*, \theta_2^*, -\theta_3^*, \dots)^t.$$

Each of v_d, v_{d-1} is nonzero. Using Lemma 11.9 and the fact that $\{\theta_i\}_{i=0}^d$ is antisymmetric, we obtain $Av_d = \theta_d v_d$ and $Av_{d-1} = \theta_{d-1} v_{d-1}$. So $v_d \in V_d$ and $v_{d-1} \in V_{d-1}$. Using the form of A^* in Definition 11.10 and (11.86), we obtain $A^*v_d = v_{d-1}$. So (11.84) and (11.85) hold for $i = d$. Assume that $1 \leq i \leq d-1$. Adjusting the proof of (11.78) and (11.79), we obtain (11.84) and (11.85). We have now shown that (11.84) and (11.85) hold for $1 \leq i \leq d$.

By (11.79) or (11.85) we see that V_i is nonzero for $0 \leq i \leq d$. Consequently, θ_i is an eigenvalue of A for $0 \leq i \leq d$. Comparing (11.78) and (11.84), we obtain (11.77). \square

LEMMA 11.14. *The pair A, A^* is a TB tridiagonal pair on V . Moreover, $\{\theta_i\}_{i=0}^d$ (resp. $\{\theta_i^*\}_{i=0}^d$) is a standard ordering of the eigenvalues of A (resp. A^*).*

Proof. We verify the conditions (i)–(iv) in Definition 3.1. By construction A^* is diagonalizable. By Lemma 11.13, A is diagonalizable. Thus, condition (i) holds. Conditions (ii) and (iii) hold by Lemmas 11.13 and 11.11, respectively. Condition (iv) holds by Lemma 6.5(iii). Thus, A, A^* is a TB tridiagonal pair on V . We have shown that $\{V_i\}_{i=0}^d$ (resp. $\{V_i^*\}_{i=0}^d$) is a standard ordering of the eigenspaces of A (resp. A^*). The result follows. \square

For $0 \leq i \leq d$ let $E_i \in \text{End}(V)$ denote the projection onto V_i . Define

$$\Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d).$$

LEMMA 11.15. *The above Φ is a TB tridiagonal system on V with eigenvalue array (11.68).*

Proof. By Lemma 11.14 the pair A, A^* is a TB tridiagonal pair on V . By Lemma 11.14 and the construction, $\{E_i\}_{i=0}^d$ (resp. $\{E_i^*\}_{i=0}^d$) is a standard ordering of the primitive idempotents of A (resp. A^*). The result follows. \square

By Lemmas 11.3, 11.15 we have now proved Theorem 11.1.

12. The eigenvalue array. In this section, we introduce the notion of an eigenvalue array over \mathbb{F} . Fix an integer $d \geq 1$ and consider a sequence of scalars taken from \mathbb{F} :

$$(12.87) \quad (\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d).$$

DEFINITION 12.1. *The sequence (12.87) is called an eigenvalue array over \mathbb{F} whenever it satisfies conditions (i)–(iii) in Theorem 11.1. We call d the diameter of the eigenvalue array.*

DEFINITION 12.2. Assume that (12.87) is an eigenvalue array over \mathbb{F} . By the corresponding TB tridiagonal system, we mean the one from Theorem 11.1.

DEFINITION 12.3. Assume that (12.87) is an eigenvalue array over \mathbb{F} . By an Askey–Wilson sequence for (12.87), we mean an Askey–Wilson sequence for the corresponding TB tridiagonal system.

DEFINITION 12.4. Assume that (12.87) is an eigenvalue array over \mathbb{F} . By a fundamental parameter for (12.87), we mean a fundamental parameter for the corresponding TB tridiagonal system.

We have some comments.

LEMMA 12.5. Assume that $\text{Char}(\mathbb{F}) = 2$. Then there does not exist an eigenvalue array over \mathbb{F} with diameter d .

Proof. Combine conditions (i), (iii) in Theorem 11.1. □

LEMMA 12.6. Assume that (12.87) is an eigenvalue array over \mathbb{F} .

- (i) Assume that $d \leq 2$. Then any scalar $\beta \in \mathbb{F}$ is a fundamental parameter for (12.87).
- (ii) Assume that $d \geq 3$. Then there exists a unique fundamental parameter β for (12.87).
- (iii) Assume that $d \geq 3$ and d is odd. Then $\beta \neq -2$.

Proof. (i) By Note 11.2.

(ii) By Definition 12.4 there exists at least one fundamental parameter β . This parameter satisfies $\theta_0 - \beta\theta_1 + \theta_2 = 0$ and $\theta_1 - \beta\theta_2 + \theta_3 = 0$. At least one of θ_1, θ_2 is nonzero since $\theta_1 \neq \theta_2$. By these comments, β is uniquely determined by $\theta_0, \theta_1, \theta_2, \theta_3$. The result follows.

(iii) By Lemma 10.14. □

LEMMA 12.7. Assume that (12.87) is an eigenvalue array over \mathbb{F} . Let β denote a fundamental parameter for (12.87). Then there exists a unique ordered pair ϱ, ϱ^* such that $\beta, \varrho, \varrho^*$ is an Askey–Wilson sequence for (12.87).

Proof. The scalars ϱ, ϱ^* exist by Proposition 8.23 and Definition 9.1. They are unique by (9.60), (9.61). □

LEMMA 12.8. Assume that (12.87) is an eigenvalue array over \mathbb{F} . Let $\beta, \varrho, \varrho^*$ denote an Askey–Wilson sequence for (12.87). Then the following hold.

- (i) Assume that d is even. Then

$$\varrho = \theta_r^2, \quad \varrho^* = \theta_r^{*2}, \quad \text{where } r = d/2 - 1.$$

- (ii) Assume that d is odd. Then

$$\varrho = (\beta + 2)\theta_r^2, \quad \varrho^* = (\beta + 2)\theta_r^{*2}, \quad \text{where } r = (d - 1)/2.$$

Proof. (i) By Theorem 11.1(iii), $\theta_{r+1} = 0$. By this and (9.60) at $i = r + 1$, we get $\varrho = \theta_r^2$. Similar for ϱ^* .

(ii) By Theorem 11.1(iii), $\theta_r + \theta_{r+1} = 0$. By this and (9.60) at $i = r + 1$, we get $\varrho = (\beta + 2)\theta_r^2$. Similar for ϱ^* . □

LEMMA 12.9. Assume that (12.87) is an eigenvalue array over \mathbb{F} . Then the following hold.

(i) Assume that d is even. Then

$$\theta_i = 0 \quad \text{if and only if} \quad i = d/2 \quad (0 \leq i \leq d).$$

(ii) Assume that d is odd. Then

$$\theta_i \neq 0 \quad (0 \leq i \leq d).$$

Proof. We have $\text{Char}(\mathbb{F}) \neq 2$ by Lemma 12.5. The result follows in view of Theorem 11.1(i), (iii). \square

LEMMA 12.10. Assume that (12.87) is an eigenvalue array over \mathbb{F} . Let $\beta, \varrho, \varrho^*$ denote an Askey–Wilson sequence for (12.87). Then the following are equivalent:

(i) $\varrho = 0$; (ii) $\varrho^* = 0$; (iii) $d = 1$ and $\beta = -2$.

Proof. First assume that d is even. Then each of ϱ, ϱ^* is nonzero by Lemmas 12.8(i) and 12.9(i). Next assume that d is odd. Then (i)–(iii) are equivalent by Lemmas 12.6(iii), 12.8(ii), 12.9(ii). \square

LEMMA 12.11. Assume that (12.87) is an eigenvalue array over \mathbb{F} . Then (12.87) has an Askey–Wilson sequence $\beta, \varrho, \varrho^*$ such that ϱ, ϱ^* are nonzero.

Proof. By Lemma 12.10 and since the choice of β is arbitrary for $d = 1$. \square

DEFINITION 12.12. Assume that (12.87) is an eigenvalue array over \mathbb{F} . By the corresponding intersection numbers (resp. dual intersection numbers), we mean the intersection numbers (resp. dual intersection numbers) of the corresponding TB tridiagonal system.

13. The TB tridiagonal systems in closed form. In this section, we give in closed form the eigenvalue array and the (dual) intersection numbers of a TB tridiagonal system. We also display the Askey–Wilson relations for the associated TB tridiagonal pair. Fix an integer $d \geq 1$ and consider a sequence of scalars taken from \mathbb{F} :

$$(13.88) \quad (\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d).$$

As a warmup, we will handle the cases $d = 1, d = 2$ separately.

LEMMA 13.1. For $d = 1$ the following (i), (ii) are equivalent:

(i) the sequence (13.88) is an eigenvalue array over \mathbb{F} ;
 (ii) $\text{Char}(\mathbb{F}) \neq 2$, the scalars θ_0, θ_0^* are nonzero, and $\theta_1 = -\theta_0, \theta_1^* = -\theta_0^*$.

Assume that (i), (ii) hold. Then the corresponding (dual) intersection numbers are

$$c_1 = \theta_0, \quad b_0 = \theta_0, \quad c_1^* = \theta_0^*, \quad b_0^* = \theta_0^*.$$

Moreover, the Askey–Wilson sequences for (13.88) are

$$\beta, \quad (\beta + 2)\theta_0^2, \quad (\beta + 2)\theta_0^{*2} \quad (\beta \in \mathbb{F}).$$

In addition,

$$AA^* = -A^*A, \quad A^2 = \theta_0^2 I, \quad A^{*2} = \theta_0^{*2} I.$$

Proof. (i) \Rightarrow (ii) By Theorem 11.1(iii), $\theta_1 = -\theta_0$ and $\theta_1^* = -\theta_0^*$. By this and Theorem 11.1(i), $\text{Char}(\mathbb{F}) \neq 2$ and $\theta_0 \neq 0, \theta_0^* \neq 0$.

(ii) \Rightarrow (i) One verifies the conditions (i)–(iii) in Theorem 11.1. Therefore, (13.88) is an eigenvalue array over \mathbb{F} .

Assume that (i), (ii) hold. Then the intersection numbers and dual intersection numbers are obtained using Lemma 5.4. The Askey–Wilson sequences are obtained by Lemmas 12.6(i) and 12.8(ii). By (ii) above, A has eigenvalues $\theta_0, -\theta_0$. So $A^2 = \theta_0^2 I$. Similarly, $A^{*2} = \theta_0^{*2} I$. By (3.7), A^* swaps the eigenspaces of A . By this we find that $AA^* = -A^*A$. \square

LEMMA 13.2. For $d = 2$ the following (i), (ii) are equivalent:

- (i) the sequence (13.88) is an eigenvalue array over \mathbb{F} ;
- (ii) $\text{Char}(\mathbb{F}) \neq 2$, the scalars θ_0, θ_0^* are nonzero, and

$$\theta_1 = 0, \quad \theta_2 = -\theta_0, \quad \theta_1^* = 0, \quad \theta_2^* = -\theta_0^*.$$

Assume that (i), (ii) hold. Then the corresponding (dual) intersection numbers are

$$\begin{array}{cccc} c_1 = \theta_0/2, & c_2 = \theta_0, & c_1^* = \theta_0^*/2, & c_2^* = \theta_0^*, \\ b_0 = \theta_0, & b_1 = \theta_0/2, & b_0^* = \theta_0^*, & b_1^* = \theta_0^*/2. \end{array}$$

Moreover, the Askey–Wilson sequences for (13.88) are

$$\beta, \quad \theta_0^2, \quad \theta_0^{*2} \quad (\beta \in \mathbb{F}).$$

In addition,

$$(13.89) \quad AA^*A = 0, \quad A^2A^* + A^*A^2 = \theta_0^2 A^*,$$

$$(13.90) \quad A^*AA^* = 0, \quad A^{*2}A + AA^{*2} = \theta_0^{*2} A.$$

Proof. For the assertions above (13.89), the proof is similar to Lemma 13.1. To obtain (13.89), set $\varrho = \theta_0^2$ in (8.42) and use the fact that β is arbitrary. Equation (13.90) is similarly obtained. \square

For the rest of this section, assume that $d \geq 3$. In Examples 13.3–13.6 below, we display all the parameter arrays over \mathbb{F} with diameter d . In each case, we display the (dual) intersection numbers from Definition 11.6, as well as the corresponding Askey–Wilson sequence $\beta, \varrho, \varrho^*$.

EXAMPLE 13.3. Assume that $\text{Char}(\mathbb{F})$ is 0 or greater than d . Assume that there exist nonzero $h, h^* \in \mathbb{F}$ such that

$$\theta_i = h(d - 2i), \quad \theta_i^* = h^*(d - 2i) \quad (0 \leq i \leq d).$$

Then (13.88) is an eigenvalue array over \mathbb{F} with fundamental parameter $\beta = 2$. The corresponding (dual) intersection numbers and the scalars ϱ, ϱ^* are

$$\begin{array}{ccc} c_i = hi, & c_i^* = h^*i & (1 \leq i \leq d), \\ b_i = h(d - i), & b_i^* = h^*(d - i) & (0 \leq i \leq d - 1), \\ \varrho = 4h^2, & \varrho^* = 4h^{*2}. & \end{array}$$

Proof. Define $\beta = 2$. One routinely verifies that each of $\{\theta_i\}_{i=0}^d, \{\theta_i^*\}_{i=0}^d$ is β -recurrent and antisymmetric, so contained in $\mathcal{R}^{\text{asym}}$. By Lemma 10.16, the subspace $\mathcal{R}^{\text{asym}}$ is MutDist, so each of $\{\theta_i\}_{i=0}^d, \{\theta_i^*\}_{i=0}^d$ is MutDist. By these comments (13.88) satisfies conditions (i)–(iii) in Theorem 11.1. Thus, (13.88) is an eigenvalue array over \mathbb{F} with fundamental parameter β . To get the (dual) intersection numbers and the scalars ρ, ρ^* , use Lemma 5.4 and (9.60), (9.61). \square

The following Examples can be verified in a similar way.

EXAMPLE 13.4. Assume that d is even. Assume that $\text{Char}(\mathbb{F})$ is 0 or greater than d . Assume that there exist nonzero $h, h^* \in \mathbb{F}$ such that

$$\theta_i = h(d - 2i)(-1)^i, \quad \theta_i^* = h^*(d - 2i)(-1)^i \quad (0 \leq i \leq d).$$

Then (13.88) is an eigenvalue array over \mathbb{F} with fundamental parameter $\beta = -2$. The corresponding (dual) intersection numbers and the scalars ρ, ρ^* are

$$\begin{aligned} c_i &= hi, & c_i^* &= h^*i & (1 \leq i \leq d), \\ b_i &= h(d - i), & b_i^* &= h^*(d - i) & (0 \leq i \leq d - 1), \\ \rho &= 4h^2, & \rho^* &= 4h^{*2}. \end{aligned}$$

EXAMPLE 13.5. Assume that d is even and $\text{Char}(\mathbb{F}) \neq 2$. Let q denote a nonzero scalar in $\overline{\mathbb{F}}$ such that

$$q^2 + q^{-2} \in \mathbb{F}, \quad q^{2i} \neq 1 \quad (1 \leq i \leq d), \quad q^{2i} \neq -1 \quad (1 \leq i \leq d - 1).$$

Assume that there exist nonzero $h, h^* \in \mathbb{F}$ such that

$$\theta_i = \frac{h(q^{d-2i} - q^{2i-d})}{q^2 - q^{-2}}, \quad \theta_i^* = \frac{h^*(q^{d-2i} - q^{2i-d})}{q^2 - q^{-2}} \quad (0 \leq i \leq d).$$

Then (13.88) is an eigenvalue array over \mathbb{F} with fundamental parameter $\beta = q^2 + q^{-2}$. Moreover, $\beta \neq \pm 2$. The corresponding (dual) intersection numbers and the scalars ρ, ρ^* are

$$\begin{aligned} c_i &= \frac{h(q^{2i} - q^{-2i})}{(q^2 - q^{-2})(q^{d-2i} + q^{2i-d})} & (1 \leq i \leq d - 1), & & c_d &= \frac{h(q^d - q^{-d})}{q^2 - q^{-2}}, \\ b_i &= \frac{h(q^{2d-2i} - q^{2i-2d})}{(q^2 - q^{-2})(q^{d-2i} + q^{2i-d})} & (1 \leq i \leq d - 1), & & b_0 &= \frac{h(q^d - q^{-d})}{q^2 - q^{-2}}, \\ \rho &= h^2. \end{aligned}$$

To get $\{c_i^*\}_{i=1}^d, \{b_i^*\}_{i=0}^{d-1}, \rho^*$, replace h with h^* in the above.

EXAMPLE 13.6. Assume that d is odd and $\text{Char}(\mathbb{F}) \neq 2$. Let q denote a nonzero scalar in $\overline{\mathbb{F}}$ such that

$$q^2 + q^{-2} \in \mathbb{F}, \quad q^{2i} \neq 1 \quad (1 \leq i \leq d), \quad q^{2i} \neq -1 \quad (1 \leq i \leq d-1).$$

Assume that there exist nonzero $h, h^* \in \mathbb{F}$ such that

$$\theta_i = \frac{h(q^{d-2i} - q^{2i-d})}{q - q^{-1}}, \quad \theta_i^* = \frac{h^*(q^{d-2i} - q^{2i-d})}{q - q^{-1}} \quad (0 \leq i \leq d).$$

Then (13.88) is an eigenvalue array over \mathbb{F} with fundamental parameter $\beta = q^2 + q^{-2}$. Moreover, $\beta \neq \pm 2$. The corresponding (dual) intersection numbers and the scalars ϱ, ϱ^* are

$$\begin{aligned} c_i &= \frac{h(q^{2i} - q^{-2i})}{(q - q^{-1})(q^{d-2i} + q^{2i-d})} & (1 \leq i \leq d-1), & & c_d &= \frac{h(q^d - q^{-d})}{q - q^{-1}}, \\ b_i &= \frac{h(q^{2d-2i} - q^{2i-2d})}{(q - q^{-1})(q^{d-2i} + q^{2i-d})} & (1 \leq i \leq d-1), & & b_0 &= \frac{h(q^d - q^{-d})}{q - q^{-1}}, \\ \varrho &= h^2(q + q^{-1})^2. \end{aligned}$$

To get $\{c_i^*\}_{i=1}^d, \{b_i^*\}_{i=0}^{d-1}, \varrho^*$, replace h with h^* in the above.

THEOREM 13.7. Every eigenvalue array over \mathbb{F} with diameter $d \geq 3$ is listed in exactly one of the Examples 13.3–13.6.

Proof. Assume that (13.88) is an eigenvalue array over \mathbb{F} , and let β denote its fundamental parameter. Observe that each of $\{\theta_i\}_{i=0}^d, \{\theta_i^*\}_{i=0}^d$ is contained in $\mathcal{R}^{\text{asym}}$. By this and Lemma 10.13, $\mathcal{R}^{\text{asym}}$ is MutDist. By this and Lemma 10.16 the conditions (10.67) hold. Let the sequence $\{\sigma_i\}_{i=0}^d$ be from Lemmas 10.5, 10.7. By Corollary 10.10, $\{\sigma_i\}_{i=0}^d$ is a basis for $\mathcal{R}^{\text{asym}}$. Therefore, there exist $h, h^* \in \mathbb{F}$ such that $\theta_i = h\sigma_i, \theta_i^* = h^*\sigma_i$ for $0 \leq i \leq d$. Now (13.88) is listed in the following Examples:

Case	Listed in
$\beta = 2$	Example 13.3
$\beta = -2$	Example 13.4
$\beta \neq \pm 2, d$ is even	Example 13.5
$\beta \neq \pm 2, d$ is odd	Example 13.6

The result follows. □

NOTE 13.8. For the eigenvalue array in Example 13.3 (resp. 13.4) (resp. 13.5, 13.6), the corresponding TB tridiagonal system is said to have Krawtchouk type (resp. Bannai/Ito type) (resp. q -Racah type).

We now display the Askey–Wilson relations. Let Φ denote a TB tridiagonal system over \mathbb{F} with eigenvalue array (13.88). Let A, A^* denote the TB tridiagonal pair associated with Φ .

LEMMA 13.9. Assume that $d \geq 3$. Then with the notation in Examples 13.3–13.6, the Askey–Wilson relations for A, A^* are given as follows:

Case	Askey–Wilson relations
$\beta = 2$	$A^2A^* - 2AA^*A + A^*A^2 = 4h^2A^*$ $A^{*2}A - 2A^*AA^* + AA^{*2} = 4h^{*2}A$
$\beta = -2$	$A^2A^* + 2AA^*A + A^*A^2 = 4h^2A^*$ $A^{*2}A + 2A^*AA^* + AA^{*2} = 4h^{*2}A$
$\beta \neq \pm 2, d$ even	$A^2A^* - (q^2 + q^{-2})AA^*A + A^*A^2 = h^2A^*$ $A^{*2}A - (q^2 + q^{-2})A^*AA^* + AA^{*2} = h^{*2}A$
$\beta \neq \pm 2, d$ odd	$A^2A^* - (q^2 + q^{-2})AA^*A + A^*A^2 = h^2(q + q^{-1})^2A^*$ $A^{*2}A - (q^2 + q^{-2})A^*AA^* + AA^{*2} = h^{*2}(q + q^{-1})^2A.$

Proof. Evaluate (8.42), (8.43) using the data in Examples 13.3–13.6. □

We will return to the Askey–Wilson relations in Section 15.

14. The relatives of a TB tridiagonal system. Let Φ denote a TB tridiagonal system over \mathbb{F} . Recall from Section 3 the relatives $\Phi^*, \Phi^\downarrow, \Phi^\updownarrow$. In this section, we discuss how these relatives are related to Φ at an algebraic level.

LEMMA 14.1. Consider a TB tridiagonal system over \mathbb{F} :

$$\Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d).$$

Then for $g \in \{\downarrow, \updownarrow, \updownarrow\}$ the TB tridiagonal system Φ^g is isomorphic to the TB tridiagonal system shown in the table below:

g	Φ^g is isomorphic to
\downarrow	$(A; \{E_i\}_{i=0}^d; -A^*; \{E_i^*\}_{i=0}^d)$
\updownarrow	$(-A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d)$
$\updownarrow\updownarrow$	$(-A; \{E_i\}_{i=0}^d; -A^*; \{E_i^*\}_{i=0}^d)$

Proof. Use Lemmas 3.14, 3.19 and Corollary 5.5, together with Theorem 11.1(iii). □

Referring to Lemma 14.1, the explicit isomorphisms will be displayed later in this section.

LEMMA 14.2. Let A, A^* denote a TB tridiagonal pair over \mathbb{F} . Then the following TB tridiagonal pairs are mutually isomorphic:

$$A, A^*, \quad A, -A^*, \quad -A, A^*, \quad -A, -A^*.$$

Proof. By Lemma 14.1. □

DEFINITION 14.3. Let A, A^* denote a TB tridiagonal pair over \mathbb{F} . Then A, A^* is said to be self-dual whenever A, A^* is isomorphic to A^*, A .

DEFINITION 14.4. Let Φ denote a TB tridiagonal system over \mathbb{F} . Then Φ is said to be self-dual whenever Φ is isomorphic to Φ^* .

LEMMA 14.5. Let Φ denote a TB tridiagonal system over \mathbb{F} with eigenvalue array $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d)$. Then the following are equivalent:

- (i) Φ is self-dual;
- (ii) $\theta_i = \theta_i^*$ for $0 \leq i \leq d$.

Proof. By Lemma 3.19, Φ^* has eigenvalue array $(\{\theta_i^*\}_{i=0}^d; \{\theta_i\}_{i=0}^d)$. By Corollary 5.5, Φ and Φ^* are isomorphic if and only if they have the same eigenvalue array. The result follows. \square

LEMMA 14.6. Assume that $d \geq 3$. Let Φ denote a TB tridiagonal system over \mathbb{F} with eigenvalue array $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d)$. Let the scalars h, h^* be from Examples 13.3–13.6. Then Φ is self-dual if and only if $h = h^*$.

Proof. By Lemma 14.5. \square

LEMMA 14.7. Consider a TB tridiagonal system over \mathbb{F} :

$$(14.91) \quad (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d).$$

Then there exists $0 \neq \zeta \in \mathbb{F}$ such that the TB tridiagonal system

$$(14.92) \quad (\zeta A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d)$$

is self-dual.

Proof. By Lemmas 11.4 and 14.5. \square

LEMMA 14.8. Let A, A^* denote a TB tridiagonal pair over \mathbb{F} . Then there exists $0 \neq \zeta \in \mathbb{F}$ such that the TB tridiagonal pair $\zeta A, A^*$ is self-dual.

Proof. By Lemma 14.7. \square

THEOREM 14.9. Let A, A^* denote a self-dual TB tridiagonal pair over \mathbb{F} . Then the following TB tridiagonal pairs are mutually isomorphic:

$$\begin{array}{cccc} A, A^*, & A, -A^*, & -A, A^*, & -A, -A^*, \\ A^*, A, & A^*, -A, & -A^*, A, & -A^*, -A. \end{array}$$

Proof. By Lemma 14.2 and Definition 14.3. \square

Our next goal is to display the isomorphisms in Lemma 14.1. For the rest of this section, fix a TB tridiagonal system over \mathbb{F} :

$$\Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d).$$

Let $(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d)$ denote the eigenvalue array of Φ .

DEFINITION 14.10. Define

$$(14.93) \quad S = \sum_{i=0}^d (-1)^i E_i, \quad S^* = \sum_{i=0}^d (-1)^i E_i^*.$$

LEMMA 14.11. We have $S^2 = I$ and $S^{*2} = I$. Moreover, S and S^* are invertible.

Proof. Concerning S , use $E_i E_j = \delta_{i,j} E_i$ ($0 \leq i, j \leq d$) and $I = \sum_{i=0}^d E_i$. The case of S^* is similar. \square

LEMMA 14.12. *The following (i)–(iv) hold:*

- (i) $SA = AS$;
- (ii) $SE_i = E_iS$ for $0 \leq i \leq d$;
- (iii) $SA^* = -A^*S$;
- (iv) $SE_i^* = E_{d-i}^*S$ for $0 \leq i \leq d$.

Proof. (i), (ii) By construction.

(iii) By Lemma 8.11 and Definition 14.10.

(iv) Using (2.6) we obtain

$$SE_i^*S^{-1} = \prod_{\substack{0 \leq j \leq d \\ j \neq i}} \frac{SA^*S^{-1} - \theta_j^*I}{\theta_i^* - \theta_j^*}.$$

Evaluate the above equation using $SA^*S^{-1} = -A^*$ and Theorem 11.1(iii) to get $SE_i^*S^{-1} = E_{d-i}^*$. The result follows. □

LEMMA 14.13. *The following (i)–(iv) hold:*

- (i) $S^*A^* = A^*S^*$;
- (ii) $S^*E_i^* = E_i^*S^*$ for $0 \leq i \leq d$;
- (iii) $S^*A = -AS^*$;
- (iv) $S^*E_i = E_{d-i}S^*$ for $0 \leq i \leq d$.

Proof. Apply Lemma 14.12 to Φ^* . □

LEMMA 14.14. *The following hold:*

- (i) S is an isomorphism of TB tridiagonal systems from Φ^\downarrow to

$$(A; \{E_i\}_{i=0}^d; -A^*; \{E_i^*\}_{i=0}^d).$$

- (ii) S is an isomorphism of TB tridiagonal pairs from A, A^* to $A, -A^*$.

Proof. Use Lemma 14.12. □

LEMMA 14.15. *The following hold:*

- (i) S^* is an isomorphism of TB tridiagonal systems from Φ^\downarrow to

$$(-A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d).$$

- (ii) S^* is an isomorphism of TB tridiagonal pairs from A, A^* to $-A, A^*$.

Proof. Use Lemma 14.13. □

LEMMA 14.16. *We have $SS^* = (-1)^d S^*S$.*

Proof. Using Definition 14.10 and Lemma 14.12(iv),

$$SS^* = \sum_{i=0}^d (-1)^i SE_i^* = \sum_{i=0}^d (-1)^i E_{d-i}^*S = \sum_{i=0}^d (-1)^{d-i} E_i^*S = (-1)^d S^*S. \quad \square$$

LEMMA 14.17. *The following hold:*

(i) SS^* is an isomorphism of TB tridiagonal systems from $\Phi^{\downarrow\downarrow}$ to

$$(-A; \{E_i\}_{i=0}^d; -A^*; \{E_i^*\}_{i=0}^d).$$

(ii) SS^* is an isomorphism of TB tridiagonal pairs from A, A^* to $-A, -A^*$.

Proof. Follows from Lemmas 14.14 and 14.15. □

For $0 \leq i \leq d$ we define some polynomials in $\mathbb{F}[x]$:

$$\begin{aligned} \tau_i(x) &= (x - \theta_0)(x - \theta_1) \cdots (x - \theta_{i-1}), \\ \eta_i(x) &= (x - \theta_d)(x - \theta_{d-1}) \cdots (x - \theta_{d-i+1}), \\ \tau_i^*(x) &= (x - \theta_0^*)(x - \theta_1^*) \cdots (x - \theta_{i-1}^*), \\ \eta_i^*(x) &= (x - \theta_d^*)(x - \theta_{d-1}^*) \cdots (x - \theta_{d-i+1}^*). \end{aligned}$$

THEOREM 14.18 (See [122]). *Assume that Φ is self-dual. Then the following four elements are equal, and this common element is an isomorphism of TB tridiagonal systems from Φ to Φ^* .*

$$\begin{aligned} \sum_{i=0}^d \eta_{d-i}(A) E_0^* E_d \tau_i^*(A^*), & \qquad \sum_{i=0}^d \eta_{d-i}^*(A^*) E_0 E_d^* \tau_i(A), \\ \sum_{i=0}^d \tau_i^*(A^*) E_d E_0^* \eta_{d-i}(A), & \qquad \sum_{i=0}^d \tau_i(A) E_d^* E_0 \eta_{d-i}^*(A^*). \end{aligned}$$

Moreover, the above common element is an isomorphism of TB tridiagonal pairs from A, A^* to A^*, A .

15. The \mathbb{Z}_3 -symmetric Askey–Wilson relations. For convenience, we adjust our notation as follows.

From now on we abbreviate $B = A^*$.

Let A, B denote a TB tridiagonal pair over \mathbb{F} . We saw in Section 8 that A, B satisfy the Askey–Wilson relations

$$(15.94) \qquad A^2 B - \beta A B A + B A^2 = \varrho B,$$

$$(15.95) \qquad B^2 A - \beta B A B + A B^2 = \varrho^* A.$$

A more detailed version of these relations is given in Lemma 13.9. In this section we put the Askey–Wilson relations in a form said to be \mathbb{Z}_3 -symmetric. This is done by introducing a third element C .

For the rest of this section, the following notation is in effect. Assume that \mathbb{F} is algebraically closed. Fix an integer $d \geq 1$, and let V denote a vector space over \mathbb{F} with dimension $d + 1$. Let A, B denote a TB tridiagonal pair on V . By Lemma 12.11, there exists an Askey–Wilson sequence $\beta, \varrho, \varrho^*$ for A, B such that ϱ, ϱ^* are nonzero.

DEFINITION 15.1. Let z, z', z'' denote scalars in \mathbb{F} that satisfy

Case	$z'z''$	$z''z$
$\beta = 2$	$-\varrho$	$-\varrho^*$
$\beta = -2$	ϱ	ϱ^*
$\beta \neq \pm 2$	$\varrho(4 - \beta^2)^{-1}$	$\varrho^*(4 - \beta^2)^{-1}$

Note that $zz'z'' \neq 0$.

PROPOSITION 15.2. Assume that $\beta = 2$. Then there exists $C \in \text{End}(V)$ such that

(15.97) $BC - CB = zA,$

(15.98) $CA - AC = z'B,$

(15.99) $AB - BA = z''C.$

Proof. Define C by (15.99). One verifies (15.97) and (15.98) using (15.94), (15.95), (15.96). □

PROPOSITION 15.3. Assume that $\beta = -2$. Then there exists $C \in \text{End}(V)$ such that

(15.100) $BC + CB = zA,$

(15.101) $CA + AC = z'B,$

(15.102) $AB + BA = z''C.$

Proof. Similar to the proof of Proposition 15.2. □

PROPOSITION 15.4. Assume that $\beta \neq \pm 2$. Let $0 \neq q \in \mathbb{F}$ be such that $\beta = q^2 + q^{-2}$. Then there exists $C \in \text{End}(V)$ such that

(15.103) $\frac{qBC - q^{-1}CB}{q^2 - q^{-2}} = zA,$

(15.104) $\frac{qCA - q^{-1}AC}{q^2 - q^{-2}} = z'B,$

(15.105) $\frac{qAB - q^{-1}BA}{q^2 - q^{-2}} = z''C.$

Proof. Similar to the proof of Proposition 15.2. □

LEMMA 15.5. Assume that A, B is self-dual. Then in Definition 15.1, the scalars z, z', z'' can be chosen such that $z = z' = z''$.

Proof. Use the fact that $\varrho = \varrho^*$ and \mathbb{F} is algebraically closed. □

The next three results follow from Propositions 15.2–15.4.

COROLLARY 15.6. Assume that $\beta = 2$. Then for

$$\rho = \rho^* = 4, \quad z = z' = z'' = 2\sqrt{-1},$$

the equations (15.97)–(15.99) become

(15.106) $BC - CB = 2\sqrt{-1}A,$

(15.107) $CA - AC = 2\sqrt{-1}B,$

(15.108) $AB - BA = 2\sqrt{-1}C.$

COROLLARY 15.7. Assume that $\beta = -2$. Then for

$$\rho = \rho^* = 4, \quad z = z' = z'' = 2,$$

the equations (15.100)–(15.102) become

$$(15.109) \quad BC + CB = 2A,$$

$$(15.110) \quad CA + AC = 2B,$$

$$(15.111) \quad AB + BA = 2C.$$

COROLLARY 15.8. Assume that $\beta \neq \pm 2$. Then for

$$\rho = \rho^* = 4 - \beta^2, \quad z = z' = z'' = 1,$$

the equations (15.103)–(15.105) become

$$(15.112) \quad \frac{qBC - q^{-1}CB}{q^2 - q^{-2}} = A,$$

$$(15.113) \quad \frac{qCA - q^{-1}AC}{q^2 - q^{-2}} = B,$$

$$(15.114) \quad \frac{qAB - q^{-1}BA}{q^2 - q^{-2}} = C.$$

NOTE 15.9. The equations (15.106)–(15.108) are the defining relations for \mathfrak{sl}_2 in the Pauli presentation. Here are some details. Assume that $\text{Char}(\mathbb{F}) \neq 2$. Let \mathfrak{sl}_2 denote the Lie algebra over \mathbb{F} consisting of the 2 by 2 matrices that have entries in \mathbb{F} and trace 0. The Lie bracket is $[r, s] = rs - sr$. Consider the Pauli matrices (see [124]):

$$S_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad S_2 = \begin{pmatrix} 0 & -\sqrt{-1} \\ \sqrt{-1} & 0 \end{pmatrix}, \quad S_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

These matrices form a basis for \mathfrak{sl}_2 , and satisfy the relations

$$[S_1, S_2] = 2\sqrt{-1} S_3, \quad [S_2, S_3] = 2\sqrt{-1} S_1, \quad [S_3, S_1] = 2\sqrt{-1} S_2.$$

NOTE 15.10. The equations (15.109)–(15.111) are essentially the defining relations for the anticommutator spin algebra (see [12]). Here are some details. In [12] Arik and Kayserilioglu introduced an \mathbb{F} -algebra by generators J_1, J_2, J_3 and relations

$$J_2J_3 + J_3J_2 = J_1, \quad J_3J_1 + J_1J_3 = J_2, \quad J_1J_2 + J_2J_1 = J_3.$$

This algebra is called the *anticommutator spin algebra*. Observe that the above relations coincide with (15.109)–(15.111) by setting $J_1 = A/2, J_2 = B/2, J_3 = C/2$.

NOTE 15.11. The equations (15.112)–(15.114) are essentially the defining relations for the quantum algebra $U_q(\mathfrak{so}_3)$. Here are some details. The algebra $U_q(\mathfrak{so}_3)$ has a presentation by generators I_1, I_2, I_3 and relations

$$(15.115) \quad q^{1/2}I_2I_3 - q^{-1/2}I_3I_2 = I_1,$$

$$(15.116) \quad q^{1/2}I_3I_1 - q^{-1/2}I_1I_3 = I_2,$$

$$(15.117) \quad q^{1/2}I_1I_2 - q^{-1/2}I_2I_1 = I_3.$$

As far as we know, the equations (15.115)–(15.117) were first considered by Santilli [128]. Later in [40], Fairlie discovered that (15.115)–(15.117) show up in $U_q(\mathfrak{so}_3)$. The fact that (15.115)–(15.117) give a presentation for $U_q(\mathfrak{so}_3)$ was proved by Odesski in [123]. See [58] for a more precise history. The relations (15.115)–(15.117) become (15.112)–(15.114) by first setting $I_1 = A/(q - q^{-1})$, $I_2 = B/(q - q^{-1})$, $I_3 = C/(q - q^{-1})$, and then replacing q with q^2 .

16. The elements W , W' , W'' and the automorphism ρ . Throughout this section, the following notation is in effect. Assume that \mathbb{F} is algebraically closed. Fix an integer $d \geq 1$, and let V denote a vector space over \mathbb{F} with dimension $d + 1$. Let A, B denote a TB tridiagonal pair on V . In view of Lemma 14.8, assume that A, B is self-dual. Let $\beta, \varrho, \varrho^*$ denote the Askey–Wilson sequence for A, B from above Definition 15.1 and note that $\varrho = \varrho^*$. For the case $\beta \neq \pm 2$, fix $0 \neq q \in \mathbb{F}$ as in Proposition 15.4.

LEMMA 16.1. *Let $\{\theta_i\}_{i=0}^d$ denote an eigenvalue sequence of A, B . Then there exists $0 \neq h \in \mathbb{F}$ such that the following holds:*

Case	θ_i
$\beta = 2$	$h(d - 2i)$
$\beta = -2$	$h(d - 2i)(-1)^i$
$\beta \neq \pm 2$	$h(q^{d-2i} - q^{2i-d})$

(16.118)

Proof. By Lemmas 13.1, 13.2 and Examples 13.3–13.6, together with the assumption that \mathbb{F} is algebraically closed. □

LEMMA 16.2. *Referring to Lemma 16.1,*

Case	ϱ
$\beta = 2$	$4h^2$
$\beta = -2$	$4h^2$
$\beta \neq \pm 2$	$h^2(q^2 - q^{-2})^2$

(16.119)

Proof. Use Proposition 9.4 and Lemma 16.1. □

In view of Lemma 15.5, we choose the scalars z, z', z'' in Definition 15.1 such that $z = z' = z''$. Let $C \in \text{End}(V)$ be from Propositions 15.2–15.4. Next, we define some invertible elements W, W' in $\text{End}(V)$, and use them to construct an automorphism ρ of $\text{End}(V)$ that sends

$$A \mapsto B, \quad B \mapsto C, \quad C \mapsto A.$$

For $0 \leq i \leq d$ let E_i (resp. E'_i) denote the primitive idempotent of A (resp. B) for the eigenvalue θ_i , where θ_i is from Lemma 16.1.

DEFINITION 16.3. *Define $W, W' \in \text{End}(V)$ by*

$$(16.120) \quad W = \sum_{i=0}^d t_i E_i, \quad W' = \sum_{i=0}^d t_i E'_i,$$

where

Case	t_i
$\beta = 2$	$2^i h^i z^{-i}$
$\beta = -2$	$(-1)^{\lfloor i/2 \rfloor} 2^i h^i z^{-i}$
$\beta \neq \pm 2$	$h^i z^{-i} q^{i(d-i)}$

(16.121)

LEMMA 16.4. *The elements W, W' are invertible, with inverse*

$$W^{-1} = \sum_{i=0}^d t_i^{-1} E_i, \quad (W')^{-1} = \sum_{i=0}^d t_i^{-1} E'_i.$$

Proof. By construction. □

Recall the antiautomorphism \dagger of $\text{End}(V)$, from Lemma 8.5.

LEMMA 16.5. *The antiautomorphism \dagger fixes each of W, W' .*

Proof. By (16.120) and since \dagger fixes E_i and E'_i for $0 \leq i \leq d$. □

LEMMA 16.6. *We have*

$$(16.122) \quad AW = WA, \quad BW' = W'B.$$

Proof. The element A commutes with W by Definition 16.3 and since A commutes with E_i for $0 \leq i \leq d$. Similarly, B commutes with W' . □

LEMMA 16.7. *We have*

$$(16.123) \quad BW = WC, \quad CW' = W'A.$$

Proof. We first obtain $BW = WC$. By (15.99), (15.102), (15.105), we have

$$C = eAB + e'BA,$$

where

Case	e	e'
$\beta = 2$	z^{-1}	$-z^{-1}$
$\beta = -2$	z^{-1}	z^{-1}
$\beta \neq \pm 2$	$\frac{qz^{-1}}{q^2 - q^{-2}}$	$-\frac{q^{-1}z^{-1}}{q^2 - q^{-2}}$

It suffices to show that

$$(16.125) \quad eAB + e'BA - W^{-1}BW = 0.$$

To obtain (16.125), we show that

$$(16.126) \quad E_i(eAB + e'BA - W^{-1}BW)E_j = 0,$$

for $0 \leq i, j \leq d$. Let i, j be given. Using $E_iA = \theta_i E_i$, $AE_j = \theta_j E_j$, $E_iW^{-1} = t_i^{-1} E_i$, $WE_j = t_j E_j$, we find that the left-hand side of (16.126) is equal to

$$(16.127) \quad (e\theta_i + e'\theta_j - t_i^{-1}t_j)E_iBE_j.$$

First, assume that $|i - j| \neq 1$. Then $E_iBE_j = 0$ by Lemma 3.21, so (16.127) is zero. Next assume that $|i - j| = 1$. Using (15.96), (16.118), (16.119), (16.121), (16.124), one routinely finds that in (16.127) the coefficient of E_iBE_j is zero. Therefore (16.127) is zero as desired. We have obtained $BW = WC$. The equation $CW' = W'A$ is similarly obtained. □

DEFINITION 16.8. Define $P = W'W$. Note that $P^\dagger = WW'$.

LEMMA 16.9. We have

$$(16.128) \quad AP = PB, \quad BP = PC, \quad CP = PA.$$

Proof. In the equation on the left in (16.123), multiply each side on the left by W' and use (16.122) to get $BP = PC$. In the equation on the right in (16.123), multiply each side on the right by W and use (16.122) to get $CP = PA$. Combining the equations in (16.123) we obtain $BWW' = WW'A$. By these comments and Definition 16.8, we obtain $BP^\dagger = P^\dagger A$. Applying \dagger we get $AP = PB$. \square

DEFINITION 16.10. Define the map $\rho : \text{End}(V) \rightarrow \text{End}(V)$, $X \mapsto P^{-1}XP$. Note that ρ is an automorphism of $\text{End}(V)$ that fixes P .

COROLLARY 16.11. The automorphism ρ sends

$$A \mapsto B, \quad B \mapsto C, \quad C \mapsto A.$$

Proof. By Lemma 16.9. \square

COROLLARY 16.12. We have $\rho^3 = 1$. Moreover, there exists $0 \neq \kappa \in \mathbb{F}$ such that $P^3 = \kappa I$.

Proof. By Corollary 16.11, the element P^3 commutes with both A, B . By this and Lemma 8.24, there exists $\kappa \in \mathbb{F}$ such that $P^3 = \kappa I$. We have $\kappa \neq 0$ since P is invertible. Now $\rho^3 = 1$ follows by Definition 16.10. \square

NOTE 16.13 (See [150, Corollary 14.11]). Referring to Corollary 16.12, the scalar κ is given as follows:

Case	$\beta = 2$	$\beta = -2$	$\beta \neq \pm 2$
κ	$(-1)^d 2^{-d} h^{-d} z^d$	1	$(-1)^d h^{-d} z^d q^{d(d-1)}$

By construction and Corollary 16.11, the element C is diagonalizable with eigenvalues $\{\theta_i\}_{i=0}^d$. For $0 \leq i \leq d$ let E''_i denote the primitive idempotent of C for the eigenvalue θ_i .

LEMMA 16.14. For $0 \leq i \leq d$ the automorphism ρ sends

$$E_i \mapsto E'_i, \quad E'_i \mapsto E''_i, \quad E''_i \mapsto E_i.$$

Proof. We first show that ρ sends $E_i \mapsto E'_i$. In (2.6) the element E_i is expressed as a polynomial in A ; let $f(x)$ denote the polynomial. Using the equation on the left in (16.128),

$$P^{-1}E_iP = P^{-1}f(A)P = f(P^{-1}AP) = f(B) = E'_i.$$

Thus ρ sends $E_i \mapsto E'_i$. Similarly, ρ sends $E'_i \mapsto E''_i$ and $E''_i \mapsto E_i$. \square

DEFINITION 16.15. Define $W'' \in \text{End}(V)$ by

$$W'' = \sum_{i=0}^d t_i E''_i,$$

where $\{t_i\}_{i=0}^d$ are from (16.121).

LEMMA 16.16. *The automorphism ρ sends*

$$W \mapsto W', \quad W' \mapsto W'', \quad W'' \mapsto W.$$

In other words

$$(16.129) \quad WP = PW', \quad W'P = PW'', \quad W''P = PW.$$

Proof. The first assertion is by Definitions 16.3, 16.15 and Lemma 16.14. The second assertion follows in view of Definition 16.10. \square

LEMMA 16.17. *The element P is equal to each of*

$$W'W, \quad W''W', \quad WW''.$$

Proof. By Definition 16.8, $P = W'W$. Apply ρ to this and use Lemma 16.16 to get $P = W''W'$. Similarly $P = WW''$. \square

DEFINITION 16.18. *Elements X, Y in $\text{End}(V)$ are said to satisfy the braid relation whenever $XYX = YXY$.*

LEMMA 16.19. *Any two of W, W', W'' satisfy the braid relation.*

Proof. By (16.129) and Lemma 16.17. \square

Corollary 16.11 and Lemma 16.14 show that A, B, C is a Leonard triple in the sense of Curtin [35]. We will say more about Leonard triples in Section 19.

17. Some antiautomorphisms associated with a TB tridiagonal pair. Throughout this section, the following notation is in effect. Assume that \mathbb{F} is algebraically closed. Fix an integer $d \geq 1$, and let V denote a vector space over \mathbb{F} with dimension $d + 1$. Let A, B denote a self-dual TB tridiagonal pair on V . Let $\beta, \varrho, \varrho^*$ denote the Askey–Wilson sequence for A, B from above Definition 15.1. Choose the scalars z, z', z'' in Definition 15.1 such that $z = z' = z''$ and let $C \in \text{End}(V)$ be from Propositions 15.2–15.4. In this section, we obtain some antiautomorphisms of $\text{End}(V)$ that act on A, B, C in an attractive manner. Recall the antiautomorphism \dagger of $\text{End}(V)$, from Lemma 8.5. By construction \dagger fixes A, B .

LEMMA 17.1. *For a map $\xi : \text{End}(V) \rightarrow \text{End}(V)$ the following are equivalent:*

- (i) ξ is an antiautomorphism of $\text{End}(V)$;
- (ii) there exists an invertible $T \in \text{End}(V)$ such that $X^\xi = T^{-1}X^\dagger T$ for all $X \in \text{End}(V)$.

Proof. (i) \Rightarrow (ii) Consider the composition

$$\omega : \text{End}(V) \xrightarrow{\dagger} \text{End}(V) \xrightarrow{\xi} \text{End}(V).$$

The map ω is an automorphism of $\text{End}(V)$. By the Skolem–Noether theorem, there exists an invertible $T \in \text{End}(V)$ such that $X^\omega = T^{-1}XT$ for all $X \in \text{End}(V)$. Thus, $X^\xi = (X^\dagger)^\omega = T^{-1}X^\dagger T$ for all $X \in \text{End}(V)$.

(ii) \Rightarrow (i) Clear. \square

Let $\{\theta_i\}_{i=0}^d$ denote the eigenvalue sequence of A, B from Lemma 16.1, and let E_i, E'_i be from above Definition 16.3. Let W, W' be from Definition 16.3, and let P be from Definition 16.8.

DEFINITION 17.2. Define the maps $\dagger', \dagger'' : \text{End}(V) \rightarrow \text{End}(V)$, $X \mapsto T^{-1}X\dagger T$, where T is from the table below.

$$\begin{array}{c|cc} & \dagger' & \dagger'' \\ \hline T & P\dagger P & (PP\dagger)^{-1} \end{array}.$$

Note that \dagger', \dagger'' are antiautomorphisms of $\text{End}(V)$.

Recall the automorphism ρ of $\text{End}(V)$, from Definition 16.10.

LEMMA 17.3. The map \dagger' is equal to the composition

$$\text{End}(V) \xrightarrow{\rho^{-1}} \text{End}(V) \xrightarrow{\dagger} \text{End}(V) \xrightarrow{\rho} \text{End}(V).$$

The map \dagger'' is equal to the composition

$$\text{End}(V) \xrightarrow{\rho} \text{End}(V) \xrightarrow{\dagger} \text{End}(V) \xrightarrow{\rho^{-1}} \text{End}(V).$$

Proof. Use Definitions 16.10 and 17.2. □

Our next goal is to describe how $\dagger, \dagger', \dagger''$ act on A, B, C . We will treat separately the cases $\beta = 2$, $\beta = -2$, $\beta \neq \pm 2$.

PROPOSITION 17.4. Assume that $\beta = 2$. Then the antiautomorphisms $\dagger, \dagger', \dagger''$ act on A, B, C as follows:

- (i) \dagger fixes A, B and sends $C \mapsto -C$;
- (ii) \dagger' fixes B, C and sends $A \mapsto -A$;
- (iii) \dagger'' fixes C, A and sends $B \mapsto -B$.

Proof. (i) By construction $A\dagger = A$ and $B\dagger = B$. Applying \dagger to each side of (15.99), we obtain $C\dagger = -C$.

(ii), (iii) Use Corollary 16.11, Lemma 17.3, and (i) above. □

PROPOSITION 17.5. Assume that $\beta = -2$. Then the maps $\dagger, \dagger', \dagger''$ coincide, and this map fixes each of A, B, C .

Proof. By Definitions 15.1, 16.3 and Lemma 16.2. we obtain $t_i^2 = 1$ for $0 \leq i \leq d$. So $W^2 = I$ and $(W')^2 = I$. By this and Definition 16.8, $P\dagger P = I$ and $PP\dagger = I$. By this and Definition 17.2, we obtain $\dagger' = \dagger$ and $\dagger'' = \dagger$. By construction \dagger fixes A, B . By this and (15.102), we obtain $C\dagger = C$. The result follows. □

PROPOSITION 17.6. Assume that $\beta \neq \pm 2$. Then the antiautomorphisms $\dagger, \dagger', \dagger''$ act on A, B, C as follows:

- (i) \dagger fixes A, B and sends $C \mapsto C - \frac{AB-BA}{z(q-q^{-1})}$;
- (ii) \dagger' fixes B, C and sends $A \mapsto A - \frac{BC-CB}{z(q-q^{-1})}$;
- (iii) \dagger'' fixes C, A and sends $B \mapsto B - \frac{CA-AC}{z(q-q^{-1})}$.

Proof. Similar to the proof of Proposition 17.4, using (15.105) instead of (15.99). □

DEFINITION 17.7. Define maps $\ddagger, \ddagger', \ddagger'' : \text{End}(V) \rightarrow \text{End}(V)$, $X \mapsto T^{-1}X\dagger T$, where T is from the table below.

$$\begin{array}{c|ccc} & \ddagger & \ddagger' & \ddagger'' \\ \hline T & W & (W')^{-1} & WW'W \end{array}$$

Note that $\ddagger, \ddagger', \ddagger''$ are antiautomorphisms of $\text{End}(V)$.

LEMMA 17.8. *The map \ddagger' is equal to the composition*

$$(17.130) \quad \text{End}(V) \xrightarrow{\rho^{-1}} \text{End}(V) \xrightarrow{\ddagger} \text{End}(V) \xrightarrow{\rho} \text{End}(V).$$

The map \ddagger'' is equal to the composition

$$(17.131) \quad \text{End}(V) \xrightarrow{\rho} \text{End}(V) \xrightarrow{\ddagger} \text{End}(V) \xrightarrow{\rho^{-1}} \text{End}(V).$$

Proof. Note by Lemma 16.5 that \ddagger fixes each of W, W' . We first show that \ddagger' is equal to the composition (17.130). Pick any $X \in \text{End}(V)$. By Definition 17.7, \ddagger' sends $X \mapsto W'X^\dagger(W')^{-1}$. By Definitions 16.8, 16.10, 17.7, the composition (17.130) sends $X \mapsto H^{-1}X^\dagger H$, where $H = WW'WW'W$. By Definition 16.8 and Corollary 16.12, we obtain $H = \kappa(W')^{-1}$. By these comments \ddagger' is equal to the composition (17.130). One similarly shows that \ddagger'' is equal to the composition (17.131). \square

PROPOSITION 17.9. *The antiautomorphisms $\ddagger, \ddagger', \ddagger''$ act on A, B, C as follows:*

- (i) \ddagger fixes A and swaps B, C ;
- (ii) \ddagger' fixes B and swaps C, A ;
- (iii) \ddagger'' fixes C and swaps A, B .

Proof. Note by Lemma 16.5 that \ddagger fixes each of W, W' .

(i) By Lemma 16.6 and Definition 17.7, $A^\ddagger = A$. By Lemma 16.7, $B^\ddagger = C$ and $C^\ddagger = W^{-1}C^\dagger W = (WCW^{-1})^\dagger = B$.

(ii), (iii) Use Corollary 16.11, Lemma 17.8, and (i) above. \square

The existence of the antiautomorphisms $\ddagger, \ddagger', \ddagger''$ shows that the Leonard triple A, B, C is modular in the sense of Curtin [35].

LEMMA 17.10. *We have $\ddagger^2 = 1, (\ddagger')^2 = 1, (\ddagger'')^2 = 1$.*

Proof. Each of the given squares is an automorphism of $\text{End}(V)$ that fixes each of A, B, C . The result follows in view of Lemma 8.2(i). \square

Next, we explain how $\ddagger, \ddagger', \ddagger''$ are related to the automorphism ρ from Definition 16.10.

PROPOSITION 17.11. *The automorphism ρ is equal to each of the following compositions:*

$$(17.132) \quad \text{End}(V) \xrightarrow{\ddagger'} \text{End}(V) \xrightarrow{\ddagger} \text{End}(V),$$

$$(17.133) \quad \text{End}(V) \xrightarrow{\ddagger''} \text{End}(V) \xrightarrow{\ddagger'} \text{End}(V),$$

$$(17.134) \quad \text{End}(V) \xrightarrow{\ddagger} \text{End}(V) \xrightarrow{\ddagger''} \text{End}(V).$$

Proof. We first show that ρ is equal to the composition (17.132). Pick any $X \in \text{End}(V)$. By Definition 16.10, ρ sends $X \mapsto P^{-1}XP$. By Definition 17.7, the composition (17.132) sends $X \mapsto W^{-1}(W')^{-1}XW'W$. By Definition 16.8, $P = W'W$. By these comments, ρ is equal to the composition (17.132). By this and Lemma 17.8, we find that ρ is equal to each of (17.133), (17.134). \square

18. An action of $\text{PSL}_2(\mathbb{Z})$ associated with a TB tridiagonal pair. Throughout this section, the following notation is in effect. Assume that \mathbb{F} is algebraically closed. Fix an integer $d \geq 1$, and let V denote a vector space over \mathbb{F} with dimension $d + 1$. Let A, B denote a self-dual TB tridiagonal pair on V . Let $\beta, \varrho, \varrho^*$ denote the Askey–Wilson sequence for A, B from above Definition 15.1. Choose the scalars z, z', z'' in Definition 15.1 such that $z = z' = z''$ and let $C \in \text{End}(V)$ be from Propositions 15.2–15.4.

In this section, we display an action of $\text{PSL}_2(\mathbb{Z})$ on $\text{End}(V)$ as a group of automorphisms that act on A, B, C in an attractive manner. Recall from [10] that $\text{PSL}_2(\mathbb{Z})$ has a presentation by generators r, s and relations $r^3 = 1, s^2 = 1$. To get the action of $\text{PSL}_2(\mathbb{Z})$, we need an automorphism of $\text{End}(V)$ that has order 3 and one that has order 2. In Corollary 16.12, we obtained an automorphism ρ of $\text{End}(V)$ that has order 3. Next we obtain an automorphism of $\text{End}(V)$ that has order 2. Recall the elements W, W' from Definition 16.3.

DEFINITION 18.1. Define the map $\sigma : \text{End}(V) \rightarrow \text{End}(V), X \mapsto TXT^{-1}$, where $T = WW'W$. Note that σ is an automorphism of $\text{End}(V)$.

Recall the antiautomorphism \dagger of $\text{End}(V)$ from Lemma 8.5, and the antiautomorphism \ddagger'' of $\text{End}(V)$ from Definition 17.7.

PROPOSITION 18.2. The automorphism σ is equal to the composition

$$\text{End}(V) \xrightarrow{\ddagger''} \text{End}(V) \xrightarrow{\dagger} \text{End}(V).$$

Proof. Routine verification using Definitions 17.7 and 18.1. □

PROPOSITION 18.3. The automorphism σ swaps A, B and sends $C \mapsto C^\dagger$.

Proof. By Propositions 17.9(iii) and 18.2. □

COROLLARY 18.4. We have $\sigma^2 = 1$.

Proof. By Proposition 18.3 the automorphism σ^2 fixes both A, B . The result follows by Lemma 8.2(i). □

COROLLARY 18.5. The group $\text{PSL}_2(\mathbb{Z})$ acts on $\text{End}(V)$ as a group of automorphisms, such that r acts as ρ and s acts as σ .

Proof. By Corollaries 16.12 and 18.4. □

19. Concluding remarks. In earlier sections, we discussed TB tridiagonal pairs. In this section, we summarize what is known about general tridiagonal pairs, using the TB case as a guide. The definition of a tridiagonal pair is given in the Introduction section. The definition of a tridiagonal system is analogous to Definition 3.7. For the rest of this section, let $\Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d)$ denote a tridiagonal system on V . It is known that $d = \delta$ (see [71, Lemma 4.5]). By [71, Theorem 10.1], there exists a sequence of scalars $\beta, \gamma, \gamma^*, \varrho, \varrho^*$ in \mathbb{F} such that both

$$(19.135) \quad 0 = [A, A^2A^* - \beta AA^*A + A^*A^2 - \gamma(AA^* + A^*A) - \varrho A^*],$$

$$(19.136) \quad 0 = [A^*, A^{*2}A - \beta A^*AA^* + AA^{*2} - \gamma^*(A^*A + AA^*) - \varrho^* A],$$

where $[r, s]$ means $rs - sr$. The above equations are called the *tridiagonal relations*. We now describe the eigenvalues. For $0 \leq i \leq d$ let θ_i (resp. θ_i^*) denote the eigenvalue of A (resp. A^*) associated with E_i (resp. E_i^*). By [71, Theorem 11.1], the expressions

$$\frac{\theta_{i-2} - \theta_{i+1}}{\theta_{i-1} - \theta_i}, \quad \frac{\theta_{i-2}^* - \theta_{i+1}^*}{\theta_{i-1}^* - \theta_i^*},$$

are equal and independent of i for $2 \leq i \leq d-1$. Moreover,

$$\begin{aligned} \gamma &= \theta_{i-1} - \beta\theta_i + \theta_{i+1} & (1 \leq i \leq d-1), \\ \gamma^* &= \theta_{i-1}^* - \beta\theta_i^* + \theta_{i+1}^* & (1 \leq i \leq d-1), \\ \varrho &= \theta_{i-1}^2 - \beta\theta_{i-1}\theta_i + \theta_i^2 - \gamma(\theta_{i-1} + \theta_i) & (1 \leq i \leq d), \\ \varrho^* &= \theta_{i-1}^{*2} - \beta\theta_{i-1}^*\theta_i^* + \theta_i^{*2} - \gamma^*(\theta_{i-1}^* + \theta_i^*) & (1 \leq i \leq d). \end{aligned}$$

Note that γ, γ^* are zero if Φ is TB. One topic that we did not discuss in earlier sections is the split decomposition. This is defined as follows. For $0 \leq i \leq d$ define

$$U_i = (E_0^*V + \cdots + E_i^*V) \cap (E_iV + \cdots + E_dV).$$

By [71, Theorem 4.6], we have $V = \sum_{i=0}^d U_i$ (direct sum). This sum is called the Φ -split decomposition of V . By [71, Theorem 4.6], the elements A, A^* act on $\{U_i\}_{i=0}^d$ as follows:

$$(19.137) \quad (A - \theta_i I)U_i \subseteq U_{i+1} \quad (0 \leq i \leq d-1), \quad (A - \theta_d I)U_d = 0,$$

$$(19.138) \quad (A^* - \theta_i^* I)U_i \subseteq U_{i-1} \quad (1 \leq i \leq d), \quad (A^* - \theta_0^* I)U_0 = 0.$$

By [71, Corollary 5.7], for $0 \leq i \leq d$ the dimensions of U_i, E_iV, E_i^*V are equal; denote this common dimension by ρ_i . By [71, Corollary 5.7], we have $\rho_i = \rho_{d-i}$. The sequence $(\rho_0, \rho_1, \dots, \rho_d)$ is called the *shape of Φ* . By [118, Theorem 1.3], the shape satisfies $\rho_i \leq \rho_0 \binom{d}{i}$ for $0 \leq i \leq d$. Additional results concerning the split decomposition and the shape can be found in [74, 97, 98, 106, 108, 112, 113, 140, 142, 157]. Some miscellaneous results about tridiagonal pairs and systems can be found in [1, 4, 28, 85, 87].

The tridiagonal system Φ is said to be *sharp* whenever $\rho_0 = 1$. If \mathbb{F} is algebraically closed, then Φ is sharp [116, Theorem 1.3]. In [72, Theorem 3.1], the sharp tridiagonal systems are classified up to isomorphism. This result makes heavy use of [82, 83, 114–119]. For the moment assume that Φ is sharp. By [114, Theorem 11.5] and [72, Theorem 3.1], there exists an antiautomorphism \dagger of $\text{End}(V)$ that fixes each of A, A^* . By linear algebra, there exists a nondegenerate symmetric bilinear form $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{F}$ such that $\langle Xu, v \rangle = \langle u, X^\dagger v \rangle$ for all $u, v \in V$ and $X \in \text{End}(V)$. See [6, 114, 130] for results on the bilinear form.

We now assume that $\rho_i = 1$ for $0 \leq i \leq d$. In this case, A, A^* is called a *Leonard pair*, and Φ is called a *Leonard system*. For some surveys on this topic, see [120, 138, 143]. For a Leonard pair, we can improve on the tridiagonal relations (19.135), (19.136) as follows. By [154, Theorem 1.5], there exist scalars ω, η, η^* in \mathbb{F} such that both

$$(19.139) \quad A^2A^* - \beta AA^*A + A^*A^2 - \gamma(AA^* + A^*A) - \varrho A^* = \gamma^*A^2 + \omega A + \eta I,$$

$$(19.140) \quad A^{*2}A - \beta A^*AA^* + AA^{*2} - \gamma^*(A^*A + AA^*) - \varrho^*A = \gamma A^{*2} + \omega A^* + \eta^* I.$$

These equations are called the *Askey–Wilson relations* (see [154, 166]). Observe that in (19.139) the right-hand side is a polynomial in A , and therefore commutes with A . Thus, (19.139) implies (19.135). Similarly, (19.140) implies (19.136). If A, A^* is TB, then the scalars $\gamma, \gamma^*, \omega, \eta, \eta^*$ are all zero and (19.139), (19.140) become (8.42), (8.43). In some cases, the Askey–Wilson relations can be put in \mathbb{Z}_3 -symmetric form (see [67, Theorem 10.1]). Additional results concerning the Askey–Wilson relations can be found in [158, 159]. We recall the Φ -split basis. Let $\{U_i\}_{i=0}^d$ denote the Φ -split decomposition of V . Pick any nonzero $v \in E_0^*V$. For $0 \leq i \leq d$ define $u_i = \tau_i(A)v$, where τ_i is from above Theorem 14.18. By [143, Section 21], we have

$0 \neq u_i \in U_i$. Consequently, the vectors $\{u_i\}_{i=0}^d$ form a basis for V , called the Φ -split basis for V . With respect to this basis, the matrices representing A, A^* take the form

$$A : \begin{pmatrix} \theta_0 & & & & \mathbf{0} \\ 1 & \theta_1 & & & \\ & 1 & \cdot & & \\ & & \cdot & \cdot & \\ \mathbf{0} & & & 1 & \theta_d \end{pmatrix}, \quad A^* : \begin{pmatrix} \theta_0^* & \varphi_1 & & & \mathbf{0} \\ & \theta_1^* & \varphi_2 & & \\ & & \cdot & \cdot & \\ & & & \cdot & \varphi_d \\ \mathbf{0} & & & & \theta_d^* \end{pmatrix},$$

where $\{\varphi_i\}_{i=1}^d$ are nonzero scalars in \mathbb{F} . The sequence $\{\varphi_i\}_{i=1}^d$ is called the *first split sequence* of Φ . The first split sequence $\{\phi_i\}_{i=1}^d$ of Φ^\downarrow is called the *second split sequence* of Φ . The sequence

$$(\{\theta_i\}_{i=0}^d; \{\theta_i^*\}_{i=0}^d; \{\varphi_i\}_{i=1}^d; \{\phi_i\}_{i=1}^d)$$

is called the *parameter array* of Φ . By [135, Lemma 3.11] Φ is determined up to isomorphism by its parameter array. In [135, Theorem 1.9], the Leonard systems are classified up to isomorphism in terms of the parameter array. In the TB case, the first and second split sequences look as follows. By [105, Theorem 1.5] and [143, Theorem 23.6],

$$\varphi_i = -\phi_i = (\theta_i^* - \theta_{i-1}^*)(\theta_0 + \theta_1 + \dots + \theta_{i-1}) \quad (1 \leq i \leq d).$$

In the table below, for $1 \leq i \leq d$ the scalar φ_i is displayed in closed form.

Case	φ_i
Example 13.3	$2hh^*i(i-d-1)$
Example 13.4	$hh^*(-1)^i(d+1)(d-2i+1) - hh^*(d-2i+1)^2$
Example 13.5	$hh^*(q^i - q^{-i})(q^{i-d-1} - q^{d-i+1})(q^{d-2i+1} + q^{2i-d-1})(q^2 - q^{-2})^{-2}$
Example 13.6	$hh^*(q^i - q^{-i})(q^{i-d-1} - q^{d-i+1})(q^{d-2i+1} + q^{2i-d-1})(q - q^{-1})^{-2}$

Another topic not discussed in earlier sections is the connection between Leonard systems and orthogonal polynomials. For $0 \leq i \leq d$ define a polynomial $u_i(x) \in \mathbb{F}[x]$ by

$$u_i(x) = \sum_{\ell=0}^i \frac{(x - \theta_0)(x - \theta_1) \cdots (x - \theta_{\ell-1})(\theta_i^* - \theta_0^*)(\theta_i^* - \theta_1^*) \cdots (\theta_i^* - \theta_{\ell-1}^*)}{\varphi_1 \varphi_2 \cdots \varphi_\ell}.$$

Define $U \in \text{Mat}_{d+1}(\mathbb{F})$ that has (i, j) -entry $u_i(\theta_j)$ for $0 \leq i, j \leq d$. By [139, Theorem 15.8], U is the transition matrix from an A^* -eigenbasis to an A -eigenbasis. By [141, Section 5], the polynomials $\{u_i(x)\}_{i=0}^d$ are from the terminating branch of the Askey scheme, which consists of the following polynomial families: q -Racah, q -Hahn, dual q -Hahn, q -Krawtchouk, dual q -Krawtchouk, affine q -Krawtchouk, quantum q -Krawtchouk, Racah, Hahn, dual Hahn, Krawtchouk, Bannai/Ito, Orphan. For a discussion of these polynomials, see [15, pp. 260–300]. In the TB case, Example 13.3 corresponds to a special case of Krawtchouk polynomials, Example 13.4 corresponds to a special case of Bannai/Ito polynomials, and Examples 13.5, 13.6 correspond to a special case of q -Racah polynomials. Some miscellaneous results about Leonard pairs and systems can be found in [36, 52–55, 100, 101, 107, 110, 111, 131, 137, 160, 161].

At the end of Section 16 and below Proposition 17.9, we mentioned the notion of a Leonard triple. This notion was introduced in [35, Definition 1.2]. A Leonard triple on V is a 3-tuple of elements in $\text{End}(V)$ such that for each map, there exists a basis for V with respect to which the matrix representing that map is diagonal and the matrices representing the other two maps are irreducible tridiagonal. To investigate Leonard triples, the following result is useful. For a Leonard pair A, A^* on V and for $X \in \text{End}(V)$, consider the matrices that represent X with respect to a standard eigenbasis for A and A^* . If these matrices are both tridiagonal, then X is a linear combination of I, A, A^*, AA^*, A^*A [109, Theorem 3.2]. Using this result, the Leonard triples have recently been classified up to isomorphism. The classification is summarized as follows. Using the eigenvalues one breaks down the analysis into four cases, called q -Racah, Racah, Krawtchouk, and Bannai/Ito [33, 61]. The Leonard triples are classified up to isomorphism in [67] (for q -Racah type); [45] (for Racah type); [89] (for Krawtchouk type); [65] (for Bannai/Ito type with even diameter); [63] (for Bannai/Ito type with odd diameter). Additional results on Leonard triples can be found in [60, 92, 99, 150, 155] (for q -Racah type); [9, 94, 125] (for Racah type); [14, 95] (for Krawtchouk type); [27, 32, 47, 48, 62, 66, 156] (for Bannai/Ito type); [102] (for general case).

We mention some algebras related to tridiagonal pairs and Leonard pairs. The *tridiagonal algebra* (see [136, Definition 3.9]) is defined by two generators subject to the tridiagonal relations (19.135), (19.136). The *Onsager algebra* (see [37, 38, 57, 129]) is the tridiagonal algebra for the case $\beta = 2, \gamma = 0, \gamma^* = 0, \varrho \neq 0, \varrho^* \neq 0$. The *q -Onsager algebra* (see [16, 19, 23, 151, 153]) is the tridiagonal algebra for the case $\beta = q^2 + q^{-2}, \gamma = 0, \gamma^* = 0, \varrho \neq 0, \varrho^* \neq 0$. The *Askey–Wilson algebra* (see [166]) is defined by two generators subject to the Askey–Wilson relations (19.139), (19.140). This algebra has a central extension (see [144]) that we now describe. Fix nonzero $q \in \mathbb{F}$ such that $q^4 \neq 1$. The *universal Askey–Wilson algebra* Δ_q is defined by generators and relations in the following way. The generators are A, B, C . The relations assert that each of

$$A + \frac{qBC - q^{-1}CB}{q^2 - q^{-2}}, \quad B + \frac{qCA - q^{-1}AC}{q^2 - q^{-2}}, \quad C + \frac{qAB - q^{-1}BA}{q^2 - q^{-2}},$$

is central in Δ_q . By [144, Theorem 3.1, Theorem 3.11] the group $\text{PSL}_2(\mathbb{Z})$ acts on Δ_q as a group of automorphisms, such that the $\text{PSL}_2(\mathbb{Z})$ -generator r (resp. s) sends $A \mapsto B \mapsto C \mapsto A$ (resp. $A \leftrightarrow B$). For more information on Δ_q , see [68–70, 144, 145, 155]. The double affine Hecke algebra (DAHA) for a reduced root system was defined by Cherednik (see [34]), and the definition was extended to include nonreduced root systems by Sahi (see [127]). The most general DAHA of rank 1 is said to have type (C_1^\vee, C_1) . In [86] a universal DAHA of type (C_1^\vee, C_1) was introduced and denoted \hat{H}_q . An injective algebra homomorphism $\Delta_q \rightarrow \hat{H}_q$ is given in [146, Section 4]; see also [86, 90, 91]. The paper [121] shows how \hat{H}_q is related to Leonard pairs. Additional results in the literature link tridiagonal pairs and Leonard pairs with the Lie algebra \mathfrak{sl}_2 (see [3, 7–9, 26, 81, 120]), the quantum algebras $U_q(\mathfrak{sl}_2)$ (see [2, 29, 30, 88, 147, 162]), $U_q(\widehat{\mathfrak{sl}}_2)$ (see [5, 24, 41, 75, 76, 84, 152]), the tetrahedron Lie algebra \boxtimes (see [25, 56, 79, 96]), and its q -deformation \boxtimes_q (see [42, 73, 77, 78, 80, 164, 165]).

Tridiagonal pairs have been used to investigate the open XXZ spin chain and related models in statistical mechanics (see [16–22]). The study of tridiagonal pairs has led to some conceptual advances, such as the bidiagonal pairs/triples (see [43, 44]), the billiard arrays (see [148, 163]), Hessenberg pairs (see [49–51]), and the lowering–raising triples (see [103, 104, 149]).

Acknowledgments. The authors thank Sarah Bockting-Conrad, Brian Curtin, Jae-ho Lee, and Hajime Tanaka for giving the paper a close reading and offering many valuable suggestions.

REFERENCES

- [1] H. Alnajjar. Self-duality of tridiagonal pairs of q -Serre type. *Int. Math. Forum*, 5:1409–1415, 2010.
- [2] H. Alnajjar. Leonard pairs associated with the equitable generators of the quantum algebra $U_q(\mathfrak{sl}_2)$. *Linear Multilinear Algebra* 59:1127–1142, 2011
- [3] H. Alnajjar. A tridiagonal linear map with respect to eigenbases of equitable basis of sl_2 . *Linear Multilinear Algebra* 63:1458–1467, 2015.
- [4] H. Alnajjar and B. Curtin. A family of tridiagonal pairs. *Linear Algebra Appl.*, 390:369–384, 2004.
- [5] H. Alnajjar and B. Curtin. A family of tridiagonal pairs related to the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$. *Electron. J. Linear Algebra*, 13:1–9, 2005.
- [6] H. Alnajjar and B. Curtin. A bilinear form for tridiagonal pairs of q -Serre type. *Linear Algebra Appl.*, 428:2688–2698, 2008.
- [7] H. Alnajjar and B. Curtin. Leonard pairs from the equitable basis of sl_2 . *Electron. J. Linear Algebra*, 20:490–505, 2010
- [8] H. Alnajjar and B. Curtin. A bidiagonal and tridiagonal linear map with respect to eigenbases of equitable basis of sl_2 . *Linear Multilinear Algebra*, 61:1668–1674, 2013.
- [9] H. Alnajjar and B. Curtin. Leonard triples from the equitable basis of \mathfrak{sl}_2 . *Linear Algebra Appl.*, 482:47–54, 2015.
- [10] R.C. Alperin. $PSL_2(\mathbb{Z}) = \mathbb{Z}_2 \star \mathbb{Z}_3$. *Am. Math. Monthly*, 100:385–386, 1993.
- [11] G. Andrews, R. Askey, and R. Roy. *Special Functions. Encyclopedia of Mathematics and Its Applications 71*. Cambridge University Press, 1999.
- [12] M. Arik and U. Kayserilioglu. The anticommutator spin algebra, its representations and quantum group invariance. *Int. J. Mod. Phys. A*, 18:5039–5045, 2003.
- [13] R. Askey and J. Wilson. A set of orthogonal polynomials that generalize the Racah coefficients or $6 - j$ symbols. *SIAM J. Math. Anal.*, 10:1008–1016, 1979.
- [14] J. Balmaceda and J.P. Maralit. Leonard triples from Leonard pairs constructed from the standard basis of the Lie algebra \mathfrak{sl}_2 . *Linear Algebra Appl.*, 437:1961–1977, 2012.
- [15] E. Bannai and T. Ito. *Algebraic Combinatorics I, Association Schemes*, Benjamin/Cummings Publishing Co. Inc., Menlo Park, CA, 1984.
- [16] P. Baseilhac. An integrable structure related with tridiagonal algebras. *Nucl. Phys. B*, 705:605–619, 2005.
- [17] P. Baseilhac. Deformed Dolan-Grady relations in quantum integrable models. *Nucl. Phys. B*, 709:491–521, 2005.
- [18] P. Baseilhac. The q -deformed analogue of the Onsager algebra: Beyond the Bethe Ansatz approach. *Nucl. Phys. B*, 754:309–328, 2006.
- [19] P. Baseilhac. A family of tridiagonal pairs and related symmetric functions. *J. Phys. A*, 39:11773–11791, 2006.
- [20] P. Baseilhac and S. Belliard. The half-infinite XXZ chain in Onsager’s approach. *Nucl. Phys. B*, 873:550–584, 2013.
- [21] P. Baseilhac and K. Koizumi. A deformed analogue of Onsager’s symmetry in the XXZ open spin chain. *J. Stat. Mech. Theory Exp.*, P10005:15 pp., 2005.
- [22] P. Baseilhac and K. Koizumi. Exact spectrum of the XXZ open spin chain from the q -Onsager algebra representation theory. *J. Stat. Mech. Theory Exp.*, P09006:27 pp., 2007.
- [23] P. Baseilhac and S. Kolb. Braid group action and root vectors for the q -Onsager algebra. *Transform. Groups*, 25:363–389, 2020.
- [24] G. Benkart and P. Terwilliger. Irreducible modules for the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ and its Borel subalgebra. *J. Algebra*, 282:172–194, 2004.
- [25] G. Benkart and P. Terwilliger. The universal central extension of the three-point \mathfrak{sl}_2 loop algebra. *Proc. Am. Math. Soc.*, 135(6):1659–1668, 2007.
- [26] G. Benkart and P. Terwilliger. The equitable basis for \mathfrak{sl}_2 . *Mathematische Zeitschrift*, 268:535–557, 2011.
- [27] H. De Bie, V. Genest, and L. Vinet. A Dirac-Dunkl equation on S^2 and the Bannai-Ito algebra. *Commun. Math. Phys.*, 344:447–464, 2016.
- [28] S. Bockting-Conrad. Two commuting operators associated with a tridiagonal pair. *Linear Algebra Appl.*, 437:242–270, 2012.
- [29] S. Bockting-Conrad. Tridiagonal pairs of q -Racah type, the double lowering operator ψ , and the quantum algebra $U_q(\mathfrak{sl}_2)$. *Linear Algebra Appl.*, 445:256–279, 2014.
- [30] S. Bockting-Conrad and P. Terwilliger. The algebra $U_q(\mathfrak{sl}_2)$ in disguise. *Linear Algebra Appl.*, 459:548–585, 2014.
- [31] A.E. Brower, A.M. Cohen, and A. Neumaier. *Distance-Regular Graphs*. Springer-Verlag, Berlin, 1989.
- [32] G.M.F. Brown. Totally bipartite/abipartite Leonard pairs and Leonard triples of Bannai/Ito type. *Electron. J. Linear Algebra*, 26:258–299, 2013.
- [33] S. Chang, S. Gao, and B. Hou. The Leonard triples with quantum parameter being not a unit root. *Linear Multilinear Algebra*, 63:2546–2560, 2015.

- [34] I. Cherednik. Double affine Hecke algebras, Knizhnik-Zamolodchikov equations, and Macdonald's operators. *Int. Math. Res. Not.*, 171–180, 1992.
- [35] B. Curtin. Modular Leonard triples, *Linear Algebra Appl.*, 424:510–539, 2007.
- [36] B. Curtin. Spin Leonard pairs. *Ramanujan J.*, 13:319–332, 2007.
- [37] B. Davies. Onsager's algebra and superintegrability. *J. Phys. A*, 23:2245–2261, 1990.
- [38] B. Davies. Onsager's algebra and the Dolan-Grady condition in the non-self-dual case. *J. Math. Phys.*, 32:2945–2950, 1991.
- [39] P. Delsarte. An algebraic approach to the association schemes of coding theory. *Philips Res. Rep. Suppl.*, 10, 1973.
- [40] D.B. Fairlie. Quantum deformations of $SU(2)$. *J. Phys. A*, 23, L183, 1990.
- [41] D. Funk-Neubauer. Raising/lowering maps and modules for the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$. *Commun. Algebra*, 35:2140–2159, 2007.
- [42] D. Funk-Neubauer. Tridiagonal pairs and the q -tetrahedron algebra. *Linear Algebra Appl.*, 431:903–925, 2009.
- [43] D. Funk-Neubauer. Bidiagonal pairs, the Lie algebra \mathfrak{sl}_2 , and the quantum group $U_q(\mathfrak{sl}_2)$. *J. Algebra Appl.*, 12:1250207, 46 pp, 2013.
- [44] D. Funk-Neubauer. Bidiagonal triples. *Linear Algebra Appl.*, 521:104–134, 2017.
- [45] S. Gao, Y. Wang and B. Hou. The classification of Leonard triples of Racah type. *Linear Algebra Appl.*, 439:1834–1861, 2013.
- [46] A.M. Gaviilik and A.U. Klimyk. q -Deformed orthogonal and pseudo-orthogonal algebras and their representations. *Lett. Math. Phys.*, 21:215–220, 1991.
- [47] V. Genest, L. Vinet, and A. Zhedanov. The Bannai-Ito polynomials as Racah coefficients of the $\mathfrak{sl}_{-1}(2)$ algebra. *Proc. Am. Math. Soc.*, 142:1545–1560, 2014.
- [48] V. Genest, L. Vinet, and A. Zhedanov. The Bannai-Ito algebra and a superintegrable system with reflections on the two-sphere. *J. Phys. A*, 47:205202, 13 pp, 2014.
- [49] A. Godjali. Hessenberg pairs of linear transformations. *Linear Algebra Appl.*, 431:1579–1586, 2009.
- [50] A. Godjali. Thin Hessenberg pairs. *Linear Algebra Appl.*, 432:3231–3249, 2010.
- [51] A. Godjali. Thin Hessenberg pairs and double Vandermonde matrices. *Linear Algebra Appl.*, 436:3018–3060, 2012.
- [52] E. Hanson. A characterization of Leonard pairs using the notion of a tail. *Linear Algebra Appl.*, 435:2961–2970, 2011.
- [53] E. Hanson. A characterization of Leonard pairs using the parameters $\{a_i\}_{i=0}^d$. *Linear Algebra Appl.*, 438:2289–2305, 2013.
- [54] E. Hanson. A characterization of bipartite Leonard pairs using the notion of a tail. *Linear Algebra Appl.*, 452:46–67, 2014.
- [55] B. Hartwig. Three mutually adjacent Leonard pairs. *Linear Algebra Appl.*, 408:19–39, 2005.
- [56] B. Hartwig. The Tetrahedron algebra and its finite-dimensional irreducible modules. *Linear Algebra Appl.*, 422:219–235, 2007.
- [57] B. Hartwig and P. Terwilliger. The Tetrahedron algebra, the Onsager algebra, and the \mathfrak{sl}_2 loop algebra. *J. Algebra*, 308:840–863, 2007.
- [58] M. Havliček, A.U. Klimyk, and S. Pošta. Representations of the cyclically symmetric q -deformed algebra $\mathfrak{so}_q(3)$. *J. Math. Phys.*, 40:2135–2161, 1999.
- [59] M. Havliček and S. Pošta. On the classification of irreducible finite-dimensional representations of $U'_q(\mathfrak{so}_3)$ algebra. *J. Math. Phys.*, 42:472–500, 2001.
- [60] B. Hou and S. Gao. Leonard pairs and Leonard triples of q -Racah type from the quantum algebra $U_q(\mathfrak{sl}_2)$. *Commun. Algebra*, 41:3762–3774, 2013.
- [61] B. Hou, Y. Liu, and S. Gao. The Leonard triples having classical type. *Linear Algebra Appl.*, 467:202–225, 2015.
- [62] B. Hou, M. Wang and S. Gao. The classification of finite-dimensional irreducible modules of Bannai/Ito algebra. *Commun. Algebra*, 44:919–943, 2016.
- [63] B. Hou, L. Wang, and S. Gao. The classification of Leonard triples that have Bannai/Ito type and odd diameter. *Linear Algebra Appl.*, 439:2667–2691, 2013.
- [64] B. Hou, J. Wang, and S. Gao. Totally bipartite Leonard pairs and totally bipartite Leonard triples of q -Racah type. *Linear Algebra Appl.*, 448:168–204, 2014.
- [65] B. Hou, Y. Xu, and S. Gao. The classification of Leonard triples of Bannai/Ito type with even diameter. *Commun. Algebra*, 43:3776–3798, 2015.
- [66] B. Hou, L. Zhang, and S. Gao. The Leonard triples extended from given Leonard pairs of Bannai/Ito type. *Linear Multilinear Algebra*, 62:1091–1104, 2014.
- [67] H. Huang. The classification of Leonard triples of QRacah type. *Linear Algebra Appl.*, 436:1442–1472, 2012.
- [68] H. Huang. Finite-dimensional irreducible modules of the universal Askey-Wilson algebra. *Commun. Math. Phys.*, 340:959–984, 2015.
- [69] H. Huang. Center of the universal Askey-Wilson algebra at roots of unity. *Nucl. Phys. B*, 909:260–296, 2016.

- [70] H. Huang. An embedding of the universal Askey-Wilson algebra into $U_q(\mathfrak{sl}_2) \otimes U_q(\mathfrak{sl}_2) \otimes U_q(\mathfrak{sl}_2)$. *Nucl. Phys. B*, 922:401–434, 2017.
- [71] T. Ito, K. Tanabe, and P. Terwilliger. Some algebra related to P - and Q -polynomial association schemes. In: *Codes and Association Schemes* (Piscataway NJ, 1999), American Mathematical Society, Providence RI, 167–192, 2001.
- [72] T. Ito, K. Nomura, and P. Terwilliger. A classification of sharp tridiagonal pairs. *Linear Algebra Appl.*, 435:1857–1884, 2011.
- [73] T. Ito, H. Rosengren, and P. Terwilliger. Evaluation modules for the q -tetrahedron algebra. *Linear Algebra Appl.*, 451:107–168, 2014.
- [74] T. Ito and P. Terwilliger. The shape of a tridiagonal pair. *J. Pure Appl. Algebra*, 188:145–160, 2004.
- [75] T. Ito and P. Terwilliger. Tridiagonal pairs and the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$. *Ramanujan J.*, 13:39–62, 2007.
- [76] T. Ito and P. Terwilliger. Two non-nilpotent linear transformations that satisfy the cubic q -Serre relations. *J. Algebra Appl.*, 6:477–503, 2007.
- [77] T. Ito and P. Terwilliger. The q -tetrahedron algebra and its finite dimensional irreducible modules. *Commun. Algebra*, 35:3415–3439, 2007.
- [78] T. Ito and P. Terwilliger. q -Inverting pairs of linear transformations and the q -tetrahedron algebra. *Linear Algebra Appl.*, 426:516–532, 2007.
- [79] T. Ito and P. Terwilliger. Finite-dimensional irreducible modules for the three-point \mathfrak{sl}_2 loop algebra. *Commun. Algebra*, 36:4557–4598, 2008.
- [80] T. Ito and P. Terwilliger. Distance-regular graphs and the q -tetrahedron algebra. *Eur. J. Comb.*, 30:682–697, 2009.
- [81] T. Ito and P. Terwilliger. Tridiagonal pairs of Krawtchouk type. *Linear Algebra Appl.*, 427:218–233, 2007.
- [82] T. Ito and P. Terwilliger. The Drinfel’d polynomial of a tridiagonal pair. *J. Comb., Inf. Syst. Sci.*, 34:255–292, 2009.
- [83] T. Ito and P. Terwilliger. Tridiagonal pairs of q -Racah type. *J. Algebra*, 322:68–93, 2009.
- [84] T. Ito and P. Terwilliger. The augmented tridiagonal algebra. *Kyushu J. Math.*, 64:81–144, 2010.
- [85] T. Ito and P. Terwilliger. How to sharpen a tridiagonal pair. *J. Algebra Appl.*, 9:543–552, 2010.
- [86] T. Ito and P. Terwilliger. Double affine Hecke algebras of rank 1 and the \mathbb{Z}_3 -symmetric Askey-Wilson relations. *Symmetry, Integr. Geometry Methods Appl.*, 6:065, 9 pages, 2010.
- [87] T. Ito and P. Terwilliger. Mock tridiagonal systems. *Linear Algebra Appl.*, 435:1997–2006, 2011.
- [88] T. Ito, P. Terwilliger and C. Weng. The quantum algebra $U_q(\mathfrak{sl}_2)$ and its equitable presentation. *J. Algebra*, 298:284–301, 2006.
- [89] N. Kang, B. Hou, and S. Gao. Leonard triples of Krawtchouk type. *Linear Multilinear Algebra*, 63:2341–2358, 2015.
- [90] T.H. Koornwinder. The relationship between Zhedanov’s algebra $AW(3)$ and the double affine Hecke algebra in the rank one case. *Symmetry, Integr. Geom. Methods Appl.*, 3:063, 15 pages, 2007
- [91] T.H. Koornwinder. Zhedanov’s algebra $AW(3)$ and the double affine Hecke algebra in the rank one case. II. The spherical subalgebra. *Symmetry, Integr. Geom. Methods Appl.*, 4:052,17 pages, 2008.
- [92] A. Korovnichenko and A. Zhedanov. Classical Leonard triples. In: M. Noumi and K. Takasaki (editors), *Elliptic Integrable Systems (Kyoto, 2004)*, Rokko Lectures in Mathematics, 18, Kobe University, 71–84, 2005.
- [93] D. Leonard. Orthogonal polynomials, duality, and association schemes. *SIAM J. Math. Anal.*, 13:656–663, 1982.
- [94] H. Liu, B. Hou and S. Gao. Leonard triples, the Racah algebra, and some distance-regular graphs of Racah type. *Linear Algebra Appl.*, 484:435–456, 2015.
- [95] S. Miklavic. Leonard triples and hypercubes. *J. Algebraic Comb.*, 28:397–424, 2008.
- [96] J.V.S. Morales and A.A. Pascasio. An action of the tetrahedron algebra on the standard module for the Hamming graphs and Doob graphs. *Graphs Comb.*, 30:1513–1527, 2014.
- [97] K. Nomura. A refinement of the split decomposition of a tridiagonal pair. *Linear Algebra Appl.*, 403:1–23, 2005.
- [98] K. Nomura. Tridiagonal pairs of height one. *Linear Algebra Appl.*, 403:118–142, 2005.
- [99] K. Nomura. Leonard pairs having LB-TD form. *Linear Algebra Appl.*, 455:1–21, 2014.
- [100] K. Nomura. The end-parameters of a Leonard pair. *Linear Algebra Appl.*, 462:88–109, 2014.
- [101] K. Nomura. Leonard pairs having specified end-entries. *Linear Algebra Appl.*, 465:43–64, 2015.
- [102] K. Nomura. Leonard pairs having zero-diagonal TD-TD form. *Linear Algebra Appl.*, 478:1–52, 2015.
- [103] K. Nomura. Linear transformations that are tridiagonal with respect to the three decompositions for an LR triple. *Linear Algebra Appl.*, 486:173–203, 2015.
- [104] K. Nomura. An LR pair that can be extended to an LR triple. *Linear Algebra Appl.*, 493:336–357, 2016.
- [105] K. Nomura and P. Terwilliger. Balanced Leonard pairs. *Linear Algebra Appl.*, 420:51–69, 2007.
- [106] K. Nomura and P. Terwilliger. Some trace formulae involving the split sequences of a Leonard pair. *Linear Algebra Appl.*, 413:189–201, 2006.
- [107] K. Nomura and P. Terwilliger. The determinant of $AA^* - A^*A$ for a Leonard pair A, A^* . *Linear Algebra Appl.*, 416:880–889, 2006.

- [108] K. Nomura and P. Terwilliger. Matrix units associated with the split basis of a Leonard pair. *Linear Algebra Appl.*, 418:775–787, 2006.
- [109] K. Nomura and P. Terwilliger. Linear transformations that are tridiagonal with respect to both eigenbases of a Leonard pair. *Linear Algebra Appl.*, 420:198–207, 2007.
- [110] K. Nomura and P. Terwilliger. The switching element for a Leonard pair. *Linear Algebra Appl.*, 428:1083–1108, 2008.
- [111] K. Nomura and P. Terwilliger. Affine transformations of a Leonard pair. *Electron. J. Linear Algebra*, 16:389–418, 2007.
- [112] K. Nomura and P. Terwilliger. The split decomposition of a tridiagonal pair. *Linear Algebra Appl.*, 424:339–345, 2007.
- [113] K. Nomura and P. Terwilliger. Transition maps between the 24 bases for a Leonard pair. *Linear Algebra Appl.*, 431:571–593, 2009.
- [114] K. Nomura and P. Terwilliger. Sharp tridiagonal pairs. *Linear Algebra Appl.*, 429:79–99, 2008.
- [115] K. Nomura and P. Terwilliger. Towards a classification of the tridiagonal pairs. *Linear Algebra Appl.*, 429:503–518, 2008.
- [116] K. Nomura and P. Terwilliger. The structure of a tridiagonal pair. *Linear Algebra Appl.*, 429:1647–1662, 2008.
- [117] K. Nomura and P. Terwilliger. Tridiagonal pairs and the μ -conjecture. *Linear Algebra Appl.*, 430:455–482, 2009.
- [118] K. Nomura and P. Terwilliger. On the shape of a tridiagonal pair. *Linear Algebra Appl.*, 432:615–636, 2010.
- [119] K. Nomura and P. Terwilliger. Tridiagonal pairs of q -Racah type and the μ -conjecture. *Linear Algebra Appl.*, 432:3201–3209, 2010.
- [120] K. Nomura and P. Terwilliger. Krawtchouk polynomials, the Lie algebra \mathfrak{sl}_2 , and Leonard pairs. *Linear Algebra Appl.*, 437:345–375, 2012.
- [121] K. Nomura and P. Terwilliger. The universal DAHA of type (C_1^\vee, C_1) and Leonard pairs of q -Racah type. *Linear Algebra Appl.*, 533:14–83, 2017.
- [122] K. Nomura and P. Terwilliger. Self-dual Leonard pairs. *Special Mat.*, 7:1–19, 2019.
- [123] A.V. Odesskii. An analog of the Sklyanin algebra. *Funct. Anal. Appl.*, 20:152–154, 1986.
- [124] W. Pauli. Zur Quantenmechanik des magnetischen Elektrons. *Zeitschrift für Physik*, 43:601–623, 1927.
- [125] S. Post. Racah polynomials and recoupling schemes of $\mathfrak{su}(1, 1)$. *Symmetry Integr. Geom. Methods Appl.*, 11:057, 17 pages, 2015.
- [126] J. Rotman. *Advanced Modern Algebra*, 2nd edition. American Mathematical Society, Providence, RI, 2010.
- [127] S. Sahi. Nonsymmetric Koornwinder polynomials and duality. *Ann. Math. (2)* 150:267–282, 1999.
- [128] R.M. Santilli. Imbedding of Lie algebras in nonassociative structures. *Nouvo Cimento*, 51:570–576, 1967.
- [129] K. Tanabe. The irreducible modules of the Terwilliger algebras of Doob schemes. *J. Algebraic Comb.*, 6:173–195, 1997.
- [130] H. Tanaka. A bilinear form relating two Leonard systems. *Linear Algebra Appl.*, 431:1726–1739, 2009.
- [131] H. Tanaka. The Erdős–Ko–Rado basis for a Leonard system. *Contributions to Discrete Mathematics*, 8:41–59, 2013.
- [132] P. Terwilliger. The subconstituent algebra of an association scheme I. *J. Algebraic Comb.*, 1:363–388, 1992.
- [133] P. Terwilliger. The subconstituent algebra of an association scheme II. *J. Algebraic Comb.*, 2:73–103, 1993.
- [134] P. Terwilliger. The subconstituent algebra of an association scheme III. *J. Algebraic Comb.*, 2:177–210, 1993.
- [135] P. Terwilliger. Two linear transformations each tridiagonal with respect to an eigenbasis of the other. *Linear Algebra Appl.*, 330:149–203, 2001.
- [136] P. Terwilliger. Two relations that generalize the q -Serre relations and the Dolan-Grady relations. In *Physics and Combinatorics 1999 (Nagoya)*, World Scientific Publishing, River Edge, NJ, 377–398, 2001.
- [137] P. Terwilliger. Leonard pairs from 24 points of view. *Rocky Mountain J. Math.*, 32:827–887, 2002.
- [138] P. Terwilliger. Introduction to Leonard pairs. OPSFA Rome 2001, *J. Computat. Appl. Math.*, 153:463–475, 2003.
- [139] P. Terwilliger. Leonard pairs and the q -Racah polynomials. *Linear Algebra Appl.*, 387:235–276, 2004.
- [140] P. Terwilliger. Two linear transformations each tridiagonal with respect to an eigenbasis of the other; comments on the split decomposition. *J. Computat. Appl. Math.*, 178:437–452, 2005.
- [141] P. Terwilliger. Two linear transformations each tridiagonal with respect to an eigenbasis of the other; comments on the parameter array. *Designs, Codes Cryptography*, 34:307–332, 2005.
- [142] P. Terwilliger. Two linear transformations each tridiagonal with respect to an eigenbasis of the other; the TD - D canonical form and the LB - UB canonical form. *J. Algebra*, 291:1–45, 2005.
- [143] P. Terwilliger. An algebraic approach to the Askey scheme of orthogonal polynomials. In: *Orthogonal Polynomials and Special Functions*, Lecture Notes in Math., 1883, Springer, Berlin, 255–330, 2006; [arXiv:math/0408390](https://arxiv.org/abs/math/0408390).
- [144] P. Terwilliger. The universal Askey-Wilson algebra. *Symmetry Integr. Geom. Methods Appl.*, 7:069, 24 pages, 2011.
- [145] P. Terwilliger. The universal Askey-Wilson algebra and the equitable presentation of $U_q(\mathfrak{sl}_2)$. *Symmetry Integr. Geom. Methods Appl.*, 7:099, 26 pages, 2011.
- [146] P. Terwilliger. The universal Askey-Wilson algebra and DAHA of type (C_1^\vee, C_1) . *Symmetry Integr. Geom. Methods Appl.*, 9:047, 40 pages, 2013.
- [147] P. Terwilliger. Finite-dimensional irreducible $U_q(\mathfrak{sl}_2)$ -modules from the equitable point of view. *Linear Algebra Appl.*, 439:358–400, 2013.
- [148] P. Terwilliger. Billiard arrays and finite-dimensional irreducible $U_q(\mathfrak{sl}_2)$ -modules. *Linear Algebra Appl.*, 461:21–270, 2014.

- [149] P. Terwilliger. Lowering-raising triples and $U_q(\mathfrak{sl}_2)$. *Linear Algebra Appl.*, 486:1–172, 2015.
- [150] P. Terwilliger. Leonard triples of q -Racah type and their pseudo intertwiners. *Linear Algebra Appl.*, 515:145–174, 2017.
- [151] P. Terwilliger. The q -Onsager algebra and the positive part of $U_q(\widehat{\mathfrak{sl}}_2)$. *Linear Algebra Appl.*, 521:19–56, 2017.
- [152] P. Terwilliger. Tridiagonal pairs of q -Racah type, the Bockling operator ψ , and L -operators for $U_q(L(\mathfrak{sl}_2))$. *ARS Mathematica Contemporanea*, 14:55–65, 2018.
- [153] P. Terwilliger. The Lusztig automorphism of the q -Onsager algebra. *J. Algebra*, 506:56–75, 2018.
- [154] P. Terwilliger and R. Vidunas. Leonard pairs and the Askey-Wilson relations. *J. Algebra Appl.*, 3:411–426, 2004.
- [155] P. Terwilliger and A. Žitnik. Distance-regular graphs of q -Racah type and the universal Askey-Wilson algebra. *J. Comb. Theory Ser. A*, 125:98–112, 2014.
- [156] S. Tsujimoto, L. Vinet, and A. Zhedanov. Dunkl shift operators and Bannai-Ito polynomials. *Adv. Math.*, 229:2123–2158, 2012.
- [157] M. Vidar. Tridiagonal pairs of shape $(1, 2, 1)$. *Linear Algebra Appl.*, 429:403–428, 2008.
- [158] R. Vidunas. Normalized Leonard pairs and Askey-Wilson relations. *Linear Algebra Appl.*, 422:39–57, 2007.
- [159] R. Vidunas. Askey-Wilson relations and Leonard pairs. *Disc. Math.*, 308:479–495, 2008.
- [160] Y. Wang, R. Hou, and S. Gao. Totally almost bipartite Leonard pairs and Leonard triples of q -Racah type. *Linear Multilinear Algebra*, 65:235–255, 2017.
- [161] Y. Wang, B. Hou, and S. Gao. Leonard triples extended from a given totally almost bipartite Leonard pair of Bannai/Ito type. *Linear Algebra Appl.*, 519:111–135, 2017.
- [162] C. Worawannotai. Dual polar graphs, the quantum algebra $U_q(\mathfrak{sl}_2)$, and Leonard systems of dual q -Krawtchouk type. *Linear Algebra Appl.*, 438:443–497, 2013.
- [163] Y. Yang. Upper triangular matrices and Billiard Arrays. *Linear Algebra Appl.*, 493:508–536, 2016.
- [164] Y. Yang. Some q -exponential formulas for finite-dimensional \square_q -modules. *Algebras Representation Theory*, 23:1–16, 2019.
- [165] Y. Yang. Finite-dimensional irreducible \square_q -modules and their Drinfel'd polynomials. *Linear Algebra Appl.*, 537:160–190, 2018.
- [166] A. Zhedanov. “Hidden symmetry” of Askey-Wilson polynomials. *Theoretical Math. Phys.*, 89:1146–1157, 1991.