

## SEMIDEFINITE GEOMETRY OF THE NUMERICAL RANGE\*

DIDIER HENRION<sup>†</sup>

**Abstract.** The numerical range of a matrix is studied geometrically via the cone of positive semidefinite matrices (or semidefinite cone for short). In particular, it is shown that the feasible set of a two-dimensional linear matrix inequality (LMI), an affine section of the semidefinite cone, is always dual to the numerical range of a matrix, which is therefore an affine projection of the semidefinite cone. Both primal and dual sets can also be viewed as convex hulls of explicit algebraic plane curve components. Several numerical examples illustrate this interplay between algebra, geometry and semidefinite programming duality. Finally, these techniques are used to revisit a theorem in statistics on the independence of quadratic forms in a normally distributed vector.

**Key words.** Numerical range, Semidefinite programming, LMI, Algebraic plane curves.

**AMS subject classifications.** 14H50, 14Q05, 47A12, 52A10, 90C22.

**1. Notations and definitions.** The numerical range of a matrix  $A \in \mathbb{C}^{n \times n}$  is defined as

$$(1.1) \quad \mathcal{W}(A) = \{w^*Aw \in \mathbb{C} : w \in \mathbb{C}^n, w^*w = 1\}.$$

It is a convex compact set of the complex plane which contains the spectrum of  $A$ . It is also called the field of values; see [8, Chapter 1] and [12, Chapter 1] for elementary introductions. Matlab functions for visualizing numerical ranges are freely available from [3] and [11].

Let

$$(1.2) \quad A_0 = I_n, \quad A_1 = \frac{A + A^*}{2}, \quad A_2 = \frac{A - A^*}{2i}$$

with  $I_n$  denoting the identity matrix of size  $n$  and  $i$  denoting the imaginary unit. Define

$$(1.3) \quad \mathcal{F}(A) = \{y \in \mathbb{P}_+^2 : F(y) = y_0A_0 + y_1A_1 + y_2A_2 \succeq 0\}$$

---

\*Received by the editors March 8, 2009. Accepted for publication May 28, 2010. Handling Editor: Shmuel Friedland.

<sup>†</sup>CNRS; LAAS; 7 avenue du colonel Roche, F-31077 Toulouse, France; Université de Toulouse; UPS, INSA, INP, ISAE; LAAS; F-31077 Toulouse, France (henrion@laas.fr); and Faculty of Electrical Engineering, Czech Technical University in Prague, Technická 4, CZ-16626 Prague, Czech Republic (henrion@fel.cvut.cz). Partially supported by project number 103/10/0628 of the Grant Agency of the Czech Republic.

with  $\succeq 0$  meaning positive semidefinite (since  $A_0, A_1, A_2$  are Hermitian matrices,  $F(y)$  has real eigenvalues for all  $y$ ) and  $\mathbb{P}_+^2$  denoting the oriented projective real plane (a model of the projective plane where the signs of homogeneous coordinates are significant, and which allows orientation, ordering and separation tests such as inequalities, see [18] for more details). Set  $\mathcal{F}(A)$  is a linear section of the cone of positive semidefinite matrices (or semidefinite cone for short) [1, Chapter 4]. Inequality  $F(y) \succeq 0$  is called a linear matrix inequality (LMI). In the complex plane  $\mathbb{C}$ , or equivalently, in the affine real plane  $\mathbb{R}^2$ , set  $\mathcal{F}(A)$  is a convex set including the origin, an affine section of the semidefinite cone.

Let

$$p(y) = \det(y_0 A_0 + y_1 A_1 + y_2 A_2)$$

be a trivariate form of degree  $n$  defining the algebraic plane curve

$$(1.4) \quad \mathcal{P} = \{y \in \mathbb{P}_+^2 : p(y) = 0\}.$$

Let

$$(1.5) \quad \mathcal{Q} = \{x \in \mathbb{P}_+^2 : q(x) = 0\}$$

be the algebraic plane curve dual to  $\mathcal{P}$ , in the sense that we associate to each point  $y \in \mathcal{P}$  a point  $x \in \mathcal{Q}$  of projective coordinates  $x = (\partial p(y)/\partial y_0, \partial p(y)/\partial y_1, \partial p(y)/\partial y_2)$ . Geometrically, a point in  $\mathcal{Q}$  corresponds to a tangent at the corresponding point in  $\mathcal{P}$ , and conversely; see [19, Section V.8] and [7, Section 1.1] for elementary properties of dual curves.

Let  $\mathbb{V}$  denote a vector space equipped with inner product  $\langle \cdot, \cdot \rangle$ . If  $x$  and  $y$  are vectors, then  $\langle x, y \rangle = x^* y$ . If  $X$  and  $Y$  are symmetric matrices, then  $\langle X, Y \rangle = \text{trace}(X^* Y)$ . Given a set  $\mathcal{K}$  in  $\mathbb{V}$ , its dual set consists of all linear maps from  $\mathcal{K}$  to non-negative elements in  $\mathbb{R}$ , namely,

$$\mathcal{K}^* = \{y \in \mathbb{V} : \langle x, y \rangle \geq 0, x \in \mathcal{K}\}.$$

Finally, the convex hull of a set  $\mathcal{K}$ , denoted  $\text{conv } \mathcal{K}$ , is the set of all convex combinations of elements in  $\mathcal{K}$ .

**2. Semidefinite duality.** After identifying  $\mathbb{C}$  with  $\mathbb{R}^2$  or  $\mathbb{P}_+^2$ , the first observation is that numerical range  $\mathcal{W}(A)$  is dual to LMI set  $\mathcal{F}(A)$ , and hence, it is an affine projection of the semidefinite cone.

LEMMA 2.1.  $\mathcal{W}(A) = \mathcal{F}(A)^* = \{(\langle A_0, W \rangle, \langle A_1, W \rangle, \langle A_2, W \rangle) \in \mathbb{P}_+^2 : W \in \mathbb{C}^{n \times n}, W \succeq 0\}$ .

*Proof.* The dual to  $\mathcal{F}(A)$  is

$$\begin{aligned}\mathcal{F}(A)^* &= \{x : \langle x, y \rangle = \langle F(y), W \rangle = \sum_k \langle A_k, W \rangle y_k \geq 0, W \succeq 0\} \\ &= \{x : x_k = \langle A_k, W \rangle, W \succeq 0\},\end{aligned}$$

an affine projection of the semidefinite cone. On the other hand, since  $w^*Aw = w^*A_1w + i w^*A_2w$ , the numerical range can be expressed as

$$\begin{aligned}\mathcal{W}(A) &= \{x = (w^*A_0w, w^*A_1w, w^*A_2w)\} \\ &= \{x : x_k = \langle A_k, W \rangle, W \succeq 0, \text{rank } W = 1\},\end{aligned}$$

the same affine projection as above, acting now on a subset of the semidefinite cone, namely, the non-convex variety of rank-one positive semidefinite matrices  $W = ww^*$ . Since  $w^*A_0w = 1$ , set  $\mathcal{W}(A)$  is compact, and  $\text{conv } \mathcal{W}(A) = \mathcal{F}(A)^*$ . The equality  $\mathcal{W}(A) = \mathcal{F}(A)^*$  follows from the Toeplitz-Hausdorff theorem establishing convexity of  $\mathcal{W}(A)$ ; see [12, Section 1.3] or [8, Theorem 1.1-2].  $\square$

Lemma 2.1 indicates that the numerical range has the geometry of planar projections of the semidefinite cone. In the terminology of [1, Chapter 4], the numerical range is semidefinite representable.

**3. Convex hulls of algebraic curves.** In this section, we notice that the boundaries of numerical range  $\mathcal{W}(A)$  and its dual LMI set  $\mathcal{F}(A)$  are subsets of algebraic curves  $\mathcal{P}$  and  $\mathcal{Q}$  defined respectively in (1.4) and (1.5), and explicitly given as loci of determinants of Hermitian pencils.

### 3.1. Dual curve.

LEMMA 3.1.  $\mathcal{F}(A)$  is the connected component delimited by  $\mathcal{P}$  around the origin.

*Proof.* A ray starting from the origin leaves LMI set  $\mathcal{F}(A)$  when the determinant  $p(y) = \det \sum_k y_k A_k$  vanishes. Therefore, the boundary of  $\mathcal{F}(A)$  is the subset of algebraic curve  $\mathcal{P}$  belonging to the convex connected component containing the origin.  $\square$

Note that  $\mathcal{P}$ , by definition, is the locus, or vanishing set of a determinant of a Hermitian pencil. Moreover, the pencil is definite at the origin so the corresponding polynomial  $p(y)$  satisfies a real zero (hyperbolicity) condition. Connected components delimited by such determinantal loci are studied in [9], where it is shown that they correspond to feasible sets of two-dimensional LMIs. A remarkable result of [9] is that every planar LMI set can be expressed this way. These LMI sets form a strict subset of planar convex basic semi-algebraic sets, called rigidly convex sets (see [9] for examples of convex basic semi-algebraic sets which are not rigidly convex). Rigidly convex sets are affine sections of the semidefinite cone.

### 3.2. Primal curve.

LEMMA 3.2.  $\mathcal{W}(A) = \text{conv } \mathcal{Q}$ .

*Proof.* From the proof of Lemma 2.1, a supporting line  $\{x : \sum_k x_k y_k = 0\}$  to  $\mathcal{W}(A)$  has coefficients  $y$  satisfying  $p(y) = 0$ . The boundary of  $\mathcal{W}(A)$  is therefore generated as an envelope of the supporting lines. See [16], [14, Theorem 10] and also [6, Theorem 1.3].  $\square$

$\mathcal{Q}$  is called the boundary generating curve of matrix  $A$  in [14]. An interesting feature is that, similarly to  $\mathcal{P}$ , curve  $\mathcal{Q}$  can be expressed as the locus of a determinant of a Hermitian pencil. In the case where  $\mathcal{Q}$  is irreducible (i.e., polynomial  $q(x)$  cannot be factored) and  $\mathcal{P}$  is not singular (i.e., there is no point in the complex projective plane such that the gradient of  $p(x)$  vanishes),  $q(x)$  can be written (up to a multiplicative constant) as the determinant of a symmetric pencil; see [6, Theorem 2.4]. Discrete differentials and Bézoutians can also be used to construct symmetric affine determinantal representations [10, Section 4.2]. Note however that the constructed pencils are not sign definite. Hence, the convex hull  $\mathcal{W}(A)$  is not a rigidly convex LMI set, and it cannot be an affine section of the semidefinite cone. However, as noticed in Lemma 2.1, it is an affine projection of the semidefinite cone.

## 4. Examples.

4.1. Rational cubic and quartic. Let

$$A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & i \\ 1 & i & 0 \end{bmatrix}.$$

Then

$$F(y) = \begin{bmatrix} y_0 & 0 & y_1 \\ 0 & y_0 + y_1 & y_2 \\ y_1 & y_2 & y_0 \end{bmatrix}$$

and

$$p(y) = (y_0 - y_1)(y_0 + y_1)^2 - y_0 y_2^2$$

defines a genus-zero cubic curve  $\mathcal{P}$  whose connected component containing the origin is the LMI set  $\mathcal{F}(A)$ ; see Figure 4.1. With an elimination technique (resultants or Gröbner basis with lexicographical ordering), we obtain

$$q(x) = 4x_1^4 + 32x_2^4 + 13x_1^2x_2^2 - 18x_0x_1x_2^2 + 4x_0x_1^3 - 27x_0^2x_2^2$$

defining the dual curve  $\mathcal{Q}$ , a genus-zero quartic with a cusp, whose convex hull is the numerical range  $\mathcal{W}(A)$ ; see Figure 4.1.




FIG. 4.1. Left: LMI set  $\mathcal{F}(A)$  (gray area) delimited by cubic  $\mathcal{P}$  (black). Right: numerical range  $\mathcal{W}(A)$  (gray area, dashed line) convex hull of quartic  $\mathcal{Q}$  (black solid line).

#### 4.2. Couple of two nested ovals. For

$$A = \begin{bmatrix} 0 & 2 & 1+2i & 0 \\ 0 & 0 & 1 & 0 \\ 0 & i & i & 0 \\ 0 & -1+i & i & 0 \end{bmatrix},$$

the quartic  $\mathcal{P}$  and its dual octic  $\mathcal{Q}$  both feature two nested ovals; see Figure 4.2. The




FIG. 4.2. Left: LMI set  $\mathcal{F}(A)$  (gray area) delimited by the inner oval of quartic  $\mathcal{P}$  (black line). Right: numerical range  $\mathcal{W}(A)$  (gray area) delimited by the outer oval of octic  $\mathcal{Q}$  (black line).

inner oval delimited by  $\mathcal{P}$  is rigidly convex, whereas the outer oval delimited by  $\mathcal{Q}$  is convex, but not rigidly convex.

**4.3. Cross and star.** A computer-generated representation of the numerical range as an envelope curve can be found in [8, Figure 1, p. 139] for

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ \frac{1}{2} & 0 & 0 & 0 \end{bmatrix}.$$

We obtain the quartic




FIG. 4.3. Left: LMI set  $\mathcal{F}(A)$  (gray area) delimited by the inner oval of quartic  $\mathcal{P}$  (black line). Right: numerical range  $\mathcal{W}(A)$  (gray area) delimited by the outer oval of twelfth-degree  $\mathcal{Q}$  (black line).

$$p(y) = \frac{1}{64}(64y_0^4 - 52y_0^2y_1^2 - 52y_0^2y_2^2 + y_1^4 + 34y_1^2y_2^2 + y_2^4)$$

and the dual twelfth-degree polynomial

$$\begin{aligned} q(x) = & 5184x_0^{12} - 299520x_0^{10}x_1^2 - 299520x_0^{10}x_2^2 + 1954576x_0^8x_1^4 \\ & + 16356256x_0^8x_1^2x_2^2 + 1954576x_0^8x_2^4 - 5375968x_0^6x_1^6 - 79163552x_0^6x_1^4x_2^2 \\ & - 79163552x_0^6x_1^2x_2^4 - 5375968x_0^6x_2^6 + 7512049x_0^4x_1^8 + 152829956x_0^4x_1^6x_2^2 \\ & - 2714586x_0^4x_1^4x_2^4 + 152829956x_0^4x_2^8 + 7512049x_0^4x_2^6 - 5290740x_0^2x_1^{10} \\ & - 136066372x_0^2x_1^8x_2^2 + 232523512x_0^2x_1^6x_2^4 + 232523512x_0^2x_1^4x_2^6 \\ & - 136066372x_0^2x_1^2x_2^8 - 5290740x_0^2x_2^{10} + 1498176x_1^{12} + 46903680x_1^{10}x_2^2 \\ & - 129955904x_1^8x_2^4 + 186148096x_1^6x_2^6 - 129955904x_1^4x_2^8 \\ & + 46903680x_1^2x_2^{10} + 1498176x_2^{12}, \end{aligned}$$

whose corresponding curves and convex hulls are represented in Figure 4.3.

**4.4. Decomposition into irreducible factors.** Consider the example of [8, Figure 6, p. 144] with

$$A = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The determinant of the trivariate pencil factors as follows:

$$p(y) = \frac{1}{256}(4y_0^3 - 3y_0y_1^2 - 3y_0y_2^2 + y_1^3 + y_1y_2^2)(4y_0^2 - y_1^2 - y_2^2)^3,$$

which means that the LMI set  $\mathcal{F}(A)$  is the intersection of a cubic and conic LMI.




FIG. 4.4. Left: LMI set  $\mathcal{F}(A)$  (gray area) intersection of cubic (black solid line) and conic (gray line) LMI sets. Right: numerical range  $\mathcal{W}(A)$  (gray area, black dashed line) convex hull of the union of a quartic curve (black solid line) and conic curve (gray line).

The dual curve  $\mathcal{Q}$  is the union of the quartic

$$\mathcal{Q}_1 = \{x : x_0^4 - 8x_0^3x_1 - 18x_0^2x_1^2 - 18x_0^2x_2^2 + 27x_1^4 + 54x_1^2x_2^2 + 27x_2^4 = 0\},$$

a cardioid dual to the cubic factor of  $p(y)$ , and the conic

$$\mathcal{Q}_2 = \{x : x_0^2 - 4x_1^2 - 4x_2^2 = 0\},$$

a circle dual to the quadratic factor of  $p(y)$ . The numerical range  $\mathcal{W}(A)$  is the convex hull of the union of  $\text{conv } \mathcal{Q}_1$  and  $\text{conv } \mathcal{Q}_2$ , which is here the same as  $\text{conv } \mathcal{Q}_1$ ; see Figure 4.4.

**4.5. Polytope.** Consider the example of [8, Figure 9, p. 147] with

$$A = \begin{bmatrix} 4 & 0 & 0 & -1 \\ -1 & 4 & 0 & 0 \\ 0 & -1 & 4 & 0 \\ 0 & 0 & -1 & 4 \end{bmatrix}.$$

The dual determinant factors into linear terms

$$p(y) = (y_0 + 5y_1)(y_0 + 3y_1)(y_0 + 4y_1 + y_2)(y_0 + 4y_1 - y_2)$$

and this generates an unbounded polyhedron  $\mathcal{F}(A) = \{y : y_0 + 5y_1 \geq 0, y_0 + 3y_1 \geq 0, y_0 + 4y_1 + y_2 \geq 0, y_0 + 4y_1 - y_2 \geq 0\}$ . The dual to curve  $\mathcal{P}$  is the union of the four




FIG. 4.5. Left: LMI set  $\mathcal{F}(A)$  (gray area), an unbounded polyhedron. Right: numerical range  $\mathcal{W}(A)$  (gray area) convex hull of four vertices.

points  $(1, 5, 0)$ ,  $(1, 3, 0)$ ,  $(1, 4, 1)$  and  $(1, 4, -1)$ , and hence, the numerical range  $\mathcal{W}(A)$  is the polytopic convex hull of these four vertices; see Figure 4.5.

**5. A problem in statistics.** We have seen in Example 4.5 that the numerical range can be polytopic, and this is the case in particular when  $A$  is a normal matrix (i.e., satisfying  $A^*A = AA^*$ ); see e.g. [14, Theorem 3] or [8, Theorem 1.4-4].

In this section, we study a problem that boils down to studying rectangular numerical ranges, i.e. polytopes with edges parallel to the main axes. Craig's theorem is a result from statistics on the stochastic independence of two quadratic forms in



variates following a joint normal distribution; see [4] for an historical account. In its simplest form (called the central case), the result can be stated as follows (in the sequel, we work in the affine plane  $y_0 = 1$ ):

**THEOREM 5.1.** *Let  $A_1$  and  $A_2$  be Hermitian matrices of size  $n$ . Then  $\det(I_n + y_1 A_1 + y_2 A_2) = \det(I_n + y_1 A_1) \det(I_n + y_2 A_2)$  if and only if  $A_1 A_2 = 0$ .*

*Proof.* If  $A_1 A_2 = 0$ , then obviously  $\det(I_n + y_1 A_1) \det(I_n + y_2 A_2) = \det((I_n + y_1 A_1)(I_n + y_2 A_2)) = \det(I_n + y_1 A_1 + y_2 A_2 + y_1 y_2 A_1 A_2) = \det(I_n + y_1 A_1 + y_2 A_2)$ . Let us prove the converse statement.

Let  $a_{1k}$  and  $a_{2k}$  respectively denote the eigenvalues of  $A_1$  and  $A_2$ , for  $k = 1, \dots, n$ . Then  $p(y) = \det(I_n + y_1 A_1 + y_2 A_2) = \det(I_n + y_1 A_1) \det(I_n + y_2 A_2) = \prod_k (1 + y_1 a_{1k}) \prod_k (1 + y_2 a_{2k})$  factors into linear terms, and we can write  $p(y) = \prod_k (1 + y_1 a_{1k} + y_2 a_{2k})$  with  $a_{1k} a_{2k} = 0$  for all  $k = 1, \dots, n$ . Geometrically, this means that the corresponding numerical range  $\mathcal{W}(A)$  for  $A = A_1 + i A_2$  is a rectangle with vertices  $(\min_k a_k, \min_k b_k)$ ,  $(\min_k a_k, \max_k b_k)$ ,  $(\max_k a_k, \min_k b_k)$  and  $(\max_k a_k, \max_k b_k)$ .

Following the terminology of [15],  $A_1$  and  $A_2$  satisfy property L since  $y_1 A_1 + y_2 A_2$  has eigenvalues  $y_1 a_{1k} + y_2 a_{2k}$  for  $k = 1, \dots, n$ . By [15, Theorem 2], it follows that  $A_1 A_2 = A_2 A_1$ , and hence that the two matrices are simultaneously diagonalisable: there exists a unitary matrix  $U$  such that  $U^* A_1 U = \text{diag}_k a_{1k}$  and  $U^* A_2 U = \text{diag}_k a_{2k}$ . Since  $a_{1k} a_{2k} = 0$  for all  $k$ , we have  $\sum_k a_{1k} a_{2k} = U^* A_1 U U^* A_2 U = U^* A_1 A_2 U = 0$ , and hence,  $A_1 A_2 = 0$ .  $\square$

**6. Conclusion.** The geometry of the numerical range, studied to a large extent by Kippenhahn in [14] – see [20] for an English translation with comments and corrections – is revisited here from the perspective of semidefinite programming duality. In contrast with previous studies of the geometry of the numerical range, based on differential topology [13], it is namely noticed that the numerical range is a semidefinite representable set, an affine projection of the semidefinite cone, whereas its geometric dual is an LMI set, an affine section of the semidefinite cone. The boundaries of both primal and dual sets are components of algebraic plane curves explicitly formulated as loci of determinants of Hermitian pencils. The geometry of the numerical range is therefore the geometry of (planar sections and projections of) the semidefinite cone, and hence every study of this cone is also relevant to the study of the numerical range.

The notion of numerical range can be generalized in various directions; for example in spaces of dimension greater than two, where it is non-convex in general [5]. Its convex hull is still representable as a projection of the semidefinite cone, and this was used extensively in the scope of robust control to derive computationally tractable but potentially conservative LMI stability conditions for uncertain linear systems, see e.g. [17]. The numerical range of three matrices is men-

tioned in [12, Section 1.8]. In this context, it would be interesting to derive conditions on three matrices  $A_1, A_2, A_3$  ensuring that  $\det(I_n + y_1 A_1 + y_2 A_2 + y_3 A_3) = \det(I_n + y_1 A_1) \det(I_n + y_2 A_2) \det(I_n + y_3 A_3)$ . Another extension of the numerical range to matrix polynomials (including matrix pencils) was carried out in [2], also using algebraic geometric considerations, and it could be interesting to study semidefinite representations of convex hulls of these numerical ranges.

The inverse problem of finding a matrix given its numerical range (as the convex hull of a given algebraic curve) seems to be difficult. In a sense, it is dual to the problem of finding a symmetric (or Hermitian) definite linear determinantal representation of a trivariate form: given  $p(y)$  satisfying a real zero (hyperbolicity) condition, find Hermitian matrices  $A_k$  such that  $p(y) = \det(\sum_k y_k A_k)$ , with  $A_0$  positive definite. Explicit formulas are described in [9] based on transcendental theta functions and Riemann surface theory, and the case of curves  $\{y : p(y) = 0\}$  of genus zero is settled in [10] using Bézoutians. A more direct and computationally viable approach in the positive genus case is still missing, and one may wonder whether the geometry of the dual object, namely, the numerical range  $\text{conv}\{x : q(x) = 0\}$ , could help in this context.

**Acknowledgment.** The author is grateful to Leiba Rodman for his suggestion of studying rigid convexity of the numerical range. This work also benefited from technical advice by Jean-Baptiste Hiriart-Urruty who recalled Theorem 5.1 in the September 2007 issue of the MODE newsletter of SMAI (French society for applied and industrial mathematics) and provided reference [4]. Finally, references [2, 13] were brought to the author's attention by an anonymous referee, who also pointed out several errors and inaccuracies in previous versions of this paper.

#### REFERENCES

- [1] A. Ben-Tal and A. Nemirovski. *Lectures on Modern Convex Optimization*. SIAM, Philadelphia, 2001.
- [2] M.T. Chien, H. Nakazato, and P. Psarrakos. Point equation of the boudary of the numerical range of a matrix polynomial. *Linear Algebra and its Applications*, 347:205–217, 2002.
- [3] C.C. Cowen and E. Harel. *An effective algorithm for computing the numerical range*. Technical report, Department of Mathematics, Purdue University, 1995.
- [4] M.F. Driscoll and W.R. Gungberg Jr. A history of the development of Craig's theorem. *The American Statistician*, 40(1):65–70, 1986.
- [5] M.K.H. Fan and A.L. Tits. On the generalized numerical range. *Linear and Multilinear Algebra*, 21(3):313–320, 1987.
- [6] M. Fiedler. Geometry of the numerical range of matrices. *Linear Algebra and its Applications*, 37:81–96, 1981.
- [7] I.M. Gelfand, M.M. Kapranov, and A.V. Zelevinsky. *Discriminants, Resultants and Multidimensional Determinants*. Birkhäuser, Boston, 1994.

- [8] K.E. Gustafsson and D.K.M. Rao. *Numerical Range: the Field of Values of Linear Operators and Matrices*. Springer, Berlin, 1997.
- [9] J.W. Helton and V. Vinnikov. Linear matrix inequality representation of sets. *Communications in Pure and Applied Mathematics*, 60(5):654–674, 2007.
- [10] D. Henrion. Detecting rigid convexity of bivariate polynomials. *Linear Algebra and its Applications*, 432:1218–1233, 2010.
- [11] N.J. Higham. *The Matrix Computation Toolbox*, Version 1.2, 2002.
- [12] R.A. Horn and C.R. Johnson. *Topics in Matrix Analysis*. Cambridge University Press, Cambridge, 1991.
- [13] E.A. Jonckheere, F. Ahmad, and E. Gutkin. Differential topology of numerical range. *Linear Algebra and its Applications*, 279:227–254, 1998.
- [14] R. Kippenhahn. Über den Wertevorrat einer Matrix. *Mathematische Nachrichten*, 6(3/4):193–228, 1951.
- [15] T.S. Motzkin and O. Taussky. Pairs of matrices with property L. *Transactions of the American Mathematical Society*, 73(1):108–114, 1952.
- [16] F.D. Murnaghan. On the field of values of a square matrix. *Proceedings of the National Academy of Sciences of the United States of America*, 18(3):246–248, 1932.
- [17] A. Packard and J. Doyle. The complex structured singular value. *Automatica*, 29(1):71–109, 1993.
- [18] J. Stolfi. *Primitives for Computational Geometry*. PhD Thesis, Department of Computer Science, Stanford University, California, 1988
- [19] R.J. Walker. *Algebraic Curves*. Princeton University Press, Princeton, 1950.
- [20] P.F. Zachlin and M.E. Hochstenbach. On the numerical range of a matrix. *Linear and Multilinear Algebra*, 56(1/2):185–225, 2008. English translation with comments and corrections of [14].