



SDD_1 TENSORS AND B_1 -TENSORS*

WEITING DUAN[†], DEKUN WEN[†], AND YAQIANG WANG[†]

Abstract. Strong \mathcal{H} -tensors play an important role in the fields of science and engineering. In this paper, we first propose a new subclass of strong \mathcal{H} -tensors that we call the class of SDD_1 tensors. We also prove that if a tensor is an SDD_1 tensor, it is a strong \mathcal{H} -tensor. As an application, a sufficient condition for an SDD_1 tensor to certify positive definiteness of even-order real symmetric tensors is proposed. Furthermore, we propose a new class of tensors by means of SDD_1 tensors, naming it B_1 -tensors, and show that B -tensors are a subclass of it. Meanwhile, some properties of B_1 -tensor were introduced. The numerical examples demonstrate the validity of our results.

Key words. \mathcal{H} -tensor, SDD_1 Tensor, real symmetric tensor, B_1 -tensor

AMS subject classifications. 15A15, 15A18, 15A69.

1. Introduction. Let n and m be integers, and $N = \{1, 2, \dots, n\}$ and \mathbb{C} (*resp.* \mathbb{R}) be the set of all complex (*resp.* real) numbers. We call $\mathcal{A} = (a_{i_1 i_2 \dots i_m})$ a complex (*resp.* real) tensor with order m and dimension n , if $a_{i_1 i_2 \dots i_m} \in \mathbb{C}$ (*resp.* \mathbb{R}), where $i_j = 1, 2, \dots, n$ for $j = 1, 2, \dots, m$. Let $\mathbb{C}^{[m, n]}$ (*resp.* $\mathbb{R}^{[m, n]}$) ($m, n \geq 2$) be the set of all complex (*resp.* real) order m dimension n tensors. In fact, a vector is a tensor of order 1 and a matrix is a tensor of order 2. Tensors, as a high-order extension of matrices, have important applications in many scientific fields, such as signal and image processing, continuum physics, magnetic resonance imaging, high-order statistics [1, 13, 17].

For a tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$, if there exists a complex number λ and a complex vector $x = (x_1, x_2, \dots, x_n)^T \neq (0, 0, \dots, 0)^T$ that are solutions of the following homogeneous polynomial equations:

$$\mathcal{A}x^{m-1} = \lambda x^{[m-1]},$$

then λ is called an eigenvalue of \mathcal{A} and x is its corresponding eigenvector [7, 8, 16], where $\mathcal{A}x^{m-1}$ and $\lambda x^{[m-1]}$ are vectors, and whose i th components are

$$(\mathcal{A}x^{m-1})_i = \sum_{i_2, \dots, i_m \in N} a_{i i_2 \dots i_m} x_{i_2} \cdots x_{i_m}, \text{ and } (x^{[m-1]})_i = x_i^{m-1}.$$

In particular, if λ and x are both real, then λ is called an \mathcal{H} -eigenvalue of \mathcal{A} and x is its corresponding \mathcal{H} -eigenvector [16]. The \mathcal{H} -eigenvalues of tensors are widely used in data analysis, high-order Markov chains, and positive definiteness of even-order homogeneous polynomials [5, 14, 15]. In this paper, we will discuss its applications to the positive definiteness of even-order homogeneous polynomials. The notions of homogeneous polynomials and positive definiteness are given in the following.

*Received by the editors on February 28, 2024. Accepted for publication on July 29, 2024. Handling Editor: Nick Vanieuwenhoven. Corresponding Author: Yaqiang Wang.

Funding: This work was partly supported by the National Natural Science Foundation of China (31600299), Natural Science Basic Research Program of Shaanxi, China (2020JM-622), and the Postgraduate Innovative Research Project of Baoji University of Arts and Sciences (YJSCX23YB33).

[†]School of Mathematics and Information Science, Baoji University of Arts and Sciences, Baoji, Shaanxi, 721013, P. R. China (weitingduan0393@163.com, dekun9926@163.com, yaqiangwang1004@163.com).

For an m th degree homogeneous polynomial of n variables, $f(x)$ can be usually denoted as:

$$f(x) \equiv \mathcal{A}x^m = \sum_{i_1, i_2, \dots, i_m \in N} a_{i_1 i_2 \dots i_m} x_{i_1} x_{i_2} \cdots x_{i_m},$$

where $x = (x_1, x_2, \dots, x_n)^T \in \mathbb{R}^n$ and $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$ is a symmetric tensor [15]. The tensor \mathcal{A} is called symmetric if its elements are invariant under any permutation of indices $\{i_1, i_2, \dots, i_m\}$ [16]. If m is even, and

$$f(x) > 0, \text{ for all } x \in \mathbb{R}^n, x \neq 0,$$

then we say that $f(x)$ is positive definite. The symmetric tensor \mathcal{A} is called positive definite if $f(x)$ is positive definite [15].

The positive definiteness of homogeneous polynomials play a key role in the stability study of nonlinear autonomous systems via Lyapunov's direct method in automatic control [15, 19, 26]. Concretely, for the system $\dot{x} = g(x)$, if a multivariate polynomial $f(x)$ can be found such that $f(x)$ is positive definite and

$$\left(\frac{\partial f}{\partial x}\right)^T g(x) < 0, \forall x \in \mathbb{R}^n, x \neq 0,$$

then the system $\dot{x} = g(x)$ is asymptotically stable. Hence, ascertaining whether a multivariate polynomial $f(x)$ is positive definite for all real x is often crucial to the use of Lyapunov stability tests [15]. To solve the problem of determining the positive definiteness of homogeneous polynomial, Qi proposed in [16] that $f(x)$ is positive definite if and only if the real symmetric tensor \mathcal{A} is positive definite, and Qi showed that even-order real symmetric tensor is positive definite if and only if all of its \mathcal{H} -eigenvalues are positive. However, it is hard to compute all \mathcal{H} -eigenvalues or the smallest \mathcal{H} -eigenvalue of \mathcal{A} when m and n are very large. Subsequently, a verifiable sufficient condition for positive definite tensors was proposed if a tensor has special structures; see Theorem 1.2. Before that, we give the definition of strong \mathcal{H} -tensors.

DEFINITION 1.1 (Ding, Qi, and Wei [2]). Let $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$. If there is a positive vector $x = (x_1, x_2, \dots, x_n)^T \in \mathbb{R}^n$ such that

$$|a_{ii \dots i}| x_i^{m-1} > \sum_{\substack{i_2, \dots, i_m \in N \\ \delta_{i i_2 \dots i_m} = 0}} |a_{i i_2 \dots i_m}| x_{i_2} \cdots x_{i_m}, \forall i \in N,$$

where $|a|$ is the modulus of $a \in \mathbb{C}$, then \mathcal{A} is called a strong \mathcal{H} -tensor.

THEOREM 1.2 (Li et al. [9]). Let $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ with $a_{kk \dots k} > 0$ for all $k \in N$ and m be even. If \mathcal{A} is a strong \mathcal{H} -tensor, then \mathcal{A} is positive definite.

With this theorem, for the positive definiteness of even-order real symmetric tensors, it is possible to first determine whether the given tensor is a strong \mathcal{H} -tensor. This article will propose a new subclass of strong \mathcal{H} -tensors, which are called SDD_1 tensors. Determining whether a given tensor is an SDD_1 tensor depends only on the elements of the tensor. This provides a sufficient condition for determining the positive definiteness of tensors. More criteria for judging whether a tensor is a strong \mathcal{H} -tensor have been widely proposed; see [3, 4, 10, 11, 12, 20, 21, 22, 23, 24, 25, 27].

We now give the structural arrangement of this paper. In Section 2, we will propose SDD_1 tensors, which are an extension of SDD_1 matrices. Meanwhile, we will prove that it is a subclass of strong \mathcal{H} -tensors.

As an application, a sufficient condition will be presented for using SDD_1 tensors to determine the positive definiteness of even-order real symmetric tensors. In Section 3, we propose B_1 -tensors, which include B -tensors as a subclass. At the same time, some properties of B_1 -tensors are introduced. Moreover, some numerical examples are presented to illustrate those new results. Finally, in Section 4, a brief conclusion is given.

2. SDD_1 tensors. In this section, a new class of tensors are proposed, which are called SDD_1 tensors. In particular, it is shown that SDD_1 tensors are strong \mathcal{H} -tensors. For the convenience of discussion, some notations, definitions, lemmas, and theorems are given in the following.

Calligraphic letters $\mathcal{A}, \mathcal{B}, \dots$ represent tensors; capital letters A, B, \dots denote matrices; lowercase letters x, y, \dots refer to vectors.

For a tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$, we denote

$$\begin{aligned} r_i(\mathcal{A}) &= \sum_{\substack{i_2 \dots i_m \in N^{m-1} \\ \delta_{i i_2 \dots i_m} = 0}} |a_{i i_2 \dots i_m}| = \sum_{i_2 \dots i_m \in N^{m-1}} |a_{i i_2 \dots i_m}| - |a_{i i \dots i}|, \\ N_L &= N_L(\mathcal{A}) = \{i \in N : |a_{i i \dots i}| \leq r_i(\mathcal{A})\}, \\ N_U &= N_U(\mathcal{A}) = \{i \in N : |a_{i i \dots i}| > r_i(\mathcal{A})\}, \\ N^{m-1} &= \{i_2 \dots i_m : i_j \in N, j = 2, \dots, m\}, \\ N^{m-1} \setminus S^{m-1} &= \{i_2 i_3 \dots i_m : i_2 i_3 \dots i_m \in N^{m-1} \text{ and } i_2 i_3 \dots i_m \notin S^{m-1}\}, \\ N_0^{m-1} &= N^{m-1} \setminus (N_L^{m-1} \cup N_U^{m-1}), \end{aligned}$$

where $r_i(\mathcal{A})$ is weight of the off-diagonal entries of the i th row of the flattening \mathcal{A} . N_L is the set of indices where the modulus of the diagonal entry is less than or equal to the corresponding off-diagonal weight. N_U is the set of indices where the modulus of the diagonal entry is greater than the corresponding off-diagonal weight. N^{m-1} is the set of indices in which i_2 to i_m belong to N . $N^{m-1} \setminus S^{m-1}$ denotes the difference set of N^{m-1} and S^{m-1} , where i_2 to i_m belong to N and i_2 to i_m does not belong to S . N_0^{m-1} denotes the difference set of N^{m-1} and $N_L^{m-1} \cup N_U^{m-1}$, where i_2 to i_m belong partly to N_L and partly to N_U .

DEFINITION 2.1 (Qi [16]). Let $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$. \mathcal{A} is called a diagonally dominant tensor if

$$|a_{i i \dots i}| \geq r_i(\mathcal{A}), \quad \forall i \in N.$$

\mathcal{A} is called a strictly diagonally dominant (SDD) tensor if all inequalities hold strictly.

DEFINITION 2.2 (Kolda and Bader [6]). Let $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$ and $X = \text{diag}(x_1, x_2, \dots, x_n)$. If

$$\mathcal{B} = (b_{i_1 i_2 \dots i_m}) = \mathcal{A}X^{m-1},$$

where

$$b_{i_1 i_2 \dots i_m} = a_{i_1 i_2 \dots i_m} x_{i_2} \dots x_{i_m}, \quad i_j \in N, \quad j \in \{1, 2, \dots, m\},$$

then we call \mathcal{B} the product of the tensor \mathcal{A} and the matrix X .

Let us introduce a new class of tensors containing SDD tensors, by imposing to the rows that are not strictly diagonally dominant a weaker condition.

DEFINITION 2.3. A tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$ is called an SDD_1 tensor if

$$|a_{i \dots i}| > p_i(\mathcal{A}), \quad \forall i \in N_L,$$

where

$$\begin{aligned} p_i(\mathcal{A}) = & \sum_{\substack{i_2 \dots i_m \in N_L^{m-1} \\ \delta_{i i_2 \dots i_m} = 0}} |a_{i i_2 \dots i_m}| + \sum_{\substack{i_2 \dots i_m \in N_U^{m-1} \\ \delta_{i i_2 \dots i_m} = 0}} \max_{j \in \{i_2, \dots, i_m\}} \left\{ \frac{r_j(\mathcal{A})}{|a_{j j \dots j}|} \right\} |a_{i i_2 \dots i_m}| \\ & + \sum_{\substack{i_2 \dots i_m \in N_0^{m-1} \\ \delta_{i i_2 \dots i_m} = 0}} |a_{i i_2 \dots i_m}|. \end{aligned}$$

Remark 2.4. Obviously, we have $p_i(\mathcal{A}) \leq r_i(\mathcal{A})$, so the SDD tensors form a subclass of the SDD_1 tensors.

In the following, we will give some lemmas, which will be used frequently in the subsequent proofs.

LEMMA 2.5 (Ding, Qi and Wei [2]). If $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$ is an SDD tensor, then \mathcal{A} is a strong \mathcal{H} -tensor.

LEMMA 2.6 (Wang and Song [21]). Let $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$. If there exists a positive diagonal matrix X such that $\mathcal{A}X^{m-1}$ is a strong \mathcal{H} -tensor, then \mathcal{A} is a strong \mathcal{H} -tensor.

LEMMA 2.7. A tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$ is an SDD_1 tensor if and only if $|a_{i \dots i}| > p_i(\mathcal{A})$ for all $i \in N$.

Proof. One implication is obvious. For the other one, let $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$ be an SDD_1 tensor. From Definition 2.3, we get that $|a_{i \dots i}| > p_i(\mathcal{A})$ for any $i \in N_L$ and for any $i \in N_U$, from the definition of N_U and $p_i(\mathcal{A})$, we have $|a_{i \dots i}| > r_i(\mathcal{A}) \geq p_i(\mathcal{A})$. Thus, there is $|a_{i \dots i}| > p_i(\mathcal{A})$ for all $i \in N$. \square

LEMMA 2.8. If a tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$ is an SDD_1 tensor and $N_L \neq \emptyset$, then we have $\sum_{\substack{i_2 \dots i_m \in N_U^{m-1} \\ \delta_{i i_2 \dots i_m} = 0}} |a_{i i_2 \dots i_m}| \neq 0$.

Proof. For a tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$, if $\sum_{\substack{i_2 \dots i_m \in N_U^{m-1} \\ \delta_{i i_2 \dots i_m} = 0}} |a_{i i_2 \dots i_m}| = 0$, then there is $r_i(\mathcal{A}) = p_i(\mathcal{A})$.

Because \mathcal{A} is SDD_1 tensor, then $|a_{i \dots i}| > p_i(\mathcal{A}) = r_i(\mathcal{A})$ for all $i \in N_L$, which contradicts the definition of N_L . The proof is complete. \square

In the following, we construct a positive diagonal matrix X that scales an SDD_1 tensor to an SDD tensor.

THEOREM 2.9. If a tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$ is an SDD_1 tensor, then \mathcal{A} is a strong \mathcal{H} -tensor.

Proof. Let a tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$ be an SDD_1 tensor, according to Lemma 2.7, $|a_{i \dots i}| > p_i(\mathcal{A})$ for all $i \in N$. Hence, we have

$$|a_{i \dots i}| - p_i(\mathcal{A}) > 0, \quad \forall i \in N,$$

and

$$1 - \frac{p_i(\mathcal{A})}{|a_{i \dots i}|} > 0, \quad \forall i \in N.$$

Then there exists a positive number $\varepsilon > 0$ such that

$$(2.1) \quad 0 < \varepsilon < \min \left\{ 1 - \frac{p_i(\mathcal{A})}{|a_{ii\dots i}|}, \frac{|a_{i\dots i}| - p_i(\mathcal{A})}{\sum_{\substack{i_2 \dots i_m \in N_U^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |a_{ii_2 \dots i_m}|} \right\},$$

if $\sum_{\substack{i_2 \dots i_m \in N_U^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |a_{ii_2 \dots i_m}| = 0$, then the corresponding fraction is defined to be ∞ . Construct a matrix $X = \text{diag}(x_1, x_2, \dots, x_n)$, where

$$x_i = \begin{cases} 1, & i \in N_L, \\ \left(\frac{p_i(\mathcal{A})}{|a_{ii\dots i}|} + \varepsilon \right)^{\frac{1}{m-1}}, & i \in N_U. \end{cases}$$

From inequality (2.1), we can obtain that $\frac{p_i(\mathcal{A})}{|a_{ii\dots i}|} + \varepsilon < \frac{p_i(\mathcal{A})}{|a_{ii\dots i}|} + (1 - \frac{p_i(\mathcal{A})}{|a_{ii\dots i}|}) = 1$, so $x_i \neq +\infty$, which shows that X is a positive diagonal matrix. Let $\mathcal{B} = (b_{i_1 i_2 \dots i_m}) = \mathcal{A}X^{m-1}$, then $b_{i_1 i_2 \dots i_m} = a_{i_1 i_2 \dots i_m} x_{i_2} \cdots x_{i_m}$, $\forall i_j \in N$, $j \in \{1, 2, \dots, m\}$.

Next, we will prove that \mathcal{B} is an SDD tensor:

$$\begin{aligned} r_i(\mathcal{B}) &= \sum_{\substack{i_2 \dots i_m \in N_L^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |b_{ii_2 \dots i_m}| + \sum_{\substack{i_2 \dots i_m \in N_U^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |b_{ii_2 \dots i_m}| + \sum_{i_2 \dots i_m \in N_0^{m-1}} |b_{ii_2 \dots i_m}| \\ &= \sum_{\substack{i_2 \dots i_m \in N_L^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |a_{ii_2 \dots i_m}| + \sum_{\substack{i_2 \dots i_m \in N_U^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |a_{ii_2 \dots i_m}| \left(\frac{p_{i_2}(\mathcal{A})}{|a_{i_2 \dots i_2}|} + \varepsilon \right)^{\frac{1}{m-1}} \cdots \\ &\quad \left(\frac{p_{i_m}(\mathcal{A})}{|a_{i_m \dots i_m}|} + \varepsilon \right)^{\frac{1}{m-1}} + \sum_{i_2 \dots i_m \in N_0^{m-1}} |a_{ii_2 \dots i_m}| x_{i_2} \cdots x_{i_m} \\ &\leq \sum_{\substack{i_2 \dots i_m \in N_L^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |a_{ii_2 \dots i_m}| + \sum_{\substack{i_2 \dots i_m \in N_U^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |a_{ii_2 \dots i_m}| \left(\frac{p_{i_2}(\mathcal{A})}{|a_{i_2 \dots i_2}|} + \varepsilon \right)^{\frac{1}{m-1}} \cdots \\ &\quad \left(\frac{p_{i_m}(\mathcal{A})}{|a_{i_m \dots i_m}|} + \varepsilon \right)^{\frac{1}{m-1}} + \sum_{i_2 \dots i_m \in N_0^{m-1}} |a_{ii_2 \dots i_m}| \\ &\leq \sum_{\substack{i_2 \dots i_m \in N_L^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |a_{ii_2 \dots i_m}| + \sum_{\substack{i_2 \dots i_m \in N_U^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |a_{ii_2 \dots i_m}| \left(\frac{r_{i_2}(\mathcal{A})}{|a_{i_2 \dots i_2}|} + \varepsilon \right)^{\frac{1}{m-1}} \cdots \\ &\quad \left(\frac{r_{i_m}(\mathcal{A})}{|a_{i_m \dots i_m}|} + \varepsilon \right)^{\frac{1}{m-1}} + \sum_{i_2 \dots i_m \in N_0^{m-1}} |a_{ii_2 \dots i_m}| \\ &\leq \sum_{\substack{i_2 \dots i_m \in N_L^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} |a_{ii_2 \dots i_m}| + \sum_{\substack{i_2 \dots i_m \in N_U^{m-1} \\ \delta_{ii_2 \dots i_m} = 0}} \max_{j \in \{i_2, \dots, i_m\}} \left\{ \frac{r_j(\mathcal{A})}{|a_{j \dots j}|} + \varepsilon \right\} |a_{ii_2 \dots i_m}| \end{aligned}$$

$$\begin{aligned}
 & + \sum_{i_2 \cdots i_m \in N_0^{m-1}} |a_{ii_2 \cdots i_m}| \\
 = & p_i(\mathcal{A}) + \varepsilon \sum_{\substack{i_2 \cdots i_m \in N_U^{m-1} \\ \delta_{ii_2 \cdots i_m} = 0}} |a_{ii_2 \cdots i_m}|.
 \end{aligned}$$

For any $i \in N_L$, according to Equation (2.1) and Lemma 2.8, we have

$$r_i(\mathcal{B}) \leq p_i(\mathcal{A}) + \varepsilon \sum_{\substack{i_2 \cdots i_m \in N_U^{m-1} \\ \delta_{ii_2 \cdots i_m} = 0}} |a_{ii_2 \cdots i_m}| < p_i(\mathcal{A}) + |a_{ii \cdots i}| - p_i(\mathcal{A}) = |a_{ii \cdots i}| = |b_{ii \cdots i}|.$$

And for any $i \in N_U$, we have

$$r_i(\mathcal{B}) \leq p_i(\mathcal{A}) + \varepsilon \sum_{\substack{i_2 \cdots i_m \in N_U^{m-1} \\ \delta_{ii_2 \cdots i_m} = 0}} |a_{ii_2 \cdots i_m}| < p_i(\mathcal{A}) + \varepsilon |a_{ii \cdots i}| = |b_{ii \cdots i}|.$$

Thus, we obtain $|b_{ii \cdots i}| > r_i(\mathcal{B})$ ($\forall i \in N$). That is, \mathcal{B} is a strictly diagonally dominant tensor so that it follows from Lemma 2.5 that \mathcal{B} is a strong \mathcal{H} -tensor. Furthermore, \mathcal{A} is a strong \mathcal{H} -tensor by Lemma 2.6. The proof is completed. \square

The following corollary is easily obtained from Theorems 1.2 and 2.9.

COROLLARY 2.10. *Let $\mathcal{A} = (a_{i_1 i_2 \cdots i_m}) \in \mathbb{R}^{[m, n]}$ be an even-order symmetric tensor, and $a_{pp \cdots p} > 0$ for all $p \in N$. If \mathcal{A} is an SDD_1 tensor, then \mathcal{A} is positive definite.*

Next, the relationships between the various tensor classes described in this section are easily obtained from Theorem 2.9 and Remark 2.4, see Figure 1.




FIG. 1. Relationships between some tensor classes.

We present an example of determining whether a given tensor is a strong \mathcal{H} -tensor using the definition of the SDD_1 tensor.

EXAMPLE 2.1. *Let us consider tensor $\mathcal{A} = (a_{ijk}) = [A(1, :, :), A(2, :, :), A(3, :, :)] \in \mathbb{C}^{[3, 3]}$, where*

$$A(1, :, :) = \begin{pmatrix} 6 & 0.5 & 0 \\ 0 & 2 & 0 \\ 0.5 & 1 & 20 \end{pmatrix}, \quad A(2, :, :) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 20 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad A(3, :, :) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 1 & 0 & 20 \end{pmatrix}.$$

Obviously,

$$|a_{111}| = 6, r_1(\mathcal{A}) = 24, |a_{222}| = 20, r_2(\mathcal{A}) = 4, |a_{333}| = 20 \text{ and } r_3(\mathcal{A}) = 4,$$

so $N_L = \{1\}$, $N_U = \{2, 3\}$. By calculation, we obtain

$$\max_{j \in N_U} \left\{ \frac{r_j(\mathcal{A})}{|a_{jjj}|} \right\} = \frac{1}{5},$$

when $i \in N_L$, we get

$$\begin{aligned} p_1(\mathcal{A}) &= \sum_{\substack{i_2 i_3 \in N_L^2 \\ \delta_{1i_2 i_3} = 0}} |a_{1i_2 i_3}| + \sum_{i_2 i_3 \in N_U^2} \max_{j \in \{i_2, i_3\}} \left\{ \frac{r_j(\mathcal{A})}{|a_{jjj}|} \right\} |a_{1i_2 i_3}| + \sum_{i_2 i_3 \in N_0^2} |a_{1i_2 i_3}| \\ &= 0 + \frac{23}{5} + 1 = \frac{28}{5} < 6 = |a_{111}|. \end{aligned}$$

Hence, \mathcal{A} satisfies the conditions of Theorem 2.9 so that \mathcal{A} is a strong \mathcal{H} -tensor.

In addition, an example of determining the positive definiteness of homogeneous polynomial of even degree is given.

EXAMPLE 2.2. Consider the following 4th-degree homogeneous polynomial

$$f(x) = \mathcal{A}x^4 = 17x_1^4 + 21x_2^4 + 24x_3^4 + 19x_4^4 - 8x_1^3x_4 + 12x_1^2x_2x_3 - 12x_2x_3^2x_4 - 24x_1x_2x_3x_4,$$

where $x = (x_1, x_2, x_3, x_4)^T$. Then we can obtain a symmetric tensor $\mathcal{A} = (a_{ijkl}) \in \mathbb{R}^{[4,4]}$, where

$$\begin{aligned} a_{1111} &= 17, a_{2222} = 21, a_{3333} = 24, a_{4444} = 19, \\ a_{1114} &= a_{1141} = a_{1411} = a_{4111} = -2, \\ a_{1123} &= a_{1132} = a_{1213} = a_{1312} = a_{1231} = a_{1321} = 1, \\ a_{2113} &= a_{2131} = a_{2311} = a_{3112} = a_{3121} = a_{3211} = 1, \\ a_{3234} &= a_{3243} = a_{3324} = a_{3342} = a_{3423} = a_{3432} = -1, \\ a_{2334} &= a_{2343} = a_{2433} = a_{4233} = a_{4323} = a_{4332} = -1, \\ a_{1234} &= a_{1243} = a_{1324} = a_{1342} = a_{1423} = a_{1432} = -1, \\ a_{2134} &= a_{2143} = a_{2314} = a_{2341} = a_{2413} = a_{2431} = -1, \\ a_{3124} &= a_{3142} = a_{3214} = a_{3241} = a_{3412} = a_{3421} = -1, \\ a_{4123} &= a_{4132} = a_{4213} = a_{4231} = a_{4312} = a_{4321} = -1, \end{aligned}$$

and others are zeros. Then,

$$\begin{aligned} |a_{1111}| &= 17, r_1(\mathcal{A}) = 18, |a_{2222}| = 21, r_2(\mathcal{A}) = 12, \\ |a_{3333}| &= 24, r_3(\mathcal{A}) = 15, |a_{4444}| = 19, r_4(\mathcal{A}) = 11, \end{aligned}$$

so $N_L = \{1\}$, $N_U = \{2, 3, 4\}$. By calculation, we obtain

$$\max_{j \in N_U} \left\{ \frac{r_j(\mathcal{A})}{|a_{jjj}|} \right\} = \frac{5}{8},$$

when $i \in N_L$, we get

$$\begin{aligned} p_1(\mathcal{A}) &= \sum_{\substack{i_2 i_3 i_4 \in N_L^3 \\ \delta_{1 i_2 i_3 i_4} = 0}} |a_{1 i_2 i_3 i_4}| + \sum_{i_2 i_3 i_4 \in N_U^3} \max_{j \in \{i_2, i_3, i_4\}} \left\{ \frac{r_j(\mathcal{A})}{|a_{jjjj}|} \right\} |a_{1 i_2 i_3 i_4}| \\ &\quad + \sum_{i_2 i_3 i_4 \in N_0^3} |a_{1 i_2 i_3 i_4}| \\ &= 0 + \frac{15}{4} + 12 = \frac{63}{4} < 17 = |a_{1111}|. \end{aligned}$$

Hence, \mathcal{A} satisfies the conditions of SDD_1 tensor, by Theorem 2.9, \mathcal{A} is a strong \mathcal{H} -tensor. Moreover, the diagonal element is greater than 0. Further, from Corollary 2.10, we have that \mathcal{A} is positive definite, that is $f(x)$ is positive definite.

3. B_1 -tensor. At the beginning of this section, we first recall the definition of the B -tensor.

DEFINITION 3.1 (Song and Qi [18]). A tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ is called a B -tensor if and only if for all $i \in N$,

$$\sum_{i_2, \dots, i_m \in N} a_{i i_2 \dots i_m} > 0,$$

and

$$\frac{1}{n^{m-1}} \left(\sum_{i_2, \dots, i_m \in N} a_{i i_2 \dots i_m} \right) > a_{i j_2 \dots j_m}, \quad \forall (j_2, \dots, j_m) \neq (i, \dots, i).$$

In this section, we will introduce a new tensor class inspired by SDD_1 tensors and call it the B_1 -tensor, which includes B -tensors as its subclass. Further, we introduce some of its properties. For the convenience of discussion, we also first provide some notations, definitions, and theorems in the following.

Given a tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$, for each row i , denote

$$(3.1) \quad r_i^+(\mathcal{A}) = \max\{0, a_{i j_2 \dots j_m} : (j_2, \dots, j_m) \neq (i, \dots, i)\}.$$

Let $\mathcal{A}^+ = (b_{i_1 i_2 \dots i_m})$ be the tensor defined as:

$$(3.2) \quad b_{i_1 i_2 \dots i_m} = a_{i_1 i_2 \dots i_m} - r_{i_1}^+(\mathcal{A}).$$

Clearly, \mathcal{A}^+ is a Z -tensor. A tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{C}^{[m, n]}$ is called a Z -tensor if and only if $a_{i_1 i_2 \dots i_m} \leq 0$ for all $(i_1, \dots, i_m) \neq (i, \dots, i)$ [26].

In this section, we listed the basic definition of a B -tensor. In [18], it was also proved that a B -tensor can be characterized by the following equivalent definition.

DEFINITION 3.2 (Song and Qi [18]). A tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ is called a B -tensor if and only if for each $i \in N$,

$$\sum_{i_2 \dots i_m \in N^{m-1}} |a_{i i_2 \dots i_m}| > n^{m-1} r_i^+(\mathcal{A}),$$

that is,

$$(a_{i i \dots i} - r_i^+(\mathcal{A})) > \sum_{(i_2, \dots, i_m) \neq (i, \dots, i)} (r_i^+(\mathcal{A}) - a_{i i_2 \dots i_m}) = r_i(\mathcal{A}^+).$$

Inspired by the SDD_1 tensor, in the following we give a new class of tensors and call it B_1 -tensors.

DEFINITION 3.3. A tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ is a B_1 -tensor if for all $i \in N$,

$$a_{i \dots i} - r_i^+(\mathcal{A}) > p_i(\mathcal{A}^+).$$

Next, we investigate some very interesting properties of B_1 -tensors.

PROPOSITION 3.4. If a tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ is B -tensor, then \mathcal{A} is a B_1 -tensor.

Proof. If $\mathcal{A} = (a_{i_1 i_2 \dots i_m})$ is B -tensor, then by Definition 3.2,

$$a_{i \dots i} - r_i^+(\mathcal{A}) > \sum_{\substack{i_2 \dots i_m \in N^{m-1} \\ \delta_{i i_2 \dots i_m} = 0}} (r_i^+(\mathcal{A}) - a_{i i_2 \dots i_m}) = r_i(\mathcal{A}^+).$$

However, $r_i(\mathcal{A}^+) \geq p_i(\mathcal{A}^+)$, and hence $a_{i \dots i} - r_i^+(\mathcal{A}) > p_i(\mathcal{A}^+)$, that is, \mathcal{A} is a B_1 -tensor. \square

Based on the definition of B_1 -tensors, it is easy to obtain the following properties.

PROPOSITION 3.5. Let tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ be a B_1 -tensor, then \mathcal{A}^+ is an SDD_1 tensor.

PROPOSITION 3.6. If tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ is a B_1 -tensor, then \mathcal{A}^+ has positive diagonal elements.

Proof. If tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m})$ is a B_1 -tensor, then we have $a_{i \dots i} - r_i^+(\mathcal{A}) > p_i(\mathcal{A}^+) \geq 0$, i.e., $a_{i \dots i} - r_i^+(\mathcal{A}) > 0$. The proof is complete. \square

From Proposition 3.6, we can obtain the following corollary.

COROLLARY 3.7. If tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ is a B_1 -tensor, then there must be $a_{i \dots i} > r_i^+(\mathcal{A})$.

PROPOSITION 3.8. A tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ is a B_1 -tensor if and only if \mathcal{A}^+ is an SDD_1 tensor with positive diagonal entries.

Proof. If tensor $\mathcal{A} = (a_{i_1 i_2 \dots i_m})$ is a B_1 -tensor, then by Propositions 3.5 and 3.6, \mathcal{A}^+ is an SDD_1 tensor with positive diagonal entries. Conversely, if \mathcal{A}^+ is an SDD_1 tensor, then $|a_{i \dots i} - r_i^+(\mathcal{A})| > p_i(\mathcal{A}^+)$. Since \mathcal{A}^+ has positive diagonal entries, that is, $a_{i \dots i} - r_i^+(\mathcal{A}) > 0$, then we have $a_{i \dots i} - r_i^+(\mathcal{A}) > p_i(\mathcal{A}^+)$. Therefore, \mathcal{A} is B_1 -tensor. \square

PROPOSITION 3.9. Let $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ be a Z -tensor with positive diagonal entries. Then \mathcal{A} is a B_1 -tensor if and only if \mathcal{A} is an SDD_1 tensor.

Proof. Since $\mathcal{A} = (a_{i_1 i_2 \dots i_m})$ is a Z -tensor, we have $r_i^+(\mathcal{A}) = 0$ for all $i \in N$, and $\mathcal{A}^+ = \mathcal{A}$. Then \mathcal{A} is a B_1 -tensor if and only if $a_{i \dots i} - r_i^+(\mathcal{A}) > p_i(\mathcal{A}^+)$, that is, $a_{i \dots i} > p_i(\mathcal{A})$; therefore, \mathcal{A} is an SDD_1 tensor. Conversely, if \mathcal{A} is an SDD_1 tensor, we have $|a_{i \dots i}| > p_i(\mathcal{A})$. Since \mathcal{A} has positive diagonal entries, then $a_{i \dots i} > p_i(\mathcal{A})$. Thus, $a_{i \dots i} - r_i^+(\mathcal{A}) > p_i(\mathcal{A}^+)$ can be obtained immediately. That is to say \mathcal{A} is a B_1 -tensor. \square

The following corollary can be obtained from Proposition 3.9 and Theorem 2.9.

COROLLARY 3.10. Let $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ be a Z -tensor with positive diagonal entries. If \mathcal{A} is a B_1 -tensor, then \mathcal{A} is a strong \mathcal{H} -tensor.

PROPOSITION 3.11. $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ is a B_1 -tensor if and only if \mathcal{A}^+ is a B_1 -tensor.

Proof. By Proposition 3.8, $\mathcal{A} = (a_{i_1 i_2 \dots i_m})$ is B_1 -tensor if and only if \mathcal{A}^+ is an SDD_1 tensor with positive diagonal entries. Since \mathcal{A}^+ is a Z -tensor, according to Proposition 3.9, the conclusion follows immediately. \square

PROPOSITION 3.12. *If $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m,n]}$ is a B_1 -tensor, and $\mathcal{D} \in \mathbb{R}^{[m,n]}$ is a nonnegative diagonal tensor of the same order and dimension, then $\mathcal{A} + \mathcal{D}$ is a B_1 -tensor.*

Proof. Let $\mathcal{D} = (d_{i_1 i_2 \dots i_m})$, where

$$d_{i_2 \dots i_m} = \begin{cases} d_i, & (i_2, \dots, i_m) = (i, \dots, i), \\ 0, & \text{otherwise,} \end{cases}$$

and $d_i \geq 0$. Let $\mathcal{C} = \mathcal{A} + \mathcal{D}$, where

$$c_{i_2 \dots i_m} = \begin{cases} a_{i \dots i} + d_i, & (i_2, \dots, i_m) = (i, \dots, i), \\ a_{i_2 \dots i_m}, & \text{otherwise.} \end{cases}$$

Then, \mathcal{A} and \mathcal{C} have the same nondiagonal elements so that $r_i^+(\mathcal{A}) = r_i^+(\mathcal{C})$.

Next, let us prove $c_{i \dots i} - r_i^+(\mathcal{C}) > p_i(\mathcal{C}^+)$ for all $i \in N$, where $\mathcal{C}^+ = (v_{i_1 i_2 \dots i_m})$ with $v_{i_2 \dots i_m} = c_{i_2 \dots i_m} - r_i^+(\mathcal{C})$.

Since \mathcal{A} is a B_1 -tensor, for all $i \in N$, we have

$$(3.3) \quad c_{i \dots i} - r_i^+(\mathcal{C}) = a_{i \dots i} + d_i - r_i^+(\mathcal{A}) \geq a_{i \dots i} - r_i^+(\mathcal{A}) > p_i(\mathcal{A}^+) \geq 0.$$

Meanwhile, we have the following equation that holds:

$$\begin{aligned} r_i(\mathcal{C}^+) &= \sum_{\substack{i_2 \dots i_m \in N^{m-1} \\ \delta_{i_2 \dots i_m} = 0}} |v_{i_2 \dots i_m}| = \sum_{\substack{i_2 \dots i_m \in N^{m-1} \\ \delta_{i_2 \dots i_m} = 0}} |c_{i_2 \dots i_m} - r_i^+(\mathcal{C})| \\ &= \sum_{\substack{i_2 \dots i_m \in N^{m-1} \\ \delta_{i_2 \dots i_m} = 0}} |a_{i_2 \dots i_m} - r_i^+(\mathcal{A})| = r_i(\mathcal{A}^+). \end{aligned}$$

Hence, for $\forall i \in N_L(\mathcal{C}^+)$, we have $|c_{i \dots i} - r_i^+(\mathcal{C})| \leq r_i(\mathcal{C}^+)$, that is, $|a_{i \dots i} + d_i - r_i^+(\mathcal{A})| \leq r_i(\mathcal{A}^+)$. We can immediately obtain that $|a_{i \dots i} - r_i^+(\mathcal{A})| \leq r_i(\mathcal{A}^+)$. Therefore, $i \in N_L(\mathcal{A}^+)$. This indicates that $N_L(\mathcal{C}^+) \subset N_L(\mathcal{A}^+)$. The same method can illustrate that $N_U(\mathcal{A}^+) \subset N_U(\mathcal{C}^+)$.

For $\forall i \in N$,

$$\begin{aligned} &p_i(\mathcal{C}^+) \\ &= \sum_{\substack{i_2 \dots i_m \in N_L^{m-1}(\mathcal{C}^+) \\ \delta_{i_2 \dots i_m} = 0}} |v_{i_2 \dots i_m}| + \sum_{\substack{i_2 \dots i_m \in N_U^{m-1}(\mathcal{C}^+) \\ \delta_{i_2 \dots i_m} = 0}} \max_{j \in \{i_2, \dots, i_m\}} \left\{ \frac{r_j(\mathcal{C}^+)}{|v_{j \dots j}|} \right\} |v_{i_2 \dots i_m}| \\ &\quad + \sum_{\substack{i_2 \dots i_m \in N_0^{m-1}(\mathcal{C}^+) \\ \delta_{i_2 \dots i_m} = 0}} |v_{i_2 \dots i_m}| \\ &= \sum_{\substack{i_2 \dots i_m \in N_L^{m-1}(\mathcal{C}^+) \\ \delta_{i_2 \dots i_m} = 0}} |c_{i_2 \dots i_m} - r_i^+(\mathcal{C})| + \sum_{\substack{i_2 \dots i_m \in N_U^{m-1}(\mathcal{C}^+) \\ \delta_{i_2 \dots i_m} = 0}} \max_{j \in \{i_2, \dots, i_m\}} \left\{ \frac{r_j(\mathcal{C}^+)}{|c_{j \dots j} - r_j^+(\mathcal{C})|} \right\} \\ &\quad |c_{i_2 \dots i_m} - r_i^+(\mathcal{C})| + \sum_{\substack{i_2 \dots i_m \in N_0^{m-1}(\mathcal{C}^+) \\ \delta_{i_2 \dots i_m} = 0}} |c_{i_2 \dots i_m} - r_i^+(\mathcal{C})| \end{aligned}$$

$$\begin{aligned}
 &= \sum_{\substack{i_2 \cdots i_m \in N_L^{m-1}(\mathcal{C}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} |a_{ii_2 \cdots i_m} - r_i^+(\mathcal{A})| + \sum_{\substack{i_2 \cdots i_m \in N_U^{m-1}(\mathcal{C}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} \max_{j \in \{i_2, \dots, i_m\}} \left\{ \frac{r_j(\mathcal{A}^+)}{|a_{jj \cdots j} + d_j - r_j^+(\mathcal{A})|} \right\} \\
 &\quad |a_{ii_2 \cdots i_m} - r_i^+(\mathcal{A})| + \sum_{\substack{i_2 \cdots i_m \in N_0^{m-1}(\mathcal{C}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} |a_{ii_2 \cdots i_m} - r_i^+(\mathcal{A})| \\
 &\leq \sum_{\substack{i_2 \cdots i_m \in N_L^{m-1}(\mathcal{A}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} |a_{ii_2 \cdots i_m} - r_i^+(\mathcal{A})| + \sum_{\substack{i_2 \cdots i_m \in N_U^{m-1}(\mathcal{A}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} \max_{j \in \{i_2, \dots, i_m\}} \left\{ \frac{r_j(\mathcal{A}^+)}{|a_{jj \cdots j} - r_j^+(\mathcal{A})|} \right\} \\
 &\quad |a_{ii_2 \cdots i_m} - r_i^+(\mathcal{A})| + \sum_{\substack{i_2 \cdots i_m \in N_0^{m-1}(\mathcal{A}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} |a_{ii_2 \cdots i_m} - r_i^+(\mathcal{A})| \\
 &= p_i(\mathcal{A}^+).
 \end{aligned}$$

Combining equation (3.3), there is $c_{i \dots i} - r_i^+(\mathcal{C}) > p_i(\mathcal{C}^+)$, and this proof is completed. \square

PROPOSITION 3.13. *If $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ is a B_1 -tensor, then we can write \mathcal{A} as $\mathcal{A} = \mathcal{B} + \mathcal{C}$, where \mathcal{B} is a Z -tensor with positive diagonal entries and \mathcal{C} is a nonnegative tensor with $c_{ii_2 \dots i_m} = c_i$, $c_i \geq 0$, $\forall i \in N$. In particular, if \mathcal{A} is a B_1 -tensor and Z -tensor, then \mathcal{C} is a zero tensor.*

Proof. Suppose $\mathcal{B} = (b_{i_1 i_2 \dots i_m})$, where $b_{ii_2 \dots i_m} = a_{ii_2 \dots i_m} - r_i^+(\mathcal{A})$. According to the definition of $r_i^+(\mathcal{A})$, it can be inferred that $b_{ii_2 \dots i_m} \leq 0$, $(i_2, \dots, i_m) \neq (i, \dots, i)$. And since \mathcal{A} is a B_1 -tensor, $b_{i \dots i} = a_{i \dots i} - r_i^+(\mathcal{A}) > p_i(\mathcal{B}) \geq 0$. Then \mathcal{B} is a Z -tensor with positive diagonal entries.

Let $\mathcal{C} = (c_{i_1 i_2 \dots i_m})$ with $c_{ii_2 \dots i_m} = r_i^+(\mathcal{A})$, obviously \mathcal{C} is a nonnegative tensor with $c_i = r_i^+(\mathcal{A})$. In particular, if \mathcal{A} is a B_1 -tensor and Z -tensor, we have $r_i^+(\mathcal{A}) = 0$, then \mathcal{C} is a zero tensor. \square

PROPOSITION 3.14. *Let $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ be a B_1 -tensor, and let $\mathcal{C} = (c_{i_1 i_2 \dots i_m})$ be a nonnegative tensor of the form $c_{ii_2 \dots i_m} = c_i$, then $\mathcal{A} + \mathcal{C}$ is a B_1 -tensor.*

Proof. Let $\mathcal{B} = \mathcal{A} + \mathcal{C}$, where $\mathcal{B} = (b_{i_1 i_2 \dots i_m})$. By Proposition 3.11, we need to prove that $\mathcal{B}^+ = (d_{i_1 i_2 \dots i_m})$ is a B_1 -tensor. By definition, we can see that for all $i \in N$,

$$r_i^+(\mathcal{B}) = r_i^+(\mathcal{A}) + c_i.$$

On the other hand, for all $i, i_2, \dots, i_m \in N$, we have

$$d_{ii_2 \dots i_m} = b_{ii_2 \dots i_m} - r_i^+(\mathcal{B}) = (a_{ii_2 \dots i_m} + c_i) - (r_i^+(\mathcal{A}) + c_i) = a_{ii_2 \dots i_m} - r_i^+(\mathcal{A}).$$

This means that $\mathcal{B}^+ = \mathcal{A}^+$. Since \mathcal{A} is a B_1 -tensor, by Proposition 3.11, \mathcal{A}^+ is a B_1 -tensor. The conclusion follows immediately. \square

PROPOSITION 3.15. *If $\mathcal{A} = (a_{i_1 i_2 \dots i_m}) \in \mathbb{R}^{[m, n]}$ is a B_1 -tensor, then every principal sub-tensor of \mathcal{A} is also a B_1 -tensor.*

Proof. Let J be a nonempty subset of N with $|J| = r$, $\mathcal{C} = \mathcal{A}_r^J \in \mathbb{R}^{[m, r]}$ be the principal sub-tensor of \mathcal{A} and $\mathcal{A}^+ = (b_{i_1 i_2 \dots i_m})$. Since \mathcal{A} is a B_1 -tensor, we have $a_{i \dots i} - r_i^+(\mathcal{A}) > p_i(\mathcal{A}^+) \geq 0$. Thus, $a_{i \dots i} > r_i^+(\mathcal{A}) \geq r_i^+(\mathcal{C})$. Next, let us prove $a_{i \dots i} - r_i^+(\mathcal{C}) > p_i(\mathcal{C}^+)$ for all $i \in J$. We have

$$\begin{aligned} r_i(\mathcal{A}^+) &= \sum_{\substack{i_2 \cdots i_m \in N^{m-1} \\ \delta_{ii_2 \cdots i_m} = 0}} |a_{ii_2 \cdots i_m} - r_i^+(\mathcal{A})| = \sum_{\substack{i_2 \cdots i_m \in N^{m-1} \\ \delta_{ii_2 \cdots i_m} = 0}} (r_i^+(\mathcal{A}) - a_{ii_2 \cdots i_m}) \\ &\geq \sum_{\substack{i_2 \cdots i_m \in J^{m-1} \\ \delta_{ii_2 \cdots i_m} = 0}} (r_i^+(\mathcal{C}) - a_{ii_2 \cdots i_m}) = r_i(\mathcal{C}^+). \end{aligned}$$

For $\forall i \in N_L(\mathcal{C}^+)$, we have $0 \leq a_{i \dots i} - r_i^+(\mathcal{C}) \leq r_i(\mathcal{C}^+) \leq r_i(\mathcal{A}^+)$, and hence $0 \leq a_{i \dots i} - r_i^+(\mathcal{A}) \leq r_i(\mathcal{A}^+)$. Therefore, $i \in N_L(\mathcal{A}^+)$. This indicates that $N_L(\mathcal{C}^+) \subset N_L(\mathcal{A}^+)$. The same method can illustrate that $N_U(\mathcal{A}^+) \subset N_U(\mathcal{C}^+)$.

For $\forall i \in J$,

$$\begin{aligned} &a_{i \dots i} - r_i^+(\mathcal{C}) \\ &\geq a_{i \dots i} - r_i^+(\mathcal{A}) > p_i(\mathcal{A}^+) \\ &= \sum_{\substack{i_2 \cdots i_m \in N_L^{m-1}(\mathcal{A}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} |b_{ii_2 \cdots i_m}| + \sum_{\substack{i_2 \cdots i_m \in N_U^{m-1}(\mathcal{A}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} \max_{j \in \{i_2, \dots, i_m\}} \left\{ \frac{r_j(\mathcal{A}^+)}{|b_{jj \dots j}|} \right\} |b_{ii_2 \cdots i_m}| \\ &\quad + \sum_{\substack{i_2 \cdots i_m \in N_0^{m-1}(\mathcal{A}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} |b_{ii_2 \cdots i_m}| \\ &= \sum_{\substack{i_2 \cdots i_m \in N_L^{m-1}(\mathcal{A}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} (r_i^+(\mathcal{A}) - a_{ii_2 \cdots i_m}) + \sum_{\substack{i_2 \cdots i_m \in N_U^{m-1}(\mathcal{A}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} \max_{j \in \{i_2, \dots, i_m\}} \left\{ \frac{r_j(\mathcal{A}^+)}{a_{jj \dots j} - r_j^+(\mathcal{A})} \right\} \\ &\quad (r_i^+(\mathcal{A}) - a_{ii_2 \cdots i_m}) + \sum_{\substack{i_2 \cdots i_m \in N_0^{m-1}(\mathcal{A}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} (r_i^+(\mathcal{A}) - a_{ii_2 \cdots i_m}) \\ &\geq \sum_{\substack{i_2 \cdots i_m \in N_L^{m-1}(\mathcal{C}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} (r_i^+(\mathcal{C}) - a_{ii_2 \cdots i_m}) + \sum_{\substack{i_2 \cdots i_m \in N_U^{m-1}(\mathcal{C}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} \max_{j \in \{i_2, \dots, i_m\}} \left\{ \frac{r_j(\mathcal{C}^+)}{a_{jj \dots j} - r_j^+(\mathcal{A})} \right\} \\ &\quad (r_i^+(\mathcal{C}) - a_{ii_2 \cdots i_m}) + \sum_{\substack{i_2 \cdots i_m \in N_0^{m-1}(\mathcal{C}^+) \\ \delta_{ii_2 \cdots i_m} = 0}} (r_i^+(\mathcal{C}) - a_{ii_2 \cdots i_m}) \\ &= p_i(\mathcal{C}^+) \end{aligned}$$

Overall, there is $a_{i \dots i} - r_i^+(\mathcal{C}) > p_i(\mathcal{C}^+)$, and this proof is completed. \square

4. Conclusions. In this paper, SDD_1 tensors were proposed, and it was proved that an SDD_1 tensor is a strong \mathcal{H} -tensor. Meanwhile, we give an application of SDD_1 tensor, which is determining the positive definiteness of even-order real symmetric tensors. Furthermore, the B_1 -tensor was proposed by the SDD_1 tensor, and some properties of B_1 -tensor were introduced. Some examples were given to illustrate those new results.

Availability of data and materials. Not applicable.

Competing interests. The authors declare that they have no competing interests.

Acknowledgments. The authors are grateful to the referee for careful reading of the paper and valuable suggestions and comments.

Authors' contributions. All authors jointly worked on the results, and they read and approved the final manuscript.

REFERENCES

- [1] A. Cichocki, R. Zdunek, A.H. Phan, and S. I. Amari. *Nonnegative Matrix and Tensor Factorizations – Applications to Exploratory Multi-way Data Analysis and Blind Source Separation*, 2009.
- [2] W. Ding, L. Qi, and Y. Wei. M-tensors and nonsingular M-tensors. *Linear Algebra Appl.*, 439:3264–3278, 2013.
- [3] W. Gong, Y. Li, and Y. Wang. An improved iterative algorithm for identifying strong \mathcal{H} -tensors. *Comm. Appl. Math. Comput.*, 1–17, 2024.
- [4] M. R. Kannan, N. Shaked-Monderer, and A. Berman. Some properties of strong \mathcal{H} -tensors and general \mathcal{H} -tensors. *Linear Algebra Appl.*, 476:42–55, 2015.
- [5] E. Kofidis and P.A. Regalia. On the best rank-1 approximation of higher-order supersymmetric tensors. *SIAM J. Matrix Anal. Appl.*, 23:863–884, 2002.
- [6] T.G. Kolda and B.W. Bader. Tensor decompositions and applications. *SIAM Rev.*, 51:455–500, 2009.
- [7] T.G. Kolda and J.R. Mayo. Shifted power method for computing tensor Eigenpairs. *SIAM J. Matrix Anal. Appl.*, 32:1095–1124, 2010.
- [8] C. Li, Y. Li, and X. Kong. New eigenvalue inclusion sets for tensors. *Numer. Linear Algebra Appl.*, 21:39–50, 2014.
- [9] C. Li, F. Wang, J. Zhao, Y. Zhu, and Y. Li. Criteria for the positive definiteness of real supersymmetric tensors. *J. Computat. Appl. Math.*, 255:1–14, 2014.
- [10] G. Li, Y. Zhang, and Y. Feng. Criteria for nonsingular \mathcal{H} -tensors. *J. Adv. Appl. Math.*, 33(3):287–297, 2018.
- [11] Y. Li, Q. Liu, and L. Qi. Programmable criteria for strong \mathcal{H} -tensors. *Numer. Algor.*, 74:199–221, 2017.
- [12] L. Liu and Y. Wang. New criteria for nonsingular \mathcal{H} -tensors. *Chin. Q. J. Math.*, 38:123–133, 2023.
- [13] M. Moakher. On the averaging of symmetric positive-definite tensors. *J. Elastic.*, 82:273–296, 2006.
- [14] M. Ng, L. Qi, and G. Zhou. Finding the largest eigenvalue of a nonnegative tensor. *SIAM J. Matrix Anal. Appl.*, 31:1090–1099, 2009.
- [15] Q. Ni, L. Qi, and F. Wang. An eigenvalue method for testing positive definiteness of a multivariate form. *IEEE Trans. Automat. Contr.*, 53:1096–1107, 2008.
- [16] L. Qi. Eigenvalues of a real supersymmetric tensor. *J. Symb. Computat.*, 40:1302–1324, 2005.
- [17] T. Schultz and H.P. Seidel. Estimating crossing fibers: A tensor decomposition approach. *IEEE Trans. Visual. Comput. Graph.*, 14:1635–1642, 2008.
- [18] Y. Song and L. Qi. Properties of some classes of structured tensors. *J. Optim. Theory Appl.*, 165:854–873, 2015.
- [19] F. Wang and L. Qi. Comments on explicit criterion for the positive definiteness of a general quartic form. *IEEE Trans. Automat. Contr.*, 50:416–418, 2005.
- [20] F. Wang and D. Sun. Criteria for \mathcal{H} -tensors and an application. *J. Numer. Methods Comput. Appl.*, 36:215–224, 2015.
- [21] F. Wang and D. Sun. New criteria for \mathcal{H} -tensors and an application. *J. Inequalities Appl.*, 96:1–12, 2016.
- [22] F. Wang, D. Sun, and Y. Xu. Some criteria for identifying \mathcal{H} -tensors and its applications. *Calcolo*, 56:1–17, 2019.
- [23] F. Wang, D. Sun, J. Zhao, and C. Li. New practical criteria for \mathcal{H} -tensors and its application. *Linear Multilinear Algebra*, 65:269–283, 2017.
- [24] Y. Wang, G. Zhou, and L. Caccetta. Nonsingular \mathcal{H} -tensor and its criteria. *J. Ind. Manag. Optim.*, 12:1173–1186, 2016.
- [25] Y. Xu, R. Zhao, and B. Zheng. Some criteria for identifying strong \mathcal{H} -tensors, *Numer. Algor.*, 80:1121–1141, 2019.
- [26] L. Zhang, L. Qi, and G. Zhou. M-tensors and some applications. *SIAM J. Matrix Anal. Appl.*, 35(2), 2014.
- [27] P. Zhao. New criterions for \mathcal{H} -tensors and its applications. *Adv. Appl. Math.*, 11:5727–5736, 2022.