

# REMOTE SENSING OF VEGETATION RECOVERY IN GRASSLANDS AFTER THE 1988 FIRES IN YELLOWSTONE NATIONAL PARK

EVELYN MERRILL ♦ DEPARTMENT OF ZOOLOGY AND PHYSIOLOGY

RONALD MARRS ♦ DEPARTMENT OF GEOLOGY  
UNIVERSITY OF WYOMING ♦ LARAMIE

## ♦ INTRODUCTION

Traditional methods for measurement of vegetative characteristics can be time-consuming and labor-intensive, especially across large areas. Yet such estimates are necessary to investigate the effects of large scale disturbances on ecosystem components and processes. Because foliage of plants differentially absorbs and reflects energy within the electromagnetic spectrum, one alternative for monitoring vegetation is to use remotely sensed spectral data (Tueller 1989). Spectral indices developed from field radiometric and Landsat data have been used successfully to quantify green leaf area, biomass, and total yields in relatively homogeneous fields for agronomic uses (Shibayama and Akiyama 1989), but have met with variable success in wildland situations (Pearson et al. 1976). Interference from soils (Hardinsky et al. 1984, Huete et al. 1985), weathered litter (Huete and Jackson 1987), and senesced vegetation (Sellers 1985) have diminished the relationship between green vegetation characteristics and various vegetation indices.

In 1987, we found that a linear combination of Landsat Multi-spectral Scanner (MSS) band 7 and the ratio of MSS bands 6 to 4 explained 63% of the variation in green herbaceous phytomass (GHP) in sagebrush-grasslands on ungulate summer range in the northeastern portion of Yellowstone National Park (Merrill et al. 1993). The extensive fires that occurred in the Park in the summer of 1988 provided

an opportunity to determine whether remote sensing could be used to estimate green phytomass in burned areas and to monitor grassland vegetation recovery in the Park after the fires. Remote sensing has previously been used to follow succession of seral stages in pine forests (Jakubauskas et al. 1990) after burning and to monitor plant cover in tundra (Hall et al. 1980) after wildfires.

The objectives of our study were to (1) develop a model for predicting GHP in sagebrush-grassland communities using 1989 and 1990 Landsat TM spectral information and field data on GHP, (2) validate the model by comparing predictions made from it to actual field data collected in 1991, and if successful, (3) compare initial vegetation recovery in burned areas relative to unburned sagebrush-grassland. We chose to use thematic mapper (TM) data rather than MSS data to increase the band options for developing a predictive model.

## ♦ STUDY AREA

The study was conducted in the northeast portion of Yellowstone National Park with major focus on the upper Lamar, Cache and Calfee River drainages and the Mirror Plateau. General descriptions of physiogamy and soils are given by Despain (1990). Elevations range from 1,500 to 3,300 m. Climate of the Park is characterized by long, cold winters and short dry summers, but

climatic patterns within the Park vary considerably (Farnes 1975 in Houston 1982). Mean annual precipitation in Cooke City, located to the northeast of the Park is 67.0 cm (26.8 in) and mean daily temperature in January and July is -10.3° C (13.5° F) and 13.9° C (57.1° F), respectively.

Descriptions of vegetation communities in the park have been given by Despain (1990). Our work focused on the non-forested plant communities within the study area. These included sagebrush *Artemisia tridentata* communities which have an understory of bluebunch wheatgrass *Agropyron spicatum* in dry areas, and Idaho fescue *Festuca idahoensis* on the more mesic sites. Silver sagebrush *Artemisia cana* with an Idaho fescue co-dominant is found on areas associated with high water table such as stream banks and seeps. High elevation grasslands are dominated by Idaho fescue/tufted hairgrass *Deschampsia cespitosa* and tufted hairgrass/sedge *Carex* spp. At intermediate elevations, Idaho fescue/wheatgrass *Agropyron spicatum* and *A. caninum* communities are encountered with the latter dominating in the more mesic sites.

Elk *Cervus elaphus*, mule deer *Odocoileus hemionus*, bison *Bison bison*, moose *Alces alces*, bighorn sheep *Ovis canadensis*, and pronghorn *Antilocapra americana* are the major ungulates in this area (Houston 1982).

## ◆ METHODS

### VEGETATION SAMPLING

Vegetation data were collected in the field from July 25 to August 10 in 1989, July 30 to August 11 in 1990, and July 30 to August 11 in 1991 across two ungulate summer ranges (Norris-Cache/Calfee ridge complex and Mirror Plateau). Each site encompassed at least 0.81 hectares (9 TM pixels) of relatively homogeneous vegetation. At each site, elevation, aspect (degrees), and average slope (%) of the plot were recorded using 1:24,000 topographic maps and the site mapped. Grassland habitat types followed Yellowstone National Park habitat mapping (Despain 1990).

We qualitatively assessed the intensity of burning in the field at each site according to the

following categories: (1) very hot:  $\geq 80$  % of the ground cover and litter consumed; presence of shrubs noted only by trunk stubs; usually heavy ash layer, (2) moderate burn:  $< 80$  % but usually  $> 35$  % of the ground cover and litter consumed; few live shrubs but standing dead shrubs present, (3) light burn:  $< 35$  % ground cover and litter consumed; many live shrubs remaining, (4) no burn.

A double sampling approach was used to estimate biomass of green forbs, green grasses, and standing dead herbaceous vegetation at each site (Eberhardt and Simmons 1987). Percent cover of graminoids, forbs, bareground, rock, moss, lichens, and wood were visually estimated and average heights of plant types (forbs, graminoids, standing herbaceous phytomass) were measured in 30 microplots (0.01 m<sup>2</sup>). An index of plant volume was calculated (canopy cover x plant height) for each of the 30 microplots. Ten of the microplots at each site were clipped to ground level. Vegetation was separated into green graminoids, green forbs, and standing dead herbaceous material. A criterion of  $\geq 25$  % "green" was used to differentiate green from senescent (standing dead) plants. Biomass samples were dried at 70° C for 48 hours and weighed to the nearest 0.1 gm. The ratio of dry plant biomass to plant volume in clipped microplots at a site was used to estimate dry plant biomass in the 20 microplots which were not clipped.

Differences in mean standing dead, green forb, green graminoid, and total (green plus standing dead) biomass at sites that were sampled in all 3 years were tested using a Wilcoxon matched-pairs signed-ranks test. Differences in plant biomass between 3 burn categories (unburned, lightly burned, moderately to severely burned) were tested within years using Kruskal-Wallis one-way analysis of variance, and between two burn categories (unburned to lightly burned, moderately to severely burned) using a Mann-Whitney U test.

### LANDSAT DATA ACQUISITION AND PREPROCESSING

We used TM data from Landsat satellite 5 to quantify spectral characteristics of our study area. TM imagery for 2 August 1989, 13 August 1990, 31 July 1990 of the study area were acquired from EOSAT by the National Park Service. Due to mechanical problems with the receiving station in

Golden California, EOSAT was unable to provide us with data from our projected 6 August 1990 satellite overpass. The closest date to our field sampling (July 30 - August 11) for which imagery was available was 13 August 1990. Data from this overpass was less than ideal because the date of the overpass was outside our sampling window and there were considerable clouds in the scene. As a result, we were unable to obtain spectral values for 6 field plots sampled in 1990.

Digital data were transferred from 9-track computer tape to the Micro-computer Image Processing System (MIPS) for data processing. Data from each scene were georeferenced to 1:24,000 USGS topographic using 8 control points. Environmental conditions that differed among years at the time of the satellites overpasses, such as sun angle and atmospheric conditions, were standardized to 1989 conditions in the following manner. First, we located 6 control sites of 9 TM pixels each in the 3 images, including bright landscape elements (Wahb hot springs and Lamar trail thermal area) and dark landscape elements (Trout and Soda Butte Lakes; rock faces of Abiathar and Thunderer peaks). Second, we recorded the spectral values of the 9 pixels for each spectral band at each site and calculated the average for the site. Third, we estimated the parameters of a linear relationship between average spectral values of each band in 1989 to the other 2 years (Appendix I). Finally, we used the relationships derived from the control points for each band to adjust reflectance values of all pixels in 1990 and 1991 to 1989.

#### RELATIONSHIP BETWEEN GREEN PHYTOMASS AND SPECTRAL VALUES

Values for each of the 6 TM bands were recorded for 9 pixels (0.81 ha) encompassing each field site and averaged to represent the spectral value of the site. Linear combinations of the TM values, as well as published vegetation indices (Jackson 1983), were related to field estimates of biomass at field sites using least squares linear regression. Three indices were based on ratios of the red and near infrared (NIR) TM bands: the ratio vegetation index ( $RVI = NIR/red$ ), the normalized difference index ( $NDVI = (NIR-red)/(NIR+red)$ ) and the transformed vegetation index ( $TVI = \sqrt{ND + 0.05}$ ) (Huete and Jackson 1987). The soil brightness index (SBI), the perpendicular vegetation index

(PVI), and the green vegetation index (GVI) were derived using the Graham-Schmidt orthogonalization process (Jackson 1983) in the MIPS software. Jackson (1983) showed that these indices minimize soil background variations while improving green vegetation signals.

We evaluated the relationship between the Landsat spectral values and the field estimates of biomass in two steps. First, the regressions between vegetative and spectral characteristics were evaluated based on their F value ( $P \leq 0.05$ ), the amount of variation in the dependant variable explained by the independent variables ( $r^2$ ), and the standard error of the estimate. Second, 21 field sites were sampled in more than one year. We used data from only one year to develop relationships between spectral characteristics and green herbaceous biomass. The remaining data were used to "validate" estimates of phytomass predicted from spectral values and actual field data.

## ◆ RESULTS

### FIELD ESTIMATES OF PHYTOMASS

Vegetation was sampled at 62 individual field sites, with 21 sites resampled in all 3 years (Table 1). Plots were distributed about equally among Lamar Flat-Norris Mount, Cache-Calfee Ridge, and the Mirror Plateau. Graminoids consistently averaged about 50% of the total green herbaceous phytomass (GHP) in the 3 years of the study (Table 2). Biomass of green forbs, green graminoids, and total herbaceous biomass (green biomass plus standing dead) on the 21 sites sampled each year was higher in 1990 than in 1989 and 1991 ( $P < 0.05$ ). The proportion of total herbaceous vegetation that was standing dead was lower in 1989 ( $0.04 \pm 0.06$ ,  $x \pm s.d.$ ) than in 1990 ( $0.12 \pm 0.11$ ) and lower in 1990 than in 1991 ( $0.24 \pm 0.17$ ) ( $P \leq 0.01$ ).

There were no significant differences in biomass of green graminoids and forbs between unburned, lightly burned, and moderately to severely burned in any year, but sample sizes within each burn category were low (5 - 8 sites). When sites were combined into severely to moderately burned ( $n = 8$ ) and lightly to unburned ( $n = 13$ ), graminoid biomass was lower but not significantly lower on severely to

Table 1. Location and characteristics of field sites sampled during August of 1989, 1990 and Yellowstone National Park.

Plot #	Year Sampled	Latitude/Longitude	Location	Burn <sup>1</sup>	Elev (m)	Asp (°)	Slp (°)	Habitat Type <sup>2</sup>
101	1989, 1991	44 50 38.6 110 08 32.7	Lower Norris	T3	7520	180	15	TFG
102	1989, 1990, 1991	44 50 51.7 110 09 03.5	Lower Norris	No	7740	179	17	TFG
103	1989, 1990, 1991	44 50 28.3 110 07 49.7	Middle Norris	T1	7800	252	15	TFG
104	1989, 1990, 1991	44 50 13.6 110 08 07.1	Middle Norris	T2	7520	302	9	TFG
105	1989, 1990, 1991	44 49 16.0 110 08 35.0	Lower Cache	T2	7460	250	14	TFG
106	1989, 1990, 1991	44 49 20.2 110 08 25.2	Lower Cache	No	7700	253	6	FN
107	1989, 1990	44 48 21.0 110 05 49.2	Upper Cache	No	8140	0	0	FN
108	1989	44 48 11.1 110 06 40.0	Upper Cache	No	7940	230	1	TFG
109	1989, 1990, 1991	44 48 47.3 110 05 58.6	Upper Cache	T3	8025	2	3	TFG
110	1989, 1990, 1991	44 48 19.1 110 06 24.4	Upper Cache	No	7960	200	18	TFG
111	1989, 1990, 1991	44 48 06.9 110 06 40.0	Upper Cache	No	7850	0	0	TFG
112	1989	44 48 18.2 110 07 19.2	Upper Cache	No	7760	295	4	TFG
113	1989, 1990, 1991	44 48 45.9 110 07 40.9	Upper Cache	T3	7680	211	11	TFG
114	1989, 1990, 1991	44 51 05.0 110 11 03.1	Lamar Flat	T2	6640	0	0	TF
115	1989, 1990, 1991	44 50 59.7 110 11 12.3	Lamar Flat	No	6640	0	0	TFG
116	1989	44 51 05.0 110 11 03.1	Upper Lamar Flat	T2	6710	0	1	TF
121	1989, 1990, 1991	44 48 30.3 110 11 59.7	Opal Creek	T2	8800	127	8	FNG
122	1989, 1990, 1991	44 48 44.3 110 11 51.5	Opal Creek	T2	8740	95	15	FNG
123	1989, 1990	44 48 26.1 110 11 53.5	Opal Creek	T3	8760	101	6	FNG
124	1989, 1990, 1991	44 47 56.8 110 11 00.3	Above Opal Camp	No	8960	90	6	FNG
125	1989, 1990, 1991	44 47 28.0 110 11 22.0	Above Opal Camp	No	8800	170	15	FNG
126	1989, 1990, 1991	44 47 16.1 110 10 55.4	Above Opal Camp	T3	8760	353	4	FNG
127	1989, 1990, 1991	44 48 10.6 110 12 13.6	Opal Creek	T3	8800	15	1	FNG
128	1989, 1990, 1991	44 48 26.7 110 11 44.5	Opal Creek	No	8680	287	8	FNG
129	1989, 1990	44 50 12.7 110 12 10.2	Specimen Ridge Trail	T2	7950	80	7	TFG
130	1989	44 48 25.3 110 13 35.8	Mirror Plateau	No	9120	192	20	FNG
131	1989	44 48 51.5 110 14 11.8	Mirror Plateau	No	9170	0	2	FNG
132	1989	44 49 09.8 110 13 45.2	Top Specimen Ridge Tr	No	8840	150	20	FNG
133	1989, 1990, 1991	44 50 27.7 110 09 46.2	Above Norris Hot Sp	T2	7000	239	7	TFG
134	1989, 1990, 1991	44 50 18.9 110 09 19.5	Lower Norris	T1	7250	213	14	TFG
136	1989	44 50 57.8 110 09 51.4	West Of Norris Cliff	T2	7440	180	15	FA
137	1989	44 51 05.8 110 06 48.6	Upper Norris	No	8130	121	5	TFG
138	1989, 1991	44 51 07.2 110 08 03.5	Pk Midway To Norris	No	8250	171	10	FNG
139	1989, 1990, 1991	44 50 54.2 110 09 06.2	Top/Draw Mid-Norris	No	7860	276	6	TFG
140	1989, 1990	44 50 57.8 110 09 30.7	Norris/Next To Cliff	No	7800	294	7	TFG
141	1990, 1991	44 50 54.3 110 09 57.0	Lower Norris	T2	7440	220	18	FA
142	1990, 1991	44 50 50.2 110 06 55.8	Upper Norris	No	8000	187	14	TFG
143	1990, 1991	44 50 46.6 110 07 56.7	Midway to Norris	No	7700	186	14	FNG
144	1990, 1991	44 50 33.6 110 10 15.0	Lower Norris	T3	6760	261	6	FNG
145	1990	44 48 46.1 110 06 57.0	Upper Cache	No	8140	211	7	FNG
146	1990	44 48 41.6 110 06 48.0	Upper Cache	No	8030	148	12	FNG
147	1990	44 48 35.7 110 07 20.3	Upper Cache	No	7720	296	1	DW
148	1990	44 48 56.1 110 07 37.3	Upper Cache	No	7800	55	2	FN
149	1990	44 48 38.3 110 06 57.5	Upper Cache	No	7920	240	82	TFG
151	1990	44 99 10.1 110 05 57.4	Upper Cache	No	7900	310	18	TFG
153	1990	44 48 15.0 110 06 02.2	Upper Cache	No	7920	0	0	TFG
155	1990	44 48 13.9 110 13 15.4	Above Opal Creek	No	9280	136	5	FN
156	1990	44 48 28.6 110 13 59.2	Mirror Plateau	No	9040	208	28	FNG
157	1990	44 48 53.1 110 14 11.3	Mirror Plateau	No	9200	12	10	FN
200	1991	44 47 51.4 110 11 51.4	Opal Creek	No	8840	200	12	FNG
201	1991	44 48 07.3 110 11 36.7	Mirror Plateau	No	8920	260	3	FNG
202	1991	44 47 23.8 110 11 36.2	Mirror Plateau	No	8820	130	5	FNG
203	1991	44 48 31.3 110 13 24.0	Opal Creek	No	9080	200	8	FNG
204	1991	44 48 08.4 110 13 21.0	Specimen Ridge Trail	No	9360	82	8	FN
205	1991	44 49 08.7 110 14 09.1	Opal Creek	No	9160	210	5	FNG
206	1991	44 49 23.2 110 13 19.0	Specimen Ridge Trail	No	8680	160	18	FNG
210	1991	44 51 43.1 110 10 09.7	Lamar Flat	No	6640	0	0	TF
211	1991	44 50 50.6 110 11 02.0	Lamar Flat	T3	6720	0	0	TF
212	1991	44 51 25.7 110 10 33.7	Lamar Flat	T2	6720	0	0	TF
213	1991	44 48 40.1 110 07 34.1	Cache Calfee Ridge	No	7720	0	0	FNG
214	1991	44 49 08.5 110 07 49.7	Cache Calfee Ridge	No	7960	0	0	FA

<sup>1</sup> Burn rankings: No - unburned; T1 - severe burn; T2 - severe-moderate burn; T3 - moderate burn; T4 - light burn. See text for complete description.

<sup>2</sup> Despain (1990)

moderately burned sites. In contrast, forb biomass was higher in burned areas with a significant difference occurring in 1990 ( $P < 0.05$ ).

## SPECTRAL INDICES AND VEGETATION CHARACTERISTICS

The normalized difference index (NDVI) was the spectral index most highly correlated with

Table 2. Plot characteristics, percent cover of bareground (Bg), rock, litter (Lit), dead wood, moss, dry weight (g/0.01 m<sup>2</sup>) of graminoids, forbs, standing dead herbaceous vegetation (SDHV), values of thematic mapper spectral bands, and spectral indices for field plots sampled in Yellowstone National Park in August 1989 - 1991.

Year	Plot #	Plot Characteristics	Percent Cover						Phytomass (g/0.01 m <sup>2</sup> )					Landsat Spectral Data					Spectral Vegetation Indices							
			Burn	Clay	Asp	Slp	CT	Bg	Rock	Lit	Wood	Moss	Grass	Forbs	SDHV	GSP	THP	TM1	TM2	TM3	TM4	TM5	TM7	NDVI	RV1	TV1
1989	8901	T3 7520 180 15 TFG	30.1	1.3	9.1	3.5	1.7	2.30	5.70	0.00	8.00	8.00	69	29	32	75	101	49	0.40	2.34	1.59	35.0	145.4	3.7		
1989	8902	T0 7720 196 25 TFG	7.6	0.1	11.8	5.9	1.4	3.62	4.00	0.55	7.62	8.17	68	30	32	42	100	41	0.12	1.28	2.86	6.6	122.9	1.8		
1989	8903	T1 7800 254 20 TFG	49.0	4.3	6.3	0.3	0.0	4.09	2.65	0.00	6.74	6.74	64	25	25	50	84	50	0.33	2.00	1.75	15.3	121.4	3.4		
1989	8904	T2 7520 228 10 TFG	53.2	5.4	7.5	0.4	0.0	1.35	2.10	0.00	3.45	3.45	65	26	29	52	88	52	0.28	1.79	1.89	15.7	127.0	2.8		
1989	8905	T2 7460 250 10 TFG	40.5	2.6	9.5	5.3	0.0	4.71	5.70	0.00	10.41	10.41	65	26	28	59	88	42	0.42	2.46	1.35	32.0	130.9	4.1		
1989	8906	T0 7680 235 5 FN	2.4	14.7	72.3	0.0	4.5	0.34	0.00	0.18	0.54	0.72	83	35	45	57	121	65	0.12	1.27	2.92	10.5	166.6	1.7		
1989	8907	T0 8100 230 20 FN	8.6	9.8	69.0	0.0	4.0	0.18	0.52	0.13	0.70	0.83	74	32	43	58	100	55	0.15	1.35	2.60	14.1	148.4	1.8		
1989	8908	T0 7940 230 1 TFG	0.5	0.0	26.3	0.0	0.7	15.17	3.45	0.21	18.62	18.83	63	27	29	90	93	37	0.51	3.10	1.41	51.8	139.8	5.3		
1989	8909	T3 8025 283 5 TFG	40.9	2.3	16.9	2.4	2.4	4.87	3.42	0.02	8.29	8.31	70	29	36	70	96	52	0.32	1.94	1.70	28.2	144.3	2.9		
1989	8910	T0 7960 195 20 TFG	36.0	0.1	43.1	6.4	0.0	0.80	2.42	0.25	3.92	4.17	74	32	43	58	100	55	0.15	1.25	2.60	14.1	148.4	1.8		
1989	8911	T0 7850 184 3 TFG	8.4	0.3	49.9	0.0	0.0	8.34	5.44	0.11	13.78	13.89	67	30	31	95	101	40	0.51	3.06	1.42	54.4	149.9	5.0		
1989	8912	T0 7760 295 4 TFG	2.9	0.0	30.0	0.0	0.0	10.65	5.04	0.47	15.69	16.16	60	29	30	94	99	40	0.52	3.13	1.41	56.1	143.9	5.2		
1989	8913	T3 7750 230 10 TFG	23.8	1.3	40.6	2.1	0.0	4.91	5.58	0.10	10.49	10.59	67	30	30	96	93	38	0.52	3.20	1.40	54.9	145.9	5.4		
1989	8914	T1 6430 0 0 TF	54.3	1.0	0.0	0.0	0.0	0.10	4.32	0.36	12.60	11.78	69	28	31	64	99	51	0.35	2.06	1.72	25.4	139.9	3.2		
1989	8915	T0 7160 206 25 TFG	3.2	0.8	70.2	12.4	1.7	2.57	4.90	0.16	7.47	7.63	76	31	38	62	107	51	0.34	1.63	2.05	19.9	150.7	2.3		
1989	8916	T2 6710 0 1 TF	33.5	0.0	26.8	0.0	0.0	5.53	14.92	0.11	20.45	20.56	74	31	35	88	99	42	0.43	2.51	1.54	44.2	152.8	3.9		
1989	8917	T3 8025 141 5 TFG	36.1	0.5	32.3	0.0	0.2	6.29	3.45	0.37	9.74	10.11	70	30	35	83	104	51	0.41	2.37	1.58	40.8	152.4	3.6		
1989	8918	T2 8740 95 15 FNG	18.4	15.0	33.7	0.0	0.0	4.4	8.0	0.23	12.10	12.13	66	28	31	82	91	40	0.31	1.91	1.59	32.0	142.7	2.8		
1989	8919	T2 8400 95 7 FNG	9.9	0.5	27.7	0.0	2.0	4.70	4.79	0.14	9.49	9.63	63	27	29	69	83	42	0.41	2.38	1.58	31.7	142.8	3.9		
1989	8920	T0 8960 90 7 FNG	17.4	0.1	39.4	0.0	0.0	4.29	3.59	0.25	7.88	8.13	67	30	31	103	101	40	0.54	3.32	1.38	61.4	152.9	5.5		
1989	8921	T0 8680 190 15 FNG	47.5	1.1	18.7	0.0	0.0	4.23	5.25	0.21	9.48	9.69	66	29	30	116	102	39	0.59	3.87	1.32	73.7	156.8	6.6		
1989	8922	T3 8760 156 1 FNG	3.8	0.0	26.6	15.0	0.0	0.0	0.0	0.0	0.0	0.0	67	29	35	82	95	34	0.39	2.00	1.43	53.0	142.5	4.9		
1989	8923	T4 8800 15 1 FNG	0.0	0.1	22.0	0.0	0.0	11.20	1.80	0.15	13.00	13.15	63	29	27	107	89	33	0.60	3.96	1.31	66.9	143.4	6.2		
1989	8924	T0 8660 305 6 FNG	0.6	0.1	26.1	0.0	0.5	10.80	2.30	0.09	13.10	13.19	63	29	27	111	86	31	0.61	4.11	1.30	70.3	143.2	7.5		
1989	8925	T0 8770 90 5 TFG	6.9	0.1	16.1	0.0	0.0	10.40	11.10	0.36	21.50	21.86	69	30	37	84	106	46	0.39	2.27	1.62	42.3	153.0	3.4		
1989	8926	T0 9170 0 2 FNG	5.7	0.0	38.9	0.0	3.3	9.10	4.60	1.35	13.70	15.05	71	32	38	89	107	50	0.40	2.34	1.59	45.0	156.4	3.5		
1989	8927	T0 8840 150 30 FNG	33.5	3.7	25.4	0.0	0.0	6.90	4.00	0.41	10.90	11.31	82	38	49	83	120	64	0.26	1.69	1.98	31.9	177.7	2.3		
1989	8928	T3 8130 23 5 TFG	6.1	0.0	17.0	0.0	0.0	6.70	16.50	0.93	23.20	24.13	68	28	30	64	81	44	0.36	2.13	1.68	24.7	129.6	3.4		
1989	8929	T1 7250 230 12 TFG	55.3	6.1	7.6	0.0	0.0	4.8	8.0	0.04	14.60	14.60	68	28	31	78	91	40	0.28	1.76	1.80	52.0	142.7	3.7		
1989	8930	T2 7440 180 15 FA	33.5	23.4	14.1	0.0	0.0	5.20	5.80	0.27	11.00	11.27	76	32	46	56	89	50	0.14	1.33	2.66	11.3	142.5	1.8		
1989	8931	T0 8130 121 5 TFG	2.3	0.0	21.9	0.0	0.0	14.70	8.30	0.35	23.00	23.35	68	31	33	104	99	41	0.52	3.15	1.41	60.8	154.5	4.1		
1989	8932	T0 8310 121 5 TFG	17.6	29.0	18.4	0.0	0.2	2.20	1.60	0.00	3.80	3.80	76	32	41	66	103	53	0.23	1.61	2.08	21.6	152.6	2.2		
1989	8933	T0 8310 121 5 TFG	4.9	9.3	14.7	0.0	0.0	4.4	8.0	0.04	14.60	14.60	68	28	31	78	91	40	0.28	1.76	1.80	52.0	142.7	3.7		
1989	8934	T0 7880 338 10 TFG	7.8	0.2	20.1	0.0	0.0	15.10	9.60	0.35	24.00	25.05	67	28	28	94	78	30	0.54	3.36	1.38	53.2	154.9	5.9		
1989	8935	T0 7760 302 5 TFG	8.0	2.0	13.0	37.0	0.0	2.20	1.50	0.60	3.70	4.30	74	30	44	65	105	49	0.19	1.48	2.29	32.4	145.2	2.0		
1990	9003	T1 7800 259 15 TFG	45.0	4.0	3.0	1.0	0.0	1.90	10.70	1.20	12.60	13.80	67	27	35	95	77	37	0.17	1.41	2.43	22.3	122.2	1.9		
1990	9004	T2 7520 202 9 TFG	10.0	1.0	9.0	4.0	5.0	13.30	5.10	1.30	18.40	19.70	68	29	39	56	86	44	0.18	1.44	2.37	24.4	127.6	2.0		
1990	9005	T0 7460 140 6 FNG	4.5	0.0	16.0	0.0	0.0	4.50	5.30	1.00	7.80	8.80	67	29	35	82	95	34	0.31	1.96	1.60	42.2	138.5	2.8		
1990	9006	T0 7700 253 6 FN	4.0	1.0	46.0	0.0	10.0	0.40	1.60	2.00	2.00	4.00	82	34	51	63	119	61	0.11	2.24	3.09	28.1	162.3	1.6		
1990	9007	T0 8140 0 0 FN	19.0	36.0	34.0	0.0	5.0	0.20	1.60	0.60	1.80	2.40	74	33	47	59	94	52	0.11	1.26	2.98	22.9	142.1	1.6		
1990	9009	T2 8760 170 3 TFG	31.0	1.0	17.0	1.0	0.0	13.40	6.50	4.90	19.90	24.80	70	31	44	76	96	47	0.27	1.73	1.95	40.1	144.6	2.4		
1990	9010	T0 7960 200 18 TFG	10.0	0.0	26.0	15.0	0.0	0.0	0.0	0.0	0.0	0.0	67	29	35	82	95	34	0.39	2.00	1.43	53.0	142.5	4.9		
1990	9011	T0 7850 0 0 FN	2.0	0.0	8.0	0.0	0.0	18.60	7.70	1.60	26.30	27.90	65	29	36	99	87	31	0.47	2.75	1.48	61.7	140.8	3.4		
1990	9012	T3 7680 211 11 TFG	23.0	3.0	14.0	1.0	0.0	4.70	7.00	0.40	11.70	12.10	68	28	33	85	76	31	0.44	2.58	1.52	48.0	151.2	4.0		
1990	9013	T2 6640 0 0 TFG	39.0	1.0	7.0	1.0	0.0	15.40	8.20	0.00	23.60	23.60	74	31	47	58	105	53	0.10	1.23	3.10	25.3	145.2	1.6		
1990	9014	T2 6640 0 0 TFG	39.0	1.0	7.0	1.0	0.0	15.40	8.20	0.00	23.60	23.60	74	31	47	58	105	53	0.10	1.23	3.10	25.3	145.2	1.6		
1990	9015	T0 8640 0 0 FN	15.0	6.0	18.0	0.0	3.0	7.30	3.70	2.50	11.00	13.50	73	32	45	70	100	44	0.22	1.56	2.16	34.8	145.3	2.1		
1990	9021	T2 8800 127 8 FD	15.0	6.0	18.0	0.0	3.0	7.30	3.70	2.50	11.00	13.50	73	32	45	70	100	44	0.22	1.56	2.16	34.8	145.3	2.1		
1990	9022	T2 8740 95 15 FNG	11.0	5.0	7.0	2.0	0.0	8.90	6.20	0.70	15.10	15.80	65	29	36	130	97	34	0.57	3.61	1.35	89.4	157.3	5.7		
1990	9023	T4 8760 101 6 FN	23.0	1.0	15.0	0.0	0.0	15.70	5.80	2.70	21.50	24.20	67	27	35	95	77	37	0.17	1.41	2.43	22.3	122.2	1.9		
1990	9024	T0 8960 90 7 FNG	11.0	5.0	7.0	2.0	0.0	8.90	6.20	0.70	15.10	15.80	65	29	36	130	97	34	0.57	3.61	1.35	89.4	157.3	5.7		
1990	9025	T0 8800 170 15 FNG	25.0	1.0	18.0	0.0	0.0	6.60	6.40	2.60	13.00	15.60	65	29	36	130	97	34	0.57	3.61	1.35	89.4	157.3	5.7		
1990	9026	T3 8760 353 4																								

years and no simple or multiple regression model could be found that explained more than 40% of the variation in NDVI in all years. In particular, data from 1990 had higher biomass for the same NDVI values as in other years. Because of differences in the timing of field sampling and satellite overpass in 1990, we did not use data collected in 1990 to develop our relationship between spectral NDVI and field estimates of phytomass. TVI and TM band 7 were most highly correlated with standing dead phytomass.

Table 3. Significant ( $P \leq 0.05$ ) correlation coefficients between TM spectral bands and spectral indices and vegetation characteristics measured during late summer 1989 - 1991 in Yellowstone National Park. NS indicates not significant.

Vegetative Characteristic	Year	Spectral Index	K
Total standing biomass	1989	NDVI <sup>1</sup>	0.53
	1990	NDVI	0.45
	1991	NDVI	0.72
Green herbaceous phytomass	1989	NDVI	0.53
	1990	NDVI	0.46
	1991	NDVI	0.74
Green graminoids	1989	NDVI	0.66
	1990	NDVI	0.48
	1991	NDVI	0.74
Green forbs	1989		NS
	1990		NS
	1991		NS
Standing dead herbaceous phytomass	1989		NS
	1990		NS
	1991		NS
Percent standing dead of total standing phytomass	1989	TVI <sup>2</sup> , TM7 <sup>3</sup>	0.61, 0.52
	1990	TM7	0.35
	1991	TVI, TM7	0.52, 0.42

<sup>1</sup>NDVI = normalized difference vegetation index: (TM Band 4 - TM Band 3)/(TM Band 4 + TM Band 3).  
<sup>2</sup>TVI = transformed vegetation index: SQRT(NDVI+0.05)  
<sup>3</sup>TM7 = thematic mapper spectral band 7

THP and GHP alone explained 45 and 46% ( $P < 0.001$ ) of the variation, respectively, in NDVI at field sites sampled in either 1989 or 1991. Elevation explained an additional 6% of the variation in NDVI (Table 4). Neither burn intensity nor standing dead herbaceous phytomass (SDHP) or the proportion of standing dead of THP explained additional variation in these data once the effects of elevation were accounted for. In contrast, the proportion of THP that was dead explained a significant amount of additional variation in NDVI if elevation were not included in the model. Burn intensity explained a significant amount of the variation in NDVI when combined with THP but not when combined with green grass GG or GHP (Table 4). Average percent canopy cover of other site characteristics that we measured, such as litter or bareground, did not contribute significantly to explaining additional

variation in NDVI.

Table 4. Significant ( $P < 0.001$ ) linear regression models that predict  $\geq 50\%$  of the variation in NDVI at 50 field sites sampled in 1989 and 1991 in Yellowstone National Park.

Independent variables	Coefficients	T	P	r <sup>2</sup>
Constant	0.272	5.79	0.00	0.56
Total biomass	0.013	4.40	0.00	
Proportion standing dead	-0.301	-3.07	0.00	
Burn intensity	-0.071	-2.10	0.04	
Constant	0.321	8.41	0.00	0.55
Green grass	0.016	4.31	0.00	
Proportion standing dead	-0.329	-3.32	0.00	
Burn intensity	-0.057	-1.62	0.11	
Constant	-0.113	-0.65	0.518	0.53
Green herbaceous phytomass	0.016	6.14	0.000	
Elevation	0.0005	1.78	0.082	
Burn intensity	-0.029	-0.78	0.438	
Constant	-0.202	-1.29	0.20	0.52
Green herbaceous phytomass	0.017	6.07	0.00	
Elevation	0.0005	2.46	0.02	
Constant	-0.216	-1.37	0.18	0.51
Total phytomass	0.016	6.02	0.00	
Elevation	0.0005	2.41	0.02	

When the linear relationship was inverted to predict GHP, less than 50% of the variation in THP and GHP was explained by NDVI and elevation. The relationship appeared weak because many high-elevation fields sites with high NDVI values had low GHP. As a result, we stratified sites by elevation and found that following linear models explained 55% of the variation in TGP and GHP at low elevational ( $\leq 2620$  m) sites:

$$\text{THP (g/0.1 m}^2\text{)} = 29.25 \times \text{NDVI} - 1.191 \quad (P < 0.001, \text{ s.e.} = 4.61) \quad \text{Eq. 1}$$

$$\text{GHP (g/0.1 m}^2\text{)} = 31.2 \times \text{NDVI} - 0.501 \quad (P < 0.001, \text{ s.e.} = 4.54) \quad \text{Eq. 2}$$

Using an exponential model, NDVI explained only an additional 1% of the variation in either THP or GHP.

The relationship between NDVI and phytomass at high elevation was nonlinear and the following curves were used to describe the relationships:

$$\text{THP (g/0.1 m}^2\text{)} = \frac{19 \times (\text{NDVI} - 0.18)}{0.110 + (\text{NDVI} - 0.18)} \quad \text{Eq. 3}$$

$$\text{GHP (g/0.1 m}^2\text{)} = \frac{17 \times (\text{NDVI} - 0.18)}{0.102 + (\text{NDVI} - 0.18)} \quad \text{Eq. 4}$$

The above equations were used to predict the

THP and GHP of 16 low elevation sites and 7 high elevation sites that were not used to develop the above predictive equations. On average, GHP was underestimated at low elevations by 0.93 g/m<sup>2</sup> and THP overestimated by 1.01 g/0.01 m<sup>2</sup>. At high elevations, GHP was overestimated by 2.34 g/0.01 m<sup>2</sup> and THP by 2.73 g/0.01 m<sup>2</sup>. Mean percent error in estimates of GHP (37%) at low elevations was greater than at high elevations (24%) because phytomass was generally lower at low elevation sites than at high elevation sites.

#### ◆ FUTURE ANALYSES

We intend to use the biomass-spectral relationships to characterize herbaceous vegetative recovery in the sagebrush-grassland areas in our study area. Our approach is to import TM spectral bands 3 and 4 from Landsat imagery for 1989, 1990, 1991 for our study area and to geo-reference the coverages to Yellowstone National Park's burned area and elevation coverages in ARC/INFO GRID. Next, we will stratify the area by elevation and identify nonforested areas of the study using the YNP habitat type map. Within each elevational stratum, we will use the NDVI-biomass algorithm developed for that elevation to predict the green and total biomass on a pixel-by-pixel basis and calculated the average for both burned and unburned areas. Initial attempts to complete this analysis were thwarted by what we believed to be either major geo-referencing problems or discrepancies between the burned areas and habitat types and what was evident on the Landsat imagery. Once these corrections have been made we can complete our analyses.

#### ◆ ACKNOWLEDGEMENTS

I would like to thank Cathy Wilson for her assistance in field work and initial data analysis. John Baldwin, Jon Hak, Robin Reid, Verda and Mike Shingleton, Greg McDaniel, and Bob Brennen provided help in collecting field data and processing samples. We also thank Rich Kluckus, Don Despain, Francis Singer, John Varley, Jennifer Whipple, George McKay, and Jack Norland of Yellowstone National Park for their administrative and field support. Drs. Kenneth Gerow and Jeff Pontius of the University of Wyoming provided statistical advice.

#### ◆ LITERATURE CITED

- Despain, D. G. 1990. *Yellowstone Vegetation: consequences of environment and history in a natural setting*. Roberts Rinehart. 239 pp.
- Eberhardt, L. L. and M. A. Simmons. 1987. Calibrating population indices by double sampling. *J. Wildl. Manage.* 51:665-675.
- Hall, D. K. J. P. Ormsby, L. Johnson, and J. Brown. 1980. Landsat digital analysis at Kokolik River, Alaska. *Remote Sens. Environ.* 10:263-272.
- Hardinsky, M.A., F. C. Daiber, C.T. Roman, and V. Klemas. 1984. Remote sensing of biomass and annual net aerial primary productivity of a salt marsh. *Remote Sens. of Environ.* 16:91-106.
- Houston, D. B. 1982. *The Northern Yellowstone elk: ecology and management*. Macmillian Publishing, New York. 474 pp.
- Huete, A. R. and R.D. Jackson. 1987. Suitability of spectral indices for evaluating vegetation characteristics on arid rangelands. *Remote Sens. of Environ.* 23:213-232.
- Huete, A. R., R. D. Jackson, D.F. Post. 1985. Spectral response of a plant canopy with different soil backgrounds. *Remote Sens. of Environ.* 17:37-53.
- Jackson, R. D. 1983. Spectral indices in n-Space. *Remote Sens. of Environ* 13: 409-421.
- Jakubauskas, M., K. P. Lulla and P. W. Mausel. 1990. Assessment of vegetation change in a fire-altered forest landscape. *Photogr. Eng. Remote Sens.* 56:371-377.
- Merrill, E. H., M. K., Bramble-Brodahl, R. W. Marrs, M. S. Boyce. 1993. Estimation of green herbaceous phytomass from Landsat MSS data in Yellowstone National Park. *J. Range Manage.* 46:151-157.

Pearson, R.L., C. J. Tucker, and L.D. Miller. 1976. Spectral mapping of short grass prairie biomass. *Photogramm. Eng. Remote Sens.* 42:317-3245.

Shibayama, M. and T. Akiyama. 1989. Seasonal visible, near-infrared and mid-infrared spectra of rice canopies in relation to LAI and above-ground dry phytomass. *Remote Sens. of Environ.* 27:119-127.

Sellers, P.J. 1985. Canopy reflectance, photosynthesis and transpiration. *Int. J. Remote Sens.* 6:1335-1372.

Tueller, P. T. 1989. Remote sensing technology for rangeland management applications. *J. Range Manage.* 42: 442-453. *Remote Sens. of Environ.* 8:127-150.

Appendix I. Equations used to calibrate spectral values from Landsat imagery for Yellowstone National Park on 13 August 1990 and 31 July 1991 to August 2, 1989 for 6 thematic mapper (TM) spectral bands.

Year	TM Band	a	b	s.e.	$r^2$
1990 to 1989	1	-10.90	1.174	3.05	0.99
	2	- 4.14	1.097	1.25	0.99
	3	0.31	1.081	16.94	0.86
	4	- 6.81	1.162	3.89	0.99
	5	- 0.27	1.018	4.51	0.99
	7	- 1.54	1.146	3.54	0.98
	1991 to 1989	1	2.53	0.962	1.39
2		- 0.87	0.955	0.82	0.99
3		4.96	0.906	17.14	0.88
4		- 2.47	1.043	3.42	0.99
5		3.32	0.901	1.19	0.99
7		2.64	0.935	5.85	0.96